1

2

3

4

5

# Original Research Article

## Estimation of Sedimentary thickness Using Spectral Analysis of Aeromagnetic Data over part of Bornu Basin, Northeast, Nigeria

### 6 ABSTRACT

7 The aeromagnetic data (High resolution) over part of Bornu Basin NE was interpreted using spectral analysis to estimate the sedimentary thickness of the study area. The study area is 8 covered by four aeromagnetic data sheets covering latitude  $12^{\circ}N$  and  $13^{\circ}N$  and Longitude 9 12°E and 13°E with an estimated total area of 12, 100 km<sup>2</sup>. Polynomial fitting method was 10 adopted for the regional-residual separation of the total magnetic intensity. The residual map 11 12 was divided into nine spectral sections. The result of the study shows a sedimentary thickness that ranges between 0.54 km and 3.35 km. The sedimentary thickness of over 3 km could be 13 14 found around the South-eastern part of the study area which corresponds to Gubio town while the minimum sedimentary thickness could be found around North-western part of the 15 16 study area which also corresponds to Borgo town area. The maximum sedimentary thickness of 3.35 km may be sufficient for hydrocarbon presence. 17

18

Keywords: Aeromagnetic data, Spectral analysis, Polynomial fitting, Sedimentary thicknessand Hydrocarbon maturation

#### 21 **1. INTRODUCTION**

22 Over the years, searching for earth embedded minerals and hydrocarbon (oil and gas) has 23 become a bone of contention in the economy of Nigeria. The bigger part of the nation's 24 income (80%) is gotten from hydrocarbon (oil and gas) profit whereupon more than 160 25 million overflowing populace relies on upon. As the hydrocarbon potential of the prolific 26 Niger delta becomes depleted or exhausted in the nearest future due to continuous 27 exploitation and inherent crises in the Niger delta region which had led to excessive reduction 28 in oil production for export and domestic use, it is of necessity to shift attention to other 29 sedimentary Basins. In particular, the Bornu Basin which is one of the inland Basins in Nigeria presumed to have high hydrocarbon potential aside other earth minerals with high 30 economic values <sup>20</sup>. The Nigerian government have directed one of its Oil-regulating Agency 31 32 (NNPC) to heighten hydrocarbon prospecting in the Northeast (Bornu Basin). As this will 33 expand Nigeria's oil and gas holds, add qualities to the hydrocarbon potentials of the Nigerian inland basins, in this way, give speculation open doors, help the economy of the nation and 34 35 also provide many new employments to lessen unemployment definitely in Nigeria. This

research will be very useful on a reconnaissance basis for oil and mineral prospecting in thearea.

38 Spectral analysis of aeromagnetic data over the area would be used to determine the 39 sedimentary thickness of the study area. The result from the spectral analysis could be used to 40 suggest areas of potential hydrocarbon presence.

41

#### 42 2. GEOLOGY OF THE STUDY AREA

43 The area of study is part of the Nigerian sector of the Chad Basin, known locally as the Bornu 44 Basin (Figure 1). It is one of the Nigerian inland Basins occupying the north-eastern part of 45 the country. It represents about one-tenth of the total area extent of the Chad Basin, which is 46 a regional large structural depression common to five countries, namely, Cameroon, Central African Republic, Niger, Chad, and Nigeria<sup>14</sup>. The Bornu Basin (Study area) falls between 47 48 latitudes 12°N and 13°N and longitudes 12°E and 13°E with an estimated area of 12,100 km<sup>2</sup>. 49 The Borno State where the area of study is located is endowed with rock mineral base resources such as clay, salt, limestone, kaolin, iron ore, uranium and mica. 50

51 Geologically, the Bornu basin has been explained as a broad sediment-filled broad depression 52 straddling North-eastern Nigeria and adjoining parts of the Republic of Chad. The 53 sedimentary rocks of the area have a cumulative thickness of over 3.6 km and rocks consist 54 of thick basal continental sequence overlaid by transitional beds followed by a thick succession of Quaternary Limnic, fluviatile and eolian sand and clays<sup>7</sup>. The stratigraphy 55 56 sequence (Figure 2) shows that Chad, Kerri-Kerri and Gombe Formations have an average 57 thickness of 130 to 400 m. Below this formations are the Fika shale with a dark grey to black 58 in colour, with an average thickness of 430 m. Others are Gongila and Bima Formations with 59 an average thickness of 320 m and 3,500 m, respectively <sup>15</sup>.



65

66

### 67 **3. MATERIALS AND METHOD**

The study area is covered by four aeromagnetic data sheets of total-field intensity in  $1/2^{\circ}$  by 1/2° procured from Nigerian Geological Survey agency (NGSA). The sheets are 43 (Borgo), 44 (Bazabure), 65 (Chungulbulturi) and 66 (Gubio). The sheets were merged together to

generate the study area using Oasis montaj software. The aeromagnetic data were obtained as part of the aeromagnetic survey carried out between 2003 and 2009 sponsored by Geological survey of Nigeria. The data were obtained at an altitude of 100 m along a flight line spacing of 500 m oriented in NW-SE and a tie line spacing of 2000 m. The maps are on a scale of 1:100,000 and half-degree sheets contoured mostly at 10nT intervals. The geomagnetic gradient was removed from the data using the International geomagnetic Reference Field (IGRF). The total area covered was about 12,100 km<sup>2</sup>.

78 The actual magnetic intensity value was reduced by 33,000 nT for handling before the 79 contour map was plotted. As a result, 33,000 nT must be added to the data so as to get the 80 actual magnetic intensity at a given point. The total magnetic intensity was contoured and 81 colour filled to show the high and low total magnetic intensity of the study area (Figure 3). 82 Polynomial fitting (order one) method was used for the regional-residual separation, Figures 83 4 and 5 shows the contoured and colour filled regional map and residual map respectively. 84 The magnetic values found on figure 4 trend northeast-southwest and the lines observed 85 indicate faults. Since minerals are structurally controlled, minerals present in the study area 86 like gypsum can be found along the fault lines.

#### 87 **3.1.Spectral Analysis**

The statistical spectral analysis of the residual field data was used to determine the depths to the buried magnetic sources within the subsurface of the study area. Spector and Grant developed a 2-D spectral depth determination method. Their model assumes that an uncorrelated distribution of magnetic sources exists at a number of depth intervals in a geologic column. The Fourier transform of the potential filed due to a prismatic body has a broad spectrum whose peak location is a function of the depth to the top and bottom surfaces and whose amplitude is determined by its density or magnetization<sup>21</sup>.

95 The peak wave number ((a) can be related to the geometry of the body according to the96 following expression.

97 
$$W' = \frac{ln \binom{h_b}{h_t}}{h_b - h_t}$$
 (1)

98 Where W' is the peak wave number in radian / ground – unit,  $h_t$  the depth to the top and  $h_b$ 99 is the depth to the bottom.

$$100 \quad f(\mathbf{\omega}) = e^{-h\omega} \tag{2}$$

101 Where  $\varpi$  is the angular wave number in radians/ground-unit and h is the depth to the top of 102 the prism. For a prism with top and bottom surface, the spectrum is:

103 
$$f(\omega) = e^{-h_t \omega} - e^{-h_b \omega}$$
(3)

104 Where  $h_t$  and  $h_b$  are the depths to top and bottom surface respectively. As the prism bottom 105 moves closer to the observation point at surface, the peak moves to a higher wave number. 106 When looking at the spectrum, it is important to note that the amplitude of a deep prism does 107 not exceed the amplitude of the same prism at shallow depth at any wavenumber. The effect 108 of increasing the depth is to shift the peak to lower wavenumbers.

Because of this characteristic, there is no way to separate the effect of deep sources from shallow sources of the same type by using wavenumber filters. The sources can only be distinguished if the deep sources have greater amplitude or if the shallow sources have less depth extent. When considering a line that is long enough to include many sources, the log spectrum of this data can be used to determine the depth to the top of a statistical ensemble of sources using the relationship.

$$115 \quad Log \ E(k) = 4\pi hk \tag{4}$$

116 Where *h* is the depth in ground – units and k is the wavenumber in cycles / ground – unit. The 117 depth of an 'ensemble' of source can be determined by measuring the slope of the energy 118 (power) spectrum and dividing by  $4\pi$ . A typical energy spectrum for magnetic data may 119 exhibit three parts – a deep source component, a shallow source component and a noise 120 component.

In this study, the graph of each energy spectral was obtained with Matlab software purposely designed to accept the longitude and latitude values alongside with its respective magnetic values for each of the nine spectral sections label (A –I) where the log of spectral energy plotted against frequency.



125

Figure 3: Total Magnetic Intensity Map over part of Bornu Basin. Magnetic 'Lows' are represented
by 'L' and magnetic 'Highs' are represented by 'H'. AA<sup>1</sup> and CC<sup>1</sup> are the identified paleo-structures
in the study area



129

130

131

132

Figure 5: Residual Map of the Study Area

### 1334. RESULTS AND DISCUSSION

Figure 4: Regional Map of the Study Area

The residual map (Figure 4) produced from this study was divided into nine (9) blocks (A-I)
of overlapping magnetic sections. Six of the divisions (section SPTA, SPTB, SPTC, SPTD,
SPTE, and SPTF) covered 55km<sup>2</sup> and three others (SPTH, SPTG, and SPTI) covered
110km<sup>2</sup>. The divisions of residual map into nine (9) spectral sections was done with Oasis

Montaj and the spectral energies were plotted, the \*.SPC file obtained were later exported into Microsoft Excel worksheets one after the other until the total number of nine (9) spectral (\*.SPC) energy files were later used as an input file into a spectral program plot (SPP) developed with Matlab. The total numbers of nine (9) spectral energies were plotted in Matlab with the developed program. A typical plot of energy against frequency (wavenumber) is shown in Figure 6.

![](_page_6_Figure_2.jpeg)

![](_page_6_Figure_3.jpeg)

145 The program has been designed to determine the first slope  $(m_1)$  and the second slope to 146 calculate for the first and second magnetic depth source using the equations below.

147  $Z_1 = -\frac{m_1}{4\pi}$  (5)

148 
$$Z_2 = -\frac{m_2}{4\pi}$$
 (6)

Where  $m_1$  and  $m_2$  are slopes of the first and second segment of the plot while  $Z_1$  and  $Z_2$  are first and second depths respectively (Table 1).

- 151 Substituting for  $m_1 = -42.2$  and  $m_2 = -6.82$  (Figure 6) into the equation (5 & 6)
- 152 Therefore,
- 153  $Z_1 = -(-42.2/4 * 3.142) = 3.35$  Km
- 154 Similarly,

### 155 $Z_2 = -(-6.82/4 * 3.124) = 0.54 \text{ Km}$

Figure 6 shows a typical plot of energy spectrum against frequency of section SPTG, two layers can be observed with their respective depths and magnitude values. The minimum depth is 0.54 km while the maximum depth source is 3.35 km (Figure 6). Table 1 shows that the shallow depth source ( $Z_2$ ) ranges from 0.39 km to 0.64 km while the deeper depth source ( $Z_1$ ) ranges from 1.32km to 3.35 km. Figure 7 and 8 shows that the maximum sedimentary thickness of 3.35 km is more pronounced around Gubio town.

S/N	Sections	Longitude	Latitude	Depth	Depth Z2
		(Degrees)	(Degrees)	<b>Z1(km)</b>	(km)
1	SPTA	12.15	12.65	1.32	0.39
2	SPTB	12.65	12.65	1.54	0.53
3	SPTC	12.15	12.15	2.10	0.54
4	SPTD	12.65	12.15	2.77	0.47
5	SPTE	12.5	12.65	1.48	0.29
6	SPTF	12.5	12.15	3.32	0.35
7	SPTG	12.15	12.5	3.35	0.54
8	SPTH	12.65	12.5	3.06	0.46
9	SPTI	12.5	12.5	3.12	0.64

162 Table 1: Estimated depth to deeper magnetic and shallow source in Km

163

164

![](_page_8_Figure_1.jpeg)

Figure 7: Contour Map of Spectral magnetic depth to top of basement over part of Bornu
 Basin.

168

169

165

![](_page_8_Figure_4.jpeg)

![](_page_8_Figure_5.jpeg)

171 5. CONCLUSION

172 Since the nearness and magnitude of minerals, oil and gas must be learned by geophysical investigations of the subsurface geologic structures and the investigation of sedimentary 173 174 basins for hydrocarbon potential relies upon the sedimentary thickness of the basin for 175 hydrocarbon maturation, it can therefore be concluded that the results from the depth 176 estimation techniques used in this study, spectral analysis with highest sedimentary thickness 177 of 3.35 km is sufficient enough for hydrocarbon maturation and the results also concur with 178 those obtained by other researchers in some part of the Bornu Basin; notably are results of studies by  $^2$  and  $^{17}$ . 179

180 181

### 6. **REFERENCES**

- 182
- 1. Afolabi, O. (2011): Interpretation of the Gravity and Magnetic Data of the Chad Basin, Northeast Nigeria for hydrocarbon prospectivity. Unpublished PhD thesis. 183 184 O.A.U. Ile-Ife.
- 2. Anakwuba, E.K., Onwumesi, A.G., Chinwuko, A.I., and Onuba, L.N. (2011). The 185 186 interpretation of aeromagnetic anomalies over Maiduguri-Dikwa depression, Chad Basin, Nigeria: A Structural Review. Archive of Applied Science Research, 3(3), 187 188 1757-1766Ajakaiye, D. E and Burke K. C (the initials) (1973). Bouguer gravity map 189 of Nigeria, Tectonophysics 16, 103-115.
- 3. Bonde, D. S., Udensi, E. E. and Momoh M. (2014). Delineation of magnetic zones of 190 191 Sokoto Basin in Northwestern Nigeria, using Aeromagnetic Data. International 192 Journal of Engineering and Science, vol.4 pp37-45.
- 193 4. Fedi, M. & Rapolla, A. (1997): Space-Frequency analysis and reduction of potential 194 field Ambiguity. Annali Di Geofisica. XL(5).
- 195 5. Isogun, M. A. (2005): Quantitative interpretation of aeromagnetic data of Chad Basin, 196 Bornu State, Nigeria. Unpublished. M.sc. Thesis, O.A.U Ile- Ife.
- 197 6. Lawal, K. M, Umego, M. N. and Ojo, S. B. (2007): Depth-to-Basement Mapping 198 using Fractal Technique: Application to the Chad Basin Northeastern Nigeria. 199 Nigerian Journal of Physics, 19(1).
- 200 7. Lawal, T.O and Nwankwo L.I, wavelet analysis of high resolution aeromagnetic data 201 over part of Chad Basin, Nigeria. *Ilorin Journal of Science*, 2014(1) pp 110-120.
- Philip, K., Brooks. M., and Hill, L. (2002): An Introduction to Geophysical 202 8. 203 Exploration, Blackwell Science, London, UK, 3rd edition.
- 204 9. Mekonnen, T. K. (2004). Interpretation & Geodatabase of Dykes Using Aeromagnetic Data of Zimbabuwe and Mozambique. M.Sc. thesis. International Institute for Geo-205 206 information science and Earth observation, Netherland. pp. 1-2.
- 207 10. Nabighian, M. N., (1972), the analytic signal of two-dimensional magnetic bodies 208 with polygonal cross-section -Its properties and use for automated anomaly interpretation: Geophysics, 37, 507 517. 209
- 11. Nabighian, M. N. (1984): Toward a three-dimensional automatic interpretation of 210 211 potential field data via generalized Hilbert transforms - Fundamental relations: 780-786. 212 Geophysics, 49,

213	12. Nwankwo, L. I., Olasehinde, P. I. & Akoshile, C.O. (2011): A new technique for
214	<i>Journal of Physics</i> 22 (1) 74-79
215	13 Nwankwo C N Ekine A S Nwosu L I (2009) Estimation of heat flow variation
217	in the Chad Basin Nigeria. Journal Applied Sciences and Environment, 13(1), 73 –
218	80.
219	<b>14.</b> Obaje, N.G., Geology and Mineral of Nigeria (Germany: Springer Verlag, 2009)
220	15. Obaje, N, Wehner H., Scheeder G., Abubakar M., and Jauro A., Hydrocarbon
221	prospectively of Nigeria's inland basins: from the viewpoint of organic geochemistry
222	and organic petrology. AAPG bulletin, 88(3), 2004, 325 - 353.
223	16. Odebode, M.O. (2010): A handout on Geology of Boron (Chad) Basin, Northeastern
224	Nigeria.
225	17. Okonkwo, C. C., Onwuemesi, A. G., Anakwuba, E. K., Chinwuko, A. I., Ikumbur, B.
226	E. & Usman, A. O. (2012): Aeromagnetic Interpretation over Maiduguri and Environs
227	of Southern Chad Basin, Nigeria. Journal of Earth Sciences and Geotechnical
228	Engineering, <b>2</b> (3), 77-93.
229	18. Okosun, E. A. (1995). Review of Geology of Bornu Basin. Journal of mining and
230	<i>Geology</i> , 31(2), 113 – 172.
231	19. Oteze, G. E, Foyose, E. A. (1988) Regional developments in the hydrogeology of the
232	Chad Basin. Water Res 1:9–29.
233	20. Salako, (2014), Depth to basement determination using Source parameter imaging
234	(SPI) of aeromagnetic data: An application to upper Benue trough and Borno basin,
235	Northeast, Nigeria. Academic Research International, vol. 5(3).
236	21. Salako K. A and Udensi E. E (2013), Spectral Depth Analysis of part of upper Benue
237	Trough and Bornu Basin, Northeast Nigeria, using Aeromagnetic data. International
238	Journal of Science and Research, Vol. 2(8).
239	22. Spector, A., Grant, F.S., (1970). Statistical models for interpretation of aeromagnetic
240	data. Geophysics, 35, 293–302.
241	23. Sunday, J. (2012). Estimation of Magnetic Source depths from Aeromagnetic data of
242	Maiduguri using gradient Inversion, M.Sc Thesis, Department of Physics, University
243	of Ilorin.
244	24. Summonu, L. (1988). Interpretation of Data from Middle Niger Basin, Nigeria. M.Sc.
245	Thesis. University of Ibadan.
246	25. Teitord, W.M, L.P. Geldart, R.E. Sheriff, (1990) Applied Geophysics, 2 <sup>nd</sup> ed.,
247	Cambridge University Press, Cambridge, 1990, p.770.