

# Spontaneous avulsion of the tibial tubercle following Osgood-Schlatter lesion.in an adult

## Abstract

Osgood Schlatter disease occurs in adolescent age group with male preponderance and persons indulging in active sports. Avulsion of tibial tuberosity with Osgood-Schlatter disease is rare but and occurs in adolescent and younger age. We describe a case of an avulsion of tibial tuberosity in a 49 year old male with no prior symptoms of this disease and was managed with excision of the bony fragment.

**Keywords:** Osgood-Schlatter, Spontaneous, Avulsion, Tibial tubercle, Operation

## Introduction

Osgood- Schlatters' disease (OSD) is characterized by pain, swelling, and tenderness in the anterior aspect of the proximal tibia. In boys, the condition appears between the ages of ten and fifteen years and especially affects those actively participating in sports. Surgical treatment of OSD, albeit rare, may occasionally be warranted if disabling symptoms persist (1 - 4). In the previous literature, excision of the ossicle(s), with or without resection of the tibial tubercle prominence, has been shown to yield better results than other methods (2, 3) (drilling, bone grafting, or refixation of the ossicle). Binazzi et al (5) examined the results of twenty-six knees, in both skeletally immature and mature patients, at an average of thirteen years after surgical treatment. They found that excision of the ossicles, with or without removal of the prominent tibial tubercle, clearly yielded better results than did various other procedures. Spontaneous avulsion of the tibial tuberosity after skeletal maturity following OSD is an extremely rare condition (6). We describe the first case of spontaneous avulsion of the tibial tuberosity following Osgood-Schlatters' disease in an adult.

## Case report

A 49 year old man with no medical history suffered spontaneous sharp pain of the left knee and consecutive swelling during walking on even ground. He had no early or remote history of trauma or unaccustomed heavy work. Physical examination

revealed swelling and tenderness in the region of the tibial tuberosity. No effusion could be palpated, collateral and cruciate ligaments were stable. Range of motion was limited to Extension / Flexion 0/0/40°. The stretched left leg could be lifted actively. Lateral x- ray of left knee (Fig.1) and MRI (Fig.2) showed avulsion of the tibial tuberosity of an unresolved OSD. After simple excision of the ossicle and uneventful postoperative course and partial weight bearing of 20 kg using two crutches for 4 weeks the patient was out of any complaints at latest follow-up after 24 months. Lateral X-ray revealed (Fig.3) the status after excision of the unresolved Osgood-Schlatter. At final follow up there was no local tenderness, range of motion of the left knee was Extension / Flexion 5/0/140°. The stretched left leg could be raised easily. During walking the patient showed no limping.

44

## 45 Discussion

Osgood-Schlatters' disease (OSD) was firstly described approximately more than 100 years ago, chronic avulsion injury caused by repetitive microtrauma and traction of tibial tuberosity (7). It almost occurs in adolescents performing sports activities and can be bilateral in 50 % of cases (8). Although various theories exist, most recently focus has been put on soft tissue component with suggestion that insult occurs at the anterior aspect of tibial tubercle at insertion of patellar tendon (9) in contrast to the general accepted theory of Ogden and Southwick who described avulsion of secondary ossification centre of the tibial tuberosity (10). The radiographs may be either normal or show fragmentation of tibial tuberosity with ossicles. Although conservative treatment is the main stay and symptoms usually subside with rest, restriction of sports in majority of patients, surgery is advisable when conservative treatment fails (11). Pihlajamäki HK et al (1) in a large retrospective study of 107 patients managed surgically by excision of the ossicle(s) with or without resection of prominent tibial tuberosity found good long term results with no deleterious results. The median age of surgery in their group was 20 (Range 18-29) years with duration of symptoms ranging from 1 - 14 years. The main indication of surgery in their group was persistent symptoms despite conservative treatment.

Avulsion fractures of tibial tuberosity occur after forceful physical activity or violent trauma (12) such as fall from height is rare but has been reported even bilaterally in

few cases (13). The usual age group of occurrence is adolescence and usually patients have been found to indulge in strenuous activities despite medical advice on the contrary. All the previous cases reported were symptomatic and were diagnosed previously except one case of asymptomatic bilateral avulsion with preexisting OSD reported by Ogden et al (14). The patient had not been diagnosed prior to the presentation which is similar to our case but of younger age. Our case depicts an atypical presentation approximately more than 30 years after the occurrence of OSD with no history of trauma or forceful knee contraction.

However in few cases knee pain has been found to persist whether treatment is conservative or surgical (1,15). Despite resolution of symptoms, mobile osseous fragment fails to unite in approximately 10% of cases and causes pain during direct pressure on the tubercle and upon kneeling which has been reported in few studies (4). Despite removal of ossicles surgically recurrent ossicle formation have been found with the passage of time in a few patients however most were asymptomatic (1). In our patient there were no symptoms of any prior surgery for OSD or any pain on kneeling prior to the fracture which is quite different from previous reported cases.

The period between diagnosis of OSD and occurrence of avulsion fracture ranges from a few weeks to 1 year (16, 17). Therefore a period of relative rest with restricted sports activity till physal fusion or radiological healing is advised as healing at skeletal maturity is the rule. In our case the patient had avulsion fracture at 49 years of age. This was unusual and has not been reported before. We agree with Niremberg et al (18) that surgical excision of the symptomatic mobile fragment should be done. At 6 years follow up our patient has normal quadriceps strength and normal range of motion.


**Conclusion:** To our knowledge the reported case is the first one of spontaneous avulsion of the tibial tubercle following OSD in an adult. Simple excision of the fragment led to an excellent result and can be recommended.

93     **Literature**

- 94     1. Pihlajamäki HK, Mattila VM, Parviainen M, Kiuru MJ, Visuri TI. Long-term  
95       outcome after surgical treatment of unresolved Osgood-Schlatter disease in  
96       young men. *J Bone Joint Surg Am*. 2009 Oct;91(10):2350-8.
- 97     2. Flowers MJ, Bhadreshwar DR. Tibial tuberosity excision for symptomatic  
98       Osgood-Schlatter disease. *J Pediatr Orthop*. 1995 May-Jun;15(3):292-7.
- 99     3. Glynn MK, Regan BF. Surgical treatment of Osgood-Schlatter's disease. *J*  
100       *Pediatr Orthop*. 1983 May;3(2):216-9
- 101    4. El-Husseini TF, Abdelgawad AA. Results of surgical treatment of unresolved  
102       Osgood-Schlatter disease in adults. *J Knee Surg*. 2010 Jun;23(2):103-7
- 103    5. Binazzi R, Felli L, Vaccari V, Borelli P. Surgical treatment of unresolved Osgood-  
104       Schlatter lesion. *Clin Orthop Relat Res*. 1993 Apr;(289):202-4.
- 105    6. Bang J, Broeng L. Spontaneous avulsion of the tibial tuberosity following  
106       Osgood-Schlatter disease. *Ugeskr Laeger*. 1995 May 22;157(21):3061-2.
- 107    7. Gottsegen CJ, Eyer BA, White EA, Learch TJ, Forrester D. Avulsion fractures of the  
108       knee:imaging findings and clinical significance. *Radiographics*. 2008 Oct;28(6):1755-  
109       70.
- 110    8. Stevens MA, El-Khoury G, Kathol MH, Brandser EA, Chow S. Imaging features of  
111       avulsion injuries.*RadioGraphics* 1999;19:655–672.
- 112    9. Rosenberg ZS, Kawelblum M, Cheung YY, Beltran J, Lehman WB, Grant AD.  
113       Osgood-Schlatter lesion: fracture or tendinitis? Scintigraphic, CT, and MR  
114       imaging features. *Radiology* 1992;185:853–858.
- 115    10. Ogden JA, Southwick WO. Osgood-Schlatter's disease and tibial tuberosity  
116       development. *Clin Orthop* 1976; 116:180–189.
- 117    11. Weiss JM, Jordan SS, Andersen JS, Lee BM, Kocher M. Surgical treatment of  
118       unresolved Osgood-Schlatter disease: ossicle resection with tibial tubercleplasty  
119       *J Pediatr Orthop*. 2007 Oct-Nov;27(7):844-7.
- 120    12. . Balmat P, Vichard P, Pem R. The treatment of avulsionfractures of the tibial  
121       tuberosity in adolescent athletes. *Sports Med* 1990;9:311–316
- 122    13. Gowda NBS, Kumar MJ. Simultaneous Bilateral Tibial Tubercle Avulsion Fracture  
123       in a case of Pre-Existing Osgood-Schlatter Disease (OSD). *Journal of*  
124       *Orthopaedic Case Reports* 2012; 2(1): 24- 27.
- 125    14. Ogden JA, Tross RB, Murphy MJ. Fractures of the tibial tuberosity in  
126       adolescents. *J Bone Joint Surg Am* 1980;62:205–215

- 127 15. Krause BL, Williams JP, Catterall A. Natural history of Osgood Schlatter disease  
128 .Pediatr Orthop 1990;10:65-8.
- 129 16. Chow SP, Lam JJ, Leong JC. Fracture of the tibial tubercle in the adolescent. J  
130 Bone Joint Surg Br. 1990 Mar;72(2):231-4.
- 131 17. Levi JH, Coleman CR. Fracture of the tibial tubercle. Am J Sports Med. 1976  
132 Nov-Dec;4(6):254-63.
- 133 18. Nierenberg G, Falah M, Keren Y, Eidelman M. Surgical treatment of residual  
134 Osgood schlatter disease in young adults: role of the mobile osseous fragment.  
135 Orthopedics. 2011 Mar 11;34(3):176.

136 Figures



137  
138 Figure 1: X- ray of the left knee ap (a) and lateral view (b) shows an avulsion of the  
139 tibial tubercle following Osgood- Schlatters' disease (red arrow)



140

141

142 Figure 2: MRI of the left knee reveals an avulsion of the tibial tubercle following

143 Osgood- Schlatters' disease with surrounding liquid



144

145

146 **Figure 3:**

147 Lateral X-ray of the left knee shows the status after excision of the unresolved

148 Osgood-Schlatter.