



SDI Review Form 1.6

Journal Name:	Asian Journal of Orthopaedic Research
Manuscript Number:	Ms_AJORR_40915
Title of the Manuscript:	Antero-medial Closed Enucleation of the Talus: A case report
Type of the Article	Case Study

General guideline for Peer Review process:

This journal's peer review policy states that **NO** manuscript should be rejected only on the basis of '**lack of Novelty**', provided the manuscript is scientifically robust and technically sound. To know the complete guideline for Peer Review process, reviewers are requested to visit this link:

(<http://www.sciencedomain.org/page.php?id=sdi-general-editorial-policy#Peer-Review-Guideline>)



SDI Review Form 1.6

PART 1: Review Comments

	Reviewer's comment	Author's comment (if agreed with reviewer, correct the manuscript and highlight that part in the manuscript. It is mandatory that authors should write his/her feedback here)
Compulsory REVISION comments		
Minor REVISION comments	<p>Line 62: You mentioned that the immobilization with splint was two weeks then by the cast for three weeks (above knee cast) and finally more three weeks by below knee cast. What is the value of immobilization by the splint to begin with over the beginning by the cast at first?</p> <p>Line 64: As the case was internally fixed, why the rehabilitation program has not been started early? The goals of rehabilitation such as reduction of stiffness and improving the blood circulation in that area may promote healing and prevent the chance of occurrence of avascular necrosis.</p> <p>Line 83: Determine which type of enucleation that conservative treatment can be succeeded to solve the problem. If you meant that all types of enucleation can be treated conservatively, please clarify that in this paragraph.</p> <p>Line 91: You mentioned that the immobilization by the cast was recommended for eight weeks. Does it mean that the splint which you mentioned at first is like the cast?</p> <p>Line 101: I think that the word "inevitable" is opposing the idea of lines 87 to 89 stating that open reduction of antero-medial and antero-lateral enucleation should be done only when the conservative treatment was failed. According to your experience, write a clear statement about this issue in the discussion section answering the question if surgeons can follow procedures of open reduction in such cases directly or wait for closed reduction and conservative treatment at first.</p> <p>Plagiarism Issue – https://app.ithenticate.com/en_us/dv/0425?lang=en_us&o=35965524 (22%)</p>	
Optional/General comments		

Reviewer Details:

Name:	Waleed Salah El-din Mahmoud
Department, University & Country	Physical Therapy and Health Rehabilitation Department, Prince Sattam bin Abdulaziz University, Saudi Arabia

