

## REVIEW OF ANTIBIOTICS USAGE IN THE MANAGEMENT OPEN FRACTURE IN A NIGERIAN HOSPITAL

### ABSTRACT

**BACKGROUND.** Antibiotics are adjuncts in the management of open fractures, and microbial characteristic of open fractures will guide the use of antibiotics. With changing pattern in microbial colonization of wounds, the need to review antibiotic usage in hospitals becomes imperative. The study aimed to evaluate the antibiotic protocol of managing open fractures at the Accident and Emergency department, with the advent of new antibiotics introduced into the hospital drug formulary.

**MATERIALS AND METHODS.** This study is a hospital-based prospective evaluation of the antibiotic sensitivity of cultured microorganisms from the patients with open fractures presenting between January 2013 and December 2013 in the Accident and Emergency Department, of a tertiary hospital in Nigeria. Swabs of superficial and deeper parts of the wound were taken at the presentation of the patients before wound debridement and commencement of antibiotics. Other two samples and biopsies were taken at the deeper parts of the wound on the 3rd and 7th day of admission. Culture and Sensitivity pattern of isolates were determined for positive cultures using antibiotics impregnated disks. Descriptive and inferential statistics of the findings are presented.

**RESULT.** One hundred and thirty patients with open fractures were recruited for the study, but 81 patients completed the study. . Forty patients discharged themselves against medical advice and while nine patients were referred to other hospitals. Sterile swabs were taken from the surface and deeper portion of the wounds at presentation before treatment was commence and at day 3 and day 7 of admission, *Staphylococcus aureus* and *Clostridium perfringens* were the most common aerobic and anaerobic isolates from the wounds respectively. The aerobic isolates and anaerobes were susceptible to ceftriaxone, ciprofloxacin, co-Amoxyclav, gentamycin, and cefotaxime and metronidazole respectively.

**CONCLUSION.** The antibiotic sensitivity pattern in the emergency department of the Hospital has changed not significantly as previously reported about 12 years earlier. Therefore, the hospital antibiotic protocol in the treatment of open fractures in the Accident and Emergency department should be retained.

**Keywords:** Open fracture. Antibiotics sensitivity, Antibiotic usage, Ibadan, Nigeria

## 7 **Introduction**

8 The choice of antibiotics in the treatment of open fractures as an adjunct to debridement and  
9 wound care, is determined by established microbial characteristics of open fractures in the  
10 locality or empirically using combination therapy to cover most of the available organisms  
11 such as Gram-positive and Gram-negative aerobes as well as the anaerobes. The choice of  
12 antibiotics in the treatment of infections is determined by the potential bacterial  
13 contamination based on historical or research documented patterns for each locality [1]. On  
14 account of their findings, Wilkins and Patzakakis recommended the use of a combination of  
15 cephalosporins, penicillins and aminoglycosides in open fractures depending on the severity  
16 of the wound and extent of contamination [2]. However, Alonge et al. in Ibadan Nigeria,  
17 found that pefloxacin, ciprofloxacin and ceftriaxone were the antibiotics which exhibited  
18 relatively higher sensitivity to the micro-organisms isolated [3], which is in agreement with  
19 the findings in other studies [4] [5] [6] [7][8].

20 Open fractures usually result from high energy trauma such as motor vehicle crashes, falls  
21 from height, gunshot injury, assault and machine injury [5] and are prone to contamination  
22 and infection [4][9][10]. Research have helped refine surgical protocols, change in antibiotic  
23 prescriptions, and in defining the appropriate timing for interventions including debridement,  
24 modalities of fracture fixation, and soft tissue coverage [11][12][13][14][15][16]. Infections  
25 in open fractures becomes more likely after six hours of injury if adequate surgical treatment  
26 is not carried out along with the administration of appropriate antibiotics early enough after  
27 the injury. Deep fracture site infections could lead to complications of chronic osteomyelitis,  
28 nonunion and sometimes limb loss. Apart from the exposure of the fractured bone, numerous  
29 predisposing factors which influence the development of infection include shock from blood  
30 loss, hypoxia and the degree of comminution [17]. Majority of infections in open fractures  
31 are caused by *Staphylococci* species especially *Staphylococcus aureus* and coagulase-  
32 negative *Staphylococci*, gram-negative bacilli which include *Acinetobacter spp*, *Escherichia*  
33 *coli*, *Pseudomonas spp*, *Klebsiella spp* and *Proteus spp* amongst others [4][14][17]. Alonge et  
34 al. in 2002 established that *E coli* was the most prevalent single isolate while *Staphylococcus*  
35 *aureus* was the most prevalent microbial isolate in poly-microbial infections [1] in their  
36 locality.

37 Resistance to available antimicrobial drugs is an established and ever-growing challenge in  
38 clinical practice. Such resistance can result from two mutually non-exclusive phenomena:  
39 mutations in house-keeping structural or regulatory genes and the horizontal acquisition of  
40 foreign genetic information [18]. Outbreaks of infections due to *Klebsiella pneumonia*  
41 harboring plasmid-encoded cephalosporinases and the spread of this resistance mechanism to  
42 bacterial species naturally susceptible to cephamycins have been reported [19].

43 An infection engrafted on a biomaterial (thick, adherent biofilm) responds poorly to  
44 antimicrobial therapy and usually is not cured until the biomaterial is removed. Bacterial  
45 isolates may not be entirely representative of the microbial components of the biofilm  
46 because the coherent properties of the adherent biofilms that are found on surfaces in these  
47 infections may prevent genuinely representative organisms from detaching in sufficient

numbers to be detected entirely and consistently by simple sampling and routine culture techniques. Therefore, antimicrobials that are chosen from the culture results may not be effective against all of the bacterial species in these biofilm infections [20].

This study aimed to review the antibiotic treatment protocol for open fractures in the A&E of a tertiary hospital in Nigeria with the view for recommendations for possible change in practice.

## **MATERIALS AND METHODS**

This study is a hospital-based prospective evaluation of antimicrobial pattern and antibiotics sensitivity pattern in open fractures presenting in the Accident and Emergency Department of the University College Hospital, Ibadan from January 2013 to December 2013 following approval of the Hospital Ethical and Research Committee according to Helsinki Declaration of 1977 modified in 2000.

**Emergency Room Protocol:** Proforma for the study was completed for all patients seen in the Accident and Emergency department of the hospital with open fracture after obtaining securing informed consent from the included patients. Patients with open fractures who had wound debridement and antibiotics before presenting at the Accident and Emergency of the University College Hospital, Ibadan were excluded.

Patients with open fractures were resuscitated and treated using the advanced trauma life support (ATLS) protocol. The associated open fractures were inspected, and clinical photographs obtained as appropriate.. Four sterile wound swabs, (superficial aerobic and anaerobic, deep aerobic and anaerobic) were collected from the superficial and deep parts of open fracture wounds using the Levine's technique. The swabs of the wounds were obtained aseptically before wound debridement and antibiotics were commenced within 30 minutes of patient's arrival at the Accident and Emergency Department. Two other samples and biopsies were taken at the deeper parts of the wounds on the 3<sup>rd</sup> and 7<sup>th</sup> day of admission. Samples were collected into sterile Stuarts transport medium, and sterile Robertson cooked meat medium for aerobic and anaerobic organisms respectively. The samples were labelled "S" for superficial swab samples, "D" for deep swab samples, "BS" and "BD" for superficial and deep biopsy samples with the patient's research number on the laboratory request form and also on the bottle. Having collected the samples, the open fracture was treated according to the hospital established protocol of early antibiotic administration, analgesics, tetanus prophylaxis, early wound debridement, fracture stabilization and early soft tissue coverage.

**Laboratory Protocol:** All obtained samples arrived the laboratory within 30 minutes to 3 hours of collection. The samples were stored at room temperature in a cupboard for less than 6 hours until ready for analysis. Microscopy, culture and sensitivity patterns of the samples to various antibiotics (penicillin, cephalosporin, quinolone, aminoglycoside, clindamycin, sulphonamides and trimethoprim, and metronidazole) were carried out. The samples for

aerobic cultures were plated out on sterile Sheep blood agar and MacConkey agar aseptically and incubated at 37°C for 24 hours. The direct Gram staining of the swabs was carried out, and the slides examined to identify the presence of organisms and pus cells. After 24 hours of incubation, the plates were analyzed for the growth of the bacteria and gram staining of the bacteria colonies were carried out.


The confirmatory test of all the isolated gram-negative bacilli was based on the use of API 20 E while the gram-positive cocci were based on the use of control organisms for coagulase test. Sensitivity testing was carried out using the disc diffusion technique (Bauer Kirby method), The anaerobic samples were inoculated aseptically into a sterile Sheep blood agar and MacConkey agar within five minutes of sample collection. The inoculated plates were incubated in the anaerobic gas chamber containing anaerobic catalytic agent, Anaero Gen kit and anaerobic control kit (Oxoid Ltd of United Kingdom). Strict anaerobic control bacteria and strict aerobic bacteria were also included as an added quality control. The anaerobic organisms were left in the chamber to incubate at 37°C for three days to isolate the fast-growing anaerobes which are mostly contaminants while the late growing anaerobes were further incubated for ten days and these are the bacteria of medical importance.

Data was managed and analysed using analysed using IBM SPSS Statistics for Windows Version 20.0 (Armonk, NY: IBM Corp). Descriptive statistics are presented as proportions and percentages as well as by use of appropriate scientific figures. Chi-square ( $\chi^2$ ) was used to test for statistical significance for observed differences for categorical variables. P - values less than 0.05 were accepted as significant.


## RESULTS

Eighty-one of the 130 recruited patients completed the study with superficial and deep swab samples taken from all patients on the first day and other swabs and biopsy samples taken on the third and seventh day of admission. Forty patients took their discharges against medical advice while nine patients were referred to other hospitals of their choice. There were 93 (71.5%) male and 37 (28.5%) female patients as shown in figure 1 while figure 2 represents open fractures in different regions of the body with the tibia and fibula constituting 78 (60%) of the cases while the femur accounted for 19 (14.6%). Gustilo and Anderson type 3B [21] was the most common grade of open fracture 48 (36.9%), while type 3A occurred in 43 (33.1%) as presented in figure 3. The microbial culture shows that *Staphylococcus aureus* and *Clostridium perfringens* were the predominant aerobic and anaerobic isolates.

**Figure 1: Showing the sex distribution**



**Figure 2: Shows open fracture in the various regions of the body**



**Figure 3: Shows the grades of open fracture**





Figure 4: Antibiotic sensitivity pattern for aerobes (blue) and anaerobes (red)



133

#### Abbreviations

134 CRO – cephtriaxone, CF – cefazolin, CN – gentamycin, LFX – levofloxacin, RS –rosoxacin,  
 135 AML –amoxycillin, CLM – clindamycin, CXM – cefuroxime, SP – sparfloxacin, TET –  
 136 tetracycline, AMC – co-Amoxycylav, AMX – amoxycillin, GP – ciprofloxacin, CAZ –  
 137 ceftazidime, PEF- pefloxacin, CTR – cefotaxime, SPX – sparfloxacillin, SN-sulphonamides,  
 138 AX – amoxycillin, AMP – ampicillin, MTZ – metronidazole and COT –cotrimoxazole.  
 139  
 140

141 The antibiotic sensitivity pattern are shown in figure 4 and tables 1and 2. Ciprofloxacin  
 142 (GP), ceftriaxone (CRO), co-amoxiclav (AMC) and gentamycin (CN) were the drugs most  
 143 aerobic organisms were sensitive to, while anaerobic microorganisms were highly sensitive to  
 144 cefotaxime (CTR), and metronidazole (MTZ).

Table 1. Aerobic Organism sensitivity

Organism	Antibiotics										$\chi^2$ ; P value
.	CRO	CN	LFX	CXM	AMG	AMX	GP	CAZ	CTR	MTZ	
SA	5	4	1	2	3	3	4	0	0	0	3.33; 0.50
EC	0	1	2	0	1	0	1	2	0	0	2.0; 0.74
KS	3	1	0	1	5	1	4	0	0	0	8.0; 0.09
PsA	2	1	1	0	0	0	2	1	0	0	2.0; 0.74
$\chi^2$ ;	2.67	3.0	2.0	2.0	5.0	6.0	1.3	2.0			
P value	0.45	0.39	0.57	0.57	0.17	0.11	.72	0.57			

146 Key: *S A* – *Staphylococcus aureus*, *E C* – *Escherichia coli*, *K S* – *klebsiella spp*, and *PsA* –  
 147 *Pseudomonas auregenosa*.

148

149

150 **Table 2. Anaerobic Organism sensitivity**

Organism	Antibiotics										$\chi^2$ ; P value
	CRO	CN	LFX	CXM	AMG	AMX	GP	CAZ	CTR	MTZ	
CP	3	0	2	2	0	0	3	1	3	20	41.33 <0.001
BS	0	0	0	0	0	0	0	0	0	5	20.00 <0.001
CT	1	0	1	0	0	0	1	0	2	9	15.33 0.004
AI	4	1	0	2	0	0	4	2	1	0	5.0 0.29
$\chi^2$ ; P value	5.0; 0.17	- -	2.0 0.57	4.0 0.26	- -	- -	5.0 .17	2.0 0.57	1.0 0.80	24.89 <0.001	

151 Key: *C P* – *Clostridium perfringens*, *C T* – *Clostridium tetani*, *B S* – *Bacteroides spp* and *A I*  
 152 – *Actinomyces israelii*.

### 153 Discussion

154 The role of early wound debridement and antibiotic administration is recognized as necessary  
 155 in the management of open fractures in the hospital. Appropriate antibiotic(s) are  
 156 administered according to the established hospital protocol following the identified historical  
 157 and sensitivity pattern of wound swabs [22]. The value of antibiotics in the treatment of open  
 158 fractures has been established, but this does not substitute for proper wound debridement and  
 159 adequate skeletal stabilization as an essential aspect of open fracture management. The  
 160 choice of antibiotic should be guided by the knowledge of possible contaminating organisms  
 161 at presentation, but subsequent infections are most likely multiple organisms which should be  
 162 covered by choice of antibiotics. Evidence-based guidelines for prophylactic antibiotic use in  
 163 open fractures recommend short-course, narrow-spectrum antibiotics for Gustilo Grade I or II  
 164 open fractures and broader gram-negative coverage for Grade III open fractures [23].

165

The antibiotic protocol in the Accident and Emergency Department of the hospital, for the treatment of open fractures, has been a combination of ceftriaxone, quinolones (ciprofloxacin) and metronidazole-based on findings of Alonge et al. in 2002 [3]. The result of Alonge et al. which had observed *Escherichia coli* as the most common single gram-negative aerobic isolate sensitive to ceftriaxone, quinolones, was slightly at variance to the findings from this study which showed that *Staphylococcus aureus* and *Clostridium perfringens* as the most common single aerobic and anaerobic isolates respectively. Since anaerobic organisms were not cultured by where as Alonge and colleagues, the inclusion of metronidazole in the hospital antibiotic protocol was based on evidence from other practices.

From this study, the predominant aerobic gram-positive organism (*Staphylococcus aureus*) was sensitive to ceftriaxone (CRO), Gentamycin (CN), co-amoxiclav (AMC), cefuroxime (CXM) and amoxycillin (AMX) while the aerobic gram-negative organisms (*Escherichia coli* and *Klebsiella spp*) were sensitive to ceftriaxone, amoxycillin, levofloxacin and ceftazidime. However, the sensitivity of the aerobic isolates to tested antibiotics was marginally statistically significant to ceftriaxone,  $P = 0.50$ , but statistically significantly sensitivity to metronidazole,  $P < 0.05$ . The antibiotic sensitivity pattern was not too different from the findings by Alonge et al. 2002 and other studies [3][4][5]. Also, isolated anaerobes were significantly sensitive to metronidazole (MTZ) and moderately sensitive to ceftriaxone, levofloxacin, cefuroxime, ciprofloxacin and cefotaxime (CTR), justifying the inclusion of metronidazole in the hospital antibiotic protocol.

The micro-organisms cultured in this study showed high resistance to ampicillin (AMP), cotrimoxazole (COT), sulphonamides (SN), clindamycin (CML), rosoxacin (RS), amoxycillin, cefazolin (CF), and tetracycline (TET). The aerobic gram-positive organisms were resistance to ceftazidime (CAZ), cefotaxime (CTR) and metronidazole while the aerobic gram-negative microorganisms were resistance to cefotaxime), metronidazole, amoxycillin, cefuroxime). The anaerobic organisms also showed significant resistance to co-amoxyclav, amoxycillin, gentamycin and Ceftazidime. These findings are comparable to a similar study in another African hospital by Sitali and colleagues in 2017 [24].

Apart from antibiotic sensitivity and microbial patterns, the hospital antibiotic protocol is also influenced by the cost and availability of the drugs. In the centre where this study was undertaken as well as in most hospitals in the region, availability of some of the antibiotics can be challenging. Even when the drugs are available, affordability often becomes another challenge as the majority of persons that in the region live below the WHO poverty line [25]. The use of generic forms of these antibiotics, therefore, is the norm in the region.

It is worth noting that cultured isolates from a wound especially in the presence of biomaterials and biofilms may not be truly representative of the actual organisms causing infections. Since an infection engrafted on a biomaterial (thick, adherent biofilm) responds



poorly to antimicrobial therapy and is not usually cured until the biomaterial is removed, the reliance on only antibiotics without appropriate debridement of dead tissue should be with caution. Antimicrobials that are chosen from the swab culture results may not be effective against all of the bacterial species in these biofilm infections [26]. Incidentally, it takes some time before biofilm develops. However, since the cultures in this study were all done within seven days of admission, the identified sensitivity patterns may not be entirely reflective of the antibiotic sensitivity and resistance in open fractures with chronic wounds where there is an existence of biofilms.

## CONCLUSION

The hospital antibiotic protocol which recommends the combination of ceftriaxone, quinolones, gentamycin, co-amoxycylav and metronidazole in treating open fractures in the Accident and Emergency department, was based on the antibiotics sensitivity to cultured microbial organisms in the hospital. The existing microbial and antibiotic sensitivity patterns had not changed significantly over the preceding 12 years when the protocol was established as such there is no reason for a change in the current practice.

## REFERENCES

1. Gustilo RB, Anderson JT. Prevention of infection in the treatment of one thousand and twenty-five open fractures of long bones: retrospective and prospective analyses. *J Bone Joint Surg. Am.* 1976;58:453-8
2. Levine NS, Lindberg B, Mason AD Jr, Pruitt BA Jr. The quantitative swab culture and smear: a quick method for determining the number of viable aerobic bacteria on open wounds. *J Trauma* 1976;16: 89–94.
3. Alonge TO, Ogunlade SO, Salawu SA, Fashina AN. Microbial isolates in open fractures seen in the accident and emergency unit of a teaching hospital in a developing country. *West African J Med.* 2002;21(4):302-4.
4. Ikem IC, Oginni LM, Bamgboye EA, Ako-Nai AK, Onipade AO. The bacteriology of open fractures in Ile-Ife, Nigeria. *Niger J Med.* 2004; 13: 359-65.
5. Bowler P, Duerden B, Armstrong D. Wound microbiology and associated approaches to wound management. *Clinical Microbiology Reviews* 2001; 14(2): 244-69.
6. Wilkins J, Patzakis M. Choice and duration of antibiotics in open fractures. *Ortho Clin N Amer.* 1991; 22(3): 433-7.
7. Akinyoola AL, Ako-Nai AK, Dosumu O, Aboderin AO, Kassim OO. Microbial isolates in early swabs of open musculoskeletal injuries. *Niger Postgrad Med J.* 2006; 13(3): 176-81.
8. Alonge TO, Salawu SA, Adebisi AT, Fashina AN. The choice of antibiotic in open fractures in a teaching hospital in a developing country. *Int J Clin Pract.* 2002; 56(5): 353 – 6.

244 9. Ifesanya AO, Alonge TO, Ogunlade SO, Omololu AB, Nottidge TE, Ayorinde RO.  
 245 Changing Trends in the pattern of tibial fractures in Nigeria: A review of 70 cases.  
 246 J.Orthopaedics 2008;5(2)e4.

247 10. Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III  
 248 (severe) open fractures: A new classification of type III open fractures. J Trauma 1984;24:  
 249 742 - 6.

250 11. Paul H. Kim MD, Seth S. Leopold MD. Gustilo-Anderson classification: Clin Orthop  
 251 Relat Res 2012;470: 3270 –4.

252 12. Zalavras CG, Marcus RE, Levin LS, Patzakis MJ. Management of open fractures and  
 253 subsequent complications. J. Bone Joint Surg. Am. 2007;89: 884-95.

254 13. Okike K, Bhattacharyya T. Trends in the management of open fractures. A critical  
 255 analysis. J Bone Joint Surg. Am. 2006;88: 2739-48.

256 14. Patzakis MJ, Wilkins J. Factors influencing infection rate in open fracture wounds.  
 257 ClinOrthopRelat Res. 1989;243: 36 – 40.

258 15. Ostermann PA, Seligson D, Henry SL. Local antibiotic therapy for severe open fractures.  
 259 A review of 1085 consecutive cases. J Bone Joint Surg Br. 1995; 77: 93 – 7.

260 16. Gosselin RA, Roberts I, Gillespie WJ. Antibiotics for preventing infection in open limb  
 261 fractures. Cochrane Database Syst Rev. 2004; (1) CD003764.

262 17. Khosravi AD, Ahmadi F, Salmanzadeh S, Dashtbozorg A, AbasiMontazeri E. Study of  
 263 bacteria isolated from orthopedic implant infections and their antimicrobial susceptibility  
 264 pattern. Res J Microbiol. 2009; 4: 158-63.

265 18. Courvalin P. Antimicrobial Drug Resistance: “Prediction Is Very Difficult, Especially  
 266 about the Future.” Emerg Infect Dis. 2005;11: 1503-6.

267 19. Garazzino S, Aprato A, Maiello A, Masse A, Biasibetti A, De Rosa FG, Di Per G.  
 268 Osteomyelitis Caused by Enterobacter cancerogenus Infection following a Traumatic  
 269 Injury: Case Report and Review of the Literature. Journal of Clinical Microbiology  
 270 2005; 43(3):1459 –61.

271 20. Gristina AG, Costerton JW. Bacterial adherence to biomaterials and tissue. The  
 272 significance of its role in clinical sepsis. J Bone Joint Surg Am. 1985; 67(2):264-73.

273 21. Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III  
 274 (severe) open fractures: A new classification of type III open fractures. J Trauma 1984; 24:  
 275 742 - 6.

276 22. Purghel F, Badea R, Ciuvica R, Anastasiu A. The use of antibiotics in traumatology and  
 277 Orthopaedic surgery. Journal of clinical medicine, 2006; 1: 58-65.

278 23. Yishak A., Wamisho BL. Microbial susceptibility of bacteria isolated from open fracture  
 279 wounds presenting to the Err of black-lion hospital, Addis Ababa University, Ethiopia  
 280 African Journal of Microbiology Research 2009; 3(12): 939 - 51.

281 24. Sitati FC, Mosi PO, Mwangi JC. Early Bacterial Cultures from Open Fractures -  
 282 Differences Before and After Debridement. Ann Afr Surg. 2017;14(2):66-70.

- 283 25. World Bank. World Bank Country indicators.  
284 Retrived from: <https://www.indexmundi.com/g/g.aspx>. Accessed: 12<sup>th</sup> December, 2017.
- 285 26. Rodriguez L, Jung HS, Goulet JA, Cicalo A, Machado-Aranda DA, Napolitano LM.  
286 Evidence-based protocol for prophylactic antibiotics in open fractures: Improved antibiotic  
287 stewardship with no increase in infection rates. Journal of Trauma and Acute Care Surgery.  
288 2014; 77(3):400–8.

289

290