

On the Properties and Applications of Transmuted Odd Generalized Exponential-Exponential Distribution

Abstract:

This article proposed a new distribution referred to as the transmuted odd generalized exponential-exponential distribution (*TOGEED*) as an extension of the popular odd generalized exponential- exponential distribution by using the Quadratic rank transmutation map (*QRTM*) proposed and studied by [12]. Using the transmutation map, we defined the probability density function (*pdf*) and cumulative distribution function (*cdf*) of the transmuted odd generalized Exponential- Exponential distribution. Some properties of the new distribution were extensively studied after derivation. The estimation of the distribution's parameters was also done using the method of maximum likelihood estimation. The performance of the proposed probability distribution was checked in comparison with some other generalizations of Exponential distribution using a real life dataset.

Keywords: *QRTM, Properties, Maximum likelihood estimation, TOGEED, Applications.*

1 Introduction

An Exponential distribution which can be used in Poisson processes gives a description of the time between events. The distribution has been applied widely life testing experiments. The distribution exhibits memoryless property with a constant failure rate which makes the distribution unsuitable for real life problems and hence creating a vital problem in statistical modeling and applications.

There are several ways of adding one or more parameters to a distribution function which makes the resulting distribution richer and more flexible for modeling data. Some of the

recent studies on the generalization of exponential distribution include the transmuted exponential distribution by [10], transmuted inverse exponential distribution by [8], the odd generalized exponential-exponential distribution by [5] and the Weibull-Exponential distribution by [9]. Of interest to us in this article is the odd generalized exponential-exponential distribution (*OGEED*) which can be use in various fields to model variables whose chances of success decreases with time whereas those of failure increases as time increases. It was also discovered that the *OGEED* is positively skewed and performed better than some existing distributions like Gamma, exponentiated exponential, Weibull and Pareto distributions.

According to [5], a random variable X is said to follow a odd generalized exponential-exponential distribution with parameters α and θ if its probability density function (*pdf*) is given as

$$f(x) = \alpha \theta e^{\theta x} e^{-\alpha(e^{\theta x} - 1)} \quad (1.1)$$

and the corresponding cumulative distribution function (*cdf*) is given as

$$F(x) = 1 - e^{-\alpha(e^{\theta x} - 1)} \quad (1.2)$$

39

The *pdf* and *cdf* of the transmuted *OGEED* are obtained using the steps proposed by [12]. According to [12], a random variable X is said to have a transmuted distribution function if its *pdf* and *cdf* are respectively given by;

$$f(x) = g(x)[1 + \lambda - 2\lambda G(x)] \quad (1.3)$$

and

$$F(x) = (1 + \lambda)G(x) - \lambda[G(x)]^2 \quad (1.4)$$

46 respectively.

47 where; $x > 0$, and $-1 \leq \lambda \leq 1$ is the transmuted parameter, $G(x)$ is the *cdf* of any continuous
48 distribution while $f(x)$ and $g(x)$ are the associated *pdf* of $F(x)$ and $G(x)$, respectively.

49 The aim of this paper is to introduce a new continuous distribution called the Transmuted
50 Odd Generalized Exponential-Exponential distribution (*TOGEED*) with a decreasing failure
51 rate useful for analyzing real life data using the proposed quadratic rank transmutation map
52 by [12]. The remaining parts of this paper are presented in sections as follows: We defined
53 the new distribution and give its plots in section 2. Section 3 derived some properties of the
54 new distribution. The estimation of parameters using maximum likelihood estimation (MLE)
55 is provided in section 4. An application of the new model with other two existing one to a
56 real life data is done in section 5 and some useful conclusions are made in section 5.

57 **2. The Transmuted Odd Generalized Exponential-Exponential Distribution** 58 **(*TOGEED*)**

59 Substituting equation (1.1) and (1.2) in (1.3) and (1.4) and simplifying, we obtain the *cdf* and
60 *pdf* of the *TOGEED* as follows:


$$61 \quad F(x) = 1 - e^{-\alpha(e^{\theta x} - 1)} \left[1 - \lambda + \lambda e^{-\alpha(e^{\theta x} - 1)} \right] \quad (2.1)$$

62
and

$$63 \quad f(x) = \alpha \theta e^{\theta x} e^{-\alpha(e^{\theta x} - 1)} \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x} - 1)} \right] \quad (2.2)$$

64 respectively. Where, $x > 0, \alpha > 0, \theta > 0, -1 \leq \lambda \leq 1$, α is the shape parameters respectively and
65 θ is the exponential parameter while λ is called the transmuted parameter.


66 The *pdf* and *cdf* of the *TOGEED* using some parameter values are displayed in **figures 2.1**
 67 and **2.2** as follows.



68

69 **Fig. 2.1:** *PDF* of the *TOGEED* for different values of the parameters α, θ —and— λ as
 70 displayed in the key on the graph.

71 Figure **2.1** indicates that the *TOGEED* distribution could be positively skewed and otherwise
 72 depending on the parameter values. This means that distribution can be very useful for
 73 datasets with right-tailed shapes.



74

75 **Fig. 2.2:** *CDF* of the *TOGEED* for different values of the parameters α, θ —and— λ as
 76 displayed in the key on the graph.

77 From the above *cdf* plot, the *cdf* increases when X increases, and approaches 1 when X
 78 becomes large, as expected.

79 3. Properties

80 In this section, we defined and discuss some properties of the *TOGEED* distribution.

81 3.1 Moments

82 Let X denote a continuous random variable, the n^{th} moment of X is given by;

$$83 \quad \mu'_n = E(X^n) = \int_0^{\infty} x^n f(x) dx \quad (3.1.1)$$

84 where $f(x)$ the *pdf* of the *TOGEED* is as given in equation (3.1.1).

$$85 \quad f(x) = \alpha \theta e^{\theta x} e^{-\alpha(e^{\theta x}-1)} \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x}-1)} \right]$$

86 Expansion and simplification of the *pdf* gives:

$$87 \quad f(x) = (1-\lambda)\alpha\theta e^{\theta x} e^{-\alpha(e^{\theta x}-1)} + 2\lambda\alpha\theta e^{\theta x} e^{-2\alpha(e^{\theta x}-1)} \quad (3.1.2)$$

88

Hence,

$$89 \quad \mu'_n = E(X^n) = \int_0^{\infty} x^n f(x) dx = \int_0^{\infty} x^n \left\{ (1-\lambda)\alpha\theta e^{\theta x} e^{-\alpha(e^{\theta x}-1)} + 2\lambda\alpha\theta e^{\theta x} e^{-2\alpha(e^{\theta x}-1)} \right\} dx$$

$$90 \quad \mu'_n = \int_0^{\infty} x^n (1-\lambda)\alpha\theta e^{\theta x} e^{-\alpha(e^{\theta x}-1)} dx + \int_0^{\infty} x^n 2\lambda\alpha\theta e^{\theta x} e^{-2\alpha(e^{\theta x}-1)} dx$$

$$91 \quad \mu'_n = (1-\lambda)\alpha\theta \int_0^{\infty} x^n e^{\theta x} e^{-\alpha(e^{\theta x}-1)} dx + 2\lambda\alpha\theta \int_0^{\infty} x^n e^{\theta x} e^{-2\alpha(e^{\theta x}-1)} dx \quad (3.1.3)$$

92

Using integration by substitution in (3.1.3) above we have,

93

Let

$$w = e^{\theta x} - 1 \Rightarrow x = \frac{\ln(w)}{\theta}$$

$$\frac{dw}{dx} = \theta e^{\theta x} \Rightarrow dx = \frac{dw}{\theta e^{\theta x}}$$

Therefore, substituting for w and dx in equation (3.1.3) above and simplifying, we have:

$$\mu'_n = \frac{(1-\lambda)\alpha}{\theta^n} \int_0^\infty (\ln(w))^n e^{-\alpha w} dw + \frac{2\lambda\alpha}{\theta^n} \int_0^\infty (\ln(w))^n e^{-2\alpha w} dw \quad (3.1.4)$$

$$\int_0^\infty (\ln(w))^n e^{-\alpha w} dw = I(n, \alpha)$$

Using a result by [7], [1] and [13],

$$I(n, \alpha) = \left(\frac{\partial}{\partial a} \right)^n \left[(\alpha)^{-a} \Gamma(a) \right] \Big|_{a=1}$$

Reduces to

Hence, substituting and simplifying in (3.1.4), we have:

$$\mu'_n = \frac{(1-\lambda)\alpha}{\theta^n} \left(\frac{\partial}{\partial a} \right)^n \left[(\alpha)^{-a} \Gamma(a) \right] \Big|_{a=1} + \frac{2\lambda\alpha}{\theta^n} \left(\frac{\partial}{\partial a} \right)^n \left[(2\alpha)^{-a} \Gamma(a) \right] \Big|_{a=1}$$

(3.1.5)

The Mean

The mean of the *TOGEED* can be obtained from the n^{th} moment of the distribution when $n=1$ as follows:

$$\mu'_1 = \frac{(1-\lambda)\alpha}{\theta} \left(\frac{\partial}{\partial a} \right) \left[(\alpha)^{-a} \Gamma(a) \right] \Big|_{a=1} + \frac{2\lambda\alpha}{\theta} \left(\frac{\partial}{\partial a} \right) \left[(2\alpha)^{-a} \Gamma(a) \right] \Big|_{a=1} \quad (3.1.6)$$

Also the second moment of the *TOGEED* is obtained from the n^{th} moment of the distribution when $n=2$ as

$$\mu'_2 = \frac{(1-\lambda)\alpha}{\theta^2} \left(\frac{\partial}{\partial a} \right)^2 \left[(\alpha)^{-a} \Gamma(a) \right] \Big|_{a=1} + \frac{2\lambda\alpha}{\theta^2} \left(\frac{\partial}{\partial a} \right)^2 \left[(2\alpha)^{-a} \Gamma(a) \right] \Big|_{a=1} \quad (3.1.7)$$

110

The Variance

111 The n^{th} central moment or moment about the mean of X , say μ_n , can be obtained as

$$112 \quad \mu_n = E(X - \mu_1')^n = \sum_{i=0}^n (-1)^i \binom{n}{i} \mu_1'^i \mu_{n-i}' \quad (3.1.8)$$

113 The variance of X for *TOGEED* is obtained from the central moment when $n=2$, that is,

$$114 \quad Var(X) = E(X^2) - \{E(X)\}^2 \quad (3.1.9)$$

$$115 \quad Var(X) = \mu_2' - \{\mu_1'\}^2 \quad (3.1.10)$$

116 Where μ_1' and μ_2' are the mean and second moment of the *TOGEED* all obtainable from
117 equation (3.1.5).

118 3.2 Moment Generating Function

119 The *mgf* of a random variable X can be defined and obtained as given below:

$$120 \quad M_x(t) = E(e^{tx}) = \int_0^{\infty} e^{tx} f(x) dx \quad (3.2.1)$$

$$121 \quad M_x(t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} \mu_n' = \sum_{n=0}^{\infty} \frac{t^n}{n!} \left\{ \frac{(1-\lambda)\alpha}{\theta^n} \left(\frac{\partial}{\partial a} \right)^n [(\alpha)^{-a} \Gamma(a)] \Big|_{a=1} + \frac{2\lambda\alpha}{\theta^n} \left(\frac{\partial}{\partial a} \right)^n [(2\alpha)^{-a} \Gamma(a)] \Big|_{a=1} \right\}$$

122 (3.2.2)

123 3.3 Characteristics Function

124 The characteristics function of a random variable X is defined and given as follows;

$$125 \quad \varphi_x(t) = E(e^{itx}) = E[\cos(tx) + i \sin(tx)] = E[\cos(tx)] + E[i \sin(tx)]$$

126 (3.3.1)

127 Simple algebra and power series expansion proves that

$$\phi_x(t) = \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n}}{(2n)!} \mu_{2n}' + i \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1}}{(2n+1)!} \mu_{2n+1}'$$

(3.3.2)

Where μ_{2n}' and μ_{2n+1}' are the moments of X for $n=2n$ and $n=2n+1$ respectively and can be obtained from μ_n' in equation (3.1.5).

3.4 Reliability analysis of the *TOGEED*.

The Survival function describes the likelihood that a system or an individual will not fail after a given time. Mathematically, the survival function is given by:

$$S(t) = 1 - F(t) \quad (3.4.1)$$

Applying the *cdf* of the *TOGEED* in (2.1), the survival function for the *TOGEED* is obtained as:

$$S(t) = e^{-\alpha(e^{\theta t} - 1)} \left[1 - \lambda + \lambda e^{-\alpha(e^{\theta t} - 1)} \right] \quad (3.4.2)$$

The following is a plot for the survival function of the *TOGEED* using different parameter values as shown in Figure 3.4.1 below;




Figure 3.4.1: Survival function of the *TOGEED* at different parameter values.

The graph in figure 3.4.1 shows that the probability of the survival equals one (1) at initial time or early age and it decreases as t increases and equals zero (0) as t becomes larger.

Hazard function is the probability that a component will fail or die for an interval of time.

The hazard function is defined as;

$$h(t) = \frac{f(t)}{S(t)} = \frac{f(t)}{1 - F(t)} \quad (3.4.3)$$

Meanwhile, the expression for the hazard rate of the *TOGEED* is given by

$$h(t) = \frac{\alpha \theta e^{\theta t} \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta t} - 1)} \right]}{\left[1 - \lambda + \lambda e^{-\alpha(e^{\theta t} - 1)} \right]} \quad (3.4.4)$$

where $\alpha, \theta > 0$ and $-1 \leq \lambda \leq 1$.

The following is a plot of the hazard function at chosen parameter values in figure 3.4.2




Figure 3.4.2: The hazard function of the *TOGEED* for different values of the parameters as displayed in the key on the graph.

Interpretation: the figure above revealed that the *TOGEED* has a decreasing failure rate which implies that the probability of failure for any random variable following a *TOGEED* decreases as time increases, that is, as time goes on, probability of death decreases.

3.5 Quantile Function

[2] defined the quantile function for any distribution in the form

$$Q(u) = F^{-1}(u) \text{ where } Q(u) \text{ is the quantile function of } F(x) \text{ for } 0 < u < 1$$

Taking $F(x)$ to be the *cdf* of the *TOGEED* and inverting it as above will give us the quantile function as follows:

$$F(x) = 1 - e^{-\alpha(e^{\theta x} - 1)} \left[1 - \lambda + \lambda e^{-\alpha(e^{\theta x} - 1)} \right] = u \quad (3.5.1)$$

Simplifying equation (3.5.1) above, we obtain:

$$Q(u) = X_q = \left\{ \left(-\frac{\ln(1-u)}{2\theta} \right)^{\frac{1}{\alpha}} - 1 \right\} \quad (3.5.2)$$

This function is derived above is used for obtaining some moments like skewness and kurtosis as well as the median and for generation of random variables from the distribution in question.

3.6 Skewness and Kurtosis

This paper presents the quantile based measures of skewness and kurtosis due to non-existence of the classical measures in some cases.

The Bowley's measure of skewness by [3] based on quartiles is given by;

$$SK = \frac{Q\left(\frac{3}{4}\right) - 2Q\left(\frac{1}{2}\right) + Q\left(\frac{1}{4}\right)}{Q\left(\frac{3}{4}\right) - Q\left(\frac{1}{4}\right)} \quad (3.6.1)$$

And the [6] kurtosis is on octiles and is given by;

$$KT = \frac{Q\left(\frac{7}{8}\right) - Q\left(\frac{5}{8}\right) - Q\left(\frac{3}{8}\right) + Q\left(\frac{1}{8}\right)}{Q\left(\frac{6}{8}\right) - Q\left(\frac{1}{8}\right)} \quad (3.6.2)$$

Where $Q(\cdot)$ is obtainable with the help of equation (3.5.2).

3.7 Order Statistics

Let $X_{(1)}$ denote the smallest of $X_1, X_2, \dots, X_n$, $X_{(2)}$ denote the second smallest of $X_1, X_2, \dots, X_n$, and similarly $X_{(i)}$ denote the i^{th} smallest of $X_1, X_2, \dots, X_n$. Then the random variables $X_{(1)}, X_{(2)}, \dots, X_{(n)}$, called the order statistics of the sample $X_1, X_2, \dots, X_n$, has probability density function of the i^{th} order statistic, $X_{(i)}$, as:

$$f_{i:n}(x) = \frac{n!}{(i-1)!(n-i)!} f(x) F(x)^{i-1} [1-F(x)]^{n-i} \quad (3.7.1)$$

Where $f(x)$ and $F(x)$ are the *pdf* and *cdf* of the *TOGEED* respectively.

190 Using (2.1) and (2.2), the *pdf* of the i^{th} order statistics $X_{i:n}$, can be expressed from (3.7.1) as;

$$191 \quad f_{i:n}(x) = \frac{n!}{(i-1)!(n-i)!} \sum_{k=0}^{n-i} (-1)^k \binom{n-i}{k} \left[\alpha \theta e^{\theta x} e^{-\alpha(e^{\theta x}-1)} \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x}-1)} \right] \right] \left[1 - e^{-\alpha(e^{\theta x}-1)} \left[1 - \lambda + \lambda e^{-\alpha(e^{\theta x}-1)} \right] \right]^{i+k-1} \quad (3.7.2)$$

192 Hence, the *pdf* of the minimum order statistic $X_{(1)}$ and maximum order statistic $X_{(n)}$ of the
193 *TOGEED* are given by;

$$194 \quad f_{1:n}(x) = n \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \left[\alpha \theta e^{\theta x} e^{-\alpha(e^{\theta x}-1)} \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x}-1)} \right] \right] \left[1 - e^{-\alpha(e^{\theta x}-1)} \left[1 - \lambda + \lambda e^{-\alpha(e^{\theta x}-1)} \right] \right]^k \quad (3.7.3)$$

195 and

$$196 \quad f_{n:n}(x) = n \left[\alpha \theta e^{\theta x} e^{-\alpha(e^{\theta x}-1)} \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x}-1)} \right] \right] \left[1 - e^{-\alpha(e^{\theta x}-1)} \left[1 - \lambda + \lambda e^{-\alpha(e^{\theta x}-1)} \right] \right]^{n-1} \quad (3.7.4)$$

197 respectively.

198 4 Estimation of Parameters

199 Let $X_1, \dots, X_n$ be a sample of size 'n' independently and identically distributed random
200 variables from the *TOGEED* with unknown parameters α , θ , and λ defined previously. The
201 *pdf* of the *TOGEED* is given as

$$f(x) = \alpha \theta e^{\theta x} e^{-\alpha(e^{\theta x} - 1)} \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x} - 1)} \right]$$

The likelihood function is given by;

$$L(X_1, X_2, \dots, X_n / \alpha, \theta, \lambda) = (\alpha \theta)^n e^{\theta \sum_{i=1}^n x_i - \alpha \sum_{i=1}^n (e^{\theta x_i} - 1)} \sum_{i=1}^n \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x_i} - 1)} \right] \quad (4.1)$$

Let the log-likelihood function, $l = \log L(X_1, X_2, \dots, X_n / \alpha, \theta, \lambda)$, therefore

$$l = n \log \alpha + n \log \theta + \theta \sum_{i=1}^n x_i - \alpha \sum_{i=1}^n (e^{\theta x_i} - 1) + \sum_{i=1}^n \log \left[1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x_i} - 1)} \right] \quad (4.2)$$

Differentiating l partially with respect to α , θ and λ respectively gives;

$$\frac{\partial l}{\partial \alpha} = \frac{n}{\alpha} - \sum_{i=1}^n (e^{\theta x_i} - 1) - 2\lambda \sum_{i=1}^n \left\{ \frac{e^{-\alpha(e^{\theta x_i} - 1)} (e^{\theta x_i} - 1)}{1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x_i} - 1)}} \right\} \quad (4.3)$$

$$\frac{\partial l}{\partial \theta} = \frac{n}{\theta} - \sum_{i=1}^n x_i - \alpha \sum_{i=1}^n \left\{ \frac{x_i e^{-\alpha(e^{\theta x_i} - 1)} e^{-\theta x_i}}{1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x_i} - 1)}} \right\} \quad (4.4)$$

$$\frac{\partial l}{\partial \lambda} = \sum_{i=1}^n \left\{ \frac{2e^{-\alpha(e^{\theta x_i} - 1)} - 1}{1 - \lambda + 2\lambda e^{-\alpha(e^{\theta x_i} - 1)}} \right\} \quad (4.5)$$

Equating equations (4.3), (4.4) and (4.5) to zero and solving for the solution of the non-linear system of equations will give us the maximum likelihood estimates of parameters α, θ , and λ respectively. However, the solution cannot be obtained analytically except numerically with the aid of suitable statistical software like Python, R, SAS, e.t.c when data sets are given.

5 Applications

In this section, we have applied and compared the performance of the Transmuted odd generalized exponential exponential distribution (*TOGEED*) to that of odd generalized

exponential exponential distribution (*OGEED*) and Weibull-exponential distribution (*WED*) using the following dataset.

Data set: This data represents the remission times (in months) of a random sample of 128 bladder cancer patients. It has previously been used by [4] and [11]. It's summarized as follows:

Table 5.1: Summary Statistics for the dataset

parameter	n	Minimum	Q_1	Median	Q_3	Mean	Maximum	Variance	Skewness	Kurtosis
Values	128	0.0800	3.348	6.395	11.840	9.366	79.05	110.425	3.3257	19.1537

From the descriptive statistics in table 5.1, we observed that the data set is positively skewed with a very high coefficient of kurtosis and therefore suitable for flexible and skewed distributions.

To compare the distributions listed above, we have used several measures of model fit such as *AIC* (Akaike Information Criterion), Cram'er-Von Mises (W^*), Anderson Darling (A^*) Kolmogorov smirnov ($K-S$) statistics.

Note that the model with the lowest values of these statistics shall be chosen as the best model to fit the data.

Table 5.2: The statistics ll , *AIC*, A^* , W^* and $K-S$ for the fitted models to the dataset.


Distributions	Parameter estimates	$ll=(\log\text{-likelihood value})$	<i>AIC</i>	A^*	W^*	$K-S$	<i>P-Value</i> ($K-S$)	Ranks
<i>TOGEED</i>	$\bar{\alpha}=0.0182$	416.5186	839.0372	1.0381	0.1747	0.1079	0.1014	1

	$\hat{\alpha}=2.7822$							
	$\hat{\lambda}=0.7591$							
WED	$\hat{\alpha}=0.0293$	546.5931	1099.186	1.4123	0.2357	0.1476	0.0076	2
	$\hat{\alpha}=1.9653$							
	$\hat{\beta}=0.6759$							
OGEED	$\hat{\theta}=0.0346$	439.5273	883.0546	3.2153	0.5463	0.2341	1.6e-06	3
	$\hat{\alpha}=1.6066$							


236

237 It is shown from Table 5.2 above that the Transmuted odd generalized exponential
 238 exponential distribution (*TOGEED*) corresponds to the lower values of ll , AIC , A^* , W^* and K -
 239 S compared to those of the odd generalized exponential-exponential distribution (*OGEED*)
 240 and the Weibull-exponential distribution (*WED*) and therefore we chose the *TOGEED* as the
 241 best model the fits the data.

242 The following figure displayed the histogram and estimated densities and CDF's of the fitted
 243 models for the bladder cancer patients' data set under study.



244



245

246 **Figure 5.1:** Histogram and estimated densities and CDF's of the *TOGEED*, *OGEED* and
247 *WED* for the bladder cancer dataset.

248 From the estimated density plots in figures 5.1 it is very clear that the performance of the
249 *TOGEED* is better than the other two Models.

250 5 Conclusion

In this article, we proposed a new distribution, derived and study some of its properties with graphical analysis and discussion on its usefulness and applications. Hence, having demonstrated earlier in the previous section, we have a conclusion based on our applications of the model to a real life data that the new distribution (*TOGEED*) has a better fit compared to the other two already existing models based on the data set considered in this study.

REFERENCES

- [1] Bourguignon, M., De Andrade, T. A. N. and Cordeiro, G.M. (2015). The Exponentiated Generalized Gumbel distribution. *Revista Colombiana de Estadística*, 38(1): 123 – 143 DOI: <http://dx.doi.org/10.15446/rce.v38n1.48806>
- [2] Hyndman. R. J. and Fan, Y. (1996). Sample quantiles in statistical packages, *The American Statistician*, 50 (4): 361-365.
- [3] Kenney, J. F. and Keeping, E. S. (1962). Mathematics of Statistics, 3 edn, *Chapman & Hall Ltd, New Jersey*.
- [4] Lee, E. and Wang, J. (2003). Statistical methods for survival data analysis, 3rd edn. Wiley, New York
- [5] Maiti, S. S. and Pramanik, S. (2015). Odds Generalized Exponential-Exponential Distribution. *J. of Data Sci.*, 13: 733-754.
- [6] Moors, J. J. (1988). A quantile alternative for kurtosis. *J. of the Royal Stat. Society: Series D*, 37: 25–32.
- [7] Nadarajah, S. (2006). The exponentiated Gumbel distribution with climate application. *Environmetrics*, 17:13–23.
- [8] Oguntunde, P. E. and Adejumo, A. O. (2015). The transmuted inverse exponential distribution. *Int. J. Adv. Stat. and Prob.*, 3(1):1–7

- 274 [9] Oguntunde, P. E., Balogun, O. S., Okagbue, H. I. and Bishop, S. A. (2015). The Weibull-
275 Exponential Distribution: Its Properties and Applications. *J. of Appl. Sci.*. 15(11):
276 1305-1311.
- 277 [10] Owoloko, E. A., oguntunde, P. E. and Adejumo, A. O. (2015). Performance rating of the
278 transmuted exponential distribution: an analytical approach. *Springerplus* 4:818.
- 279 [11] Rady, E., A., Hassanein, W. A. and Elhaddad, T. A. (2016). The power Lomax
280 distribution with an application to bladder cancer data. *SpringerPlus*, 5:1838 DOI
281 10.1186/s40064-016-3464-y
- 282 [12] Shaw, W. T. and Buckley, I. R. (2007). The alchemy of probability distributions: beyond
283 Gram-Charlier expansions and a skew-kurtotic-normal distribution from a rank
284 transmutation map. *Research report*
- 285 [13] Yahaya, A. and Ieren, T. G. (2017). The odd generalized exponential Gumbel
286 Distribution with applications to survival data. *J. of the Nig. Ass. of Math. Physics*, 39:
287 149-158.