

Clinical features and outcome of birth asphyxia in Hôpital du Cinquantenaire of Kisangani: a case-control study

ABSTRACT

Aims: There are few publications available is paucity about of literature/ studies on perinatal asphyxia, one of the three main reasons of newborn deaths in Democratic Republic of Congo (DRC). In this country, newborn mortality is among the highestmost elevated in Africa. This survey aimed to determine prevalence of newborn asphyxia, mortality its lethality and risk factors associated with it. **Study design:** A matched case-control study.

Place and Duration of Study: it was conducted in pediatric service division/ department of Hôpital du Cinquantenaire of Kisangani (HCK), from March 2013 to March 2017 (Kindly mention dates).

Methodology: Clinical and biologic data -parameters of term newborns (prenatal, perinatal and postnatal), hospitalized for neonatal -perinatal asphyxia in the HCK were retrospectively collected and analyzed. Data of dead newborns (cases) were compared to those who survived (controls) to determine risk factors associated with asphyxialetality. One case matched three controls on the basis of sex. {all babies with asphyxia should be considered as cases, rather than babies who died alone. Similarly, control group should be comprised of babies who did not have asphyxia. So there is some serious flaw in design of this study}.

Results:Results. During the period above, 612 newborns were received in the HCK and 146 died. One hundred and nineteen out of 612 had perinatal asphyxia (19.4%) and 26 out of 119 died (17.8% of all newborn deaths and 21.8% of all asphyxia cases). The most frequent perinatal antecedents were premature rupture of the membranes, meconium stained amniotic liquid, pre-eclampsia and eclampsia. In bivariate analysis, factors associated with asphyxia lethality were low birthweight ($P=.02$), reference -referral from another hospital ($P=.01$), age superior more thanto 24 hours ($P=.02$), associated infection sepsis ($P=.003$), asphyxia severity ($P<.001$) and the Stage 3 of Sarnat ($P<.001$).

Conclusion. The frequency of the asphyxia is high in the HCK and its mortality associated with avoidable factors. Networking newborn referral, improving skills of nurses about obstetrical and neonatal emergency cares, and making available equipment for newborn resuscitation can reduce that mortality.

Keywords: Perinatal asphyxia, newborn, risk factors

INTRODUCTION

Up to 7,6 millions of children under 5 years-old deceased during 2010: three million were newborns, (40% of whole deaths). Prematurity, intrapartum complications and sepsis were the main reasons of newborn death [1]. In the world health statistics 2010 published by the WHO, DRC appeared among the 3 countries having the strongest infanto-juvenile mortality, just before Angola and Chad, with a rate of 199 per for one thousand [2]. The Demographic and Health Survey of 2014 found a rate of 104 [3], what was a meaningful reduction: neonatal mortality fell from 42%o in 2007 to 28%o in 2014. However, these rates remain high very elevated and the same report underlines that neonatal mortality varied too little (???) [2]. The poor performance of health worker has multiple determinants ranging from proximal ones such as a lack of knowledge, skill and motivation to distal factors such as disabling working environment in health facilities (poor clinical practices, leadership and supervision; lack of adequate supplies and equipment; health workers' ineffective participation in planning; and lack of peer support) [4]

Formatted: Space Before: 12 pt

Formatted: Underline

Formatted: Underline

Close to 75% of newborn deaths occur in the first week of life, due to prematurity (35%), perinatal asphyxia (23%) and sepsis (23%) [1,4, 6, 7]. For other studies perinatal asphyxia was even the most frequent death reason before the 2 other (Rephrase) [7-9].

Formatted: Underline

It also causes a lot of complications: cerebral motor palsy, psychomotor development impairment, socioeconomic burden for the family. The prevalence of peripartum asphyxia is certainly low in developed countries/ world; while in low-income countries, about one newborn out of 5 newborn is suffers from asphyxiated at birth and 10 to 33% of them die [10-11].

In DRC, the health system encounters several difficulties [3] and the problem of under-equipment for neonatal resuscitation is widespread actual in many hospitals. Only few There is paucity of literature/data on asphyxia are available in the region. Biselele determined some prenatal and post-natal factors associated to mortality [12 (elaborate the need for this study with findings)]. This study was conducted to identify the clinical features and outcome aimed to identify the factors associated to death with the diagnosis of perinatal asphyxia.

Formatted: Underline

2. MATERIAL AND METHODS

2.1 Operationalization of variables

An Perinatal asphyxiated was diagnosed when newborn was all term newborn who had an Apgar score less than 7 at the 5th minutes of life. Asphyxia was light classified as mild if 5th minute Apgar score ranged between 4 and 6. If it ranged between 0 and 3, it was considered as severe. (Reference?)

Hypoxo-ischemic encephalopathy (HIE) was classified according to Levene and Sarnat & Sarnat: Grade 0 for wellbeing newborn, grade 1 or mild light HIE, grade 2 or moderate HIE and grade 3 or HIE severe (Rewrite from literature).

We considered as neonatal infection all newborn who either was symptomatic (fever for example) or asymptomatic associated to pathologic history (premature rupture of membranes, maternal fever in intrapartum) and positive C reactive protein realized after 12-hours old (Rewrite explaining early and late onset neonatal sepsis with diagnostic criteria and reference).

Formatted: Underline

Formatted: Underline

Formatted: Underline

Formatted: Underline

Formatted: Underline

2.2 Study design, sample, inclusion criteria, variables

This was a case-control study. Cases were all asphyxiated newborns who died and controls those who survived. Three controls matched 1 case on the basis of sex. (In a study on asphyxia- all babies with asphyxia should be considered as cases and those without asphyxia should be considered as controls, hence study requires redesigning and further modifications)

Data were collected from medical folders of all newborns hospitalized in the HCK with asphyxia from March 2013 to March 2017. Some were born in HCK and other referred from other hospitals of the city. Transferred cases with asphyxia who died during transportation to HCK were excluded. All premature as well as term newborns with life-threatening congenital malformations and those leading to surgical emergency (anal imperforation, primitive peritonitis, necrotizing enterocolitis) were excluded.

There is no therapeutic hypothermia facilities device of cooling in the HCK. The cases of asphyxia received a symptomatic medical treatment solely, based on oxygen, 10% glucose infusion enriched of trace elements (NaCl, KCl, calcium), phenobarbital if there were seizures.

We studied antenatal features (mode of childbirth, maternal history of pre-eclampsia or eclampsia, placenta praevia, cord proclence, dystocic presentation) and postnatal (meconium stained ial amniotic liquid fluid, Apgar at 1st and 5th minutes, clinical signs in the following hours, duration of hospitalization, mortality within 24 and 48 hours).

Treatment and analysis of Statistical data were realized analysed by Microsoft Excel® 2016 and Epi info™ 7.1.5. Pearson's chi square and odds ratio were used to check out analyse association between variables determine risk factors. The student's t- test of Student was used for comparing means of quantitative data.

3. RESULTS AND DISCUSSION

3.1 Socio-demographic and clinical data

86 From March 2013 to March 2017, 612 newborns were hospitalized in to the HCK, among whom 119
 87 were asphyxiated (19,4% of all newborn hospitalizations). One hundred and forty-six died for
 88 miscellaneous causes, including 26 with perinatal asphyxia (17.8% of all newborns deaths and 21.8%
 89 of asphyxia lethality).

90
91
92
93
94
95
96
97
98
99

Table 1. Epidemio-clinical features of newborns with perinatal asphyxia

		Frequency	%
Mean age	1.7 ± 3 days (median 1 day)		
Home address	Kabondo	24	(20.17%)
	Kisangani	6	(5.04%)
	Makiso	73	(61.34%)
	Mangobo	8	(6.72%)
	Tshopo	8	(6.72%)
Birthweight	<2500 g	16	(13.45%)
	≥2500 g	103	(86.55%)
		Mean birthweight 3120.1 ± 551.2 g (median 3095 g)	

100

101 The sex ratio F/M was 0.7 (52/67).Most of newborns were male and ≤1 day old. Weight varied
 102 between 1845 and 4250 g.

103 Table 2. Other clinical features of asphyxiated newborns

104

		freq	%
Clinical signs	Respiratory distress	59	(49.4%)
	Pallor	26	(22.1%)
	Lethargy	20	(16.9%)
	Seizures	19	(15.6%)
	Fever	19	(15.6%)
	ExcessiveEndless crying	15	(13%)
	GIDigestive hemorrhage	9	(7.8%)
	Coma-Unconsciousness/ cerebral depression	5	(4 %)

Perinatal features	Meconium stained amniotic fluid	43	(36.4%)
	<u>Cesarean section (not a risk factor for asphyxia)</u>	31	(25.9%)
	Pre-eclampsia/eclampsia	26	(22.1%)
	Premature rupture of membranes	22	(18.1%)
	<u>Too long-lasting</u> <u>Prolonged</u> labor	14	(11.6%)
	Dystocic presentations	8	(6.5%)
	<u>Twins (not a risk factor for asphyxia, unless complicated with some associated factor)</u>	5	(4%)
Origin	Born in HCK	58	(48.74%)
	Born then referred from other hospitals	61	(51.26%)
Fatal outcome	All deaths among asphyxiated newborns	26	(21.85 %)
	Deaths <u>Mortality</u> during the first 24 hours (n=26)	20	(77.6%)

Formatted: Underline

Formatted: Font: 10 pt, Not Bold, Underline

Formatted: Underline

Formatted: Underline

Respiratory disorders wasere the most common clinical feature at admissionobserved. In perinatal history, meconium stained amniotic fluid wasere the most common antecedent factorobserved, followed by premature rupture of membranes.

About HIE, we had 13 cases with grade 0 (healthy newborns), 18 cases with grade 1, 33 cases with grade 2 and 17 with grade 3. The mean hospitalization duration was 7.6 ± 4.2 days (median 5 days)

3.2 Analytic data (N =368)

Table 3. Factors associated with mortality

		Deceased	Alive	OR* (CI† 95%)	P
Birthweight	< 2500 g	8	9	3.4 (1.1 – 10.3)	.02
	> 2500 g	19	84		
Sex	Female	11	41	.9 (.3 – 2.2)	.8
	Male	15	52		
Origin	Referred	19	42	3.2 (1.2 - 9)	.01
	HCK	7	51		
Age at admission	≤ 1 day	17	79	2.9 (1.1 - 8)	.02
	> 1 day	9	14		
Sepsis	Yes	15	25	3.7 (1.5 – 9.1)	.003
	No	11	68		
Asphyxia severity	Light	7	81	.05 (.01 – .1)	< .001

	Severe	19	12	
HIE Sarnat grade	0	1	44	< .001
	1	4	28	
	2	9	19	
	3	12	2	

OR: odds ratio; CI: confidence interval.

Deceased newborns had a lower birthweight (2912.3 ± 609.8 g) than those who survived (3178.2 ± 522.5 g): $t(117) = -2.21$; $P = .02$). The low birthweight had a three-fold risk of dying with asphyxia than those who had a normal birthweight.
Referral from other hospitals, arriving at HCK at an age older than 1 day, sepsis, severe asphyxia, Sarnat grade 2 and 3 were **associated with asphyxia related at higher risk of** mortality.

3.3 Discussion

The majority of our newborns were male, from Makiso township (Table 1). The male predominance was **found reported** by other authors **as well** who found a meaningful difference in favor of the boys [13,14]. Tina Katamea in DRC found that the neonatal mortality was associated to masculine sex [15] but there were no results about association of perinatal asphyxia and mortality related to it. **(any possible expaination for mortality in males should be added)**

The peripartum asphyxia represented 19.4% of all newborns hospitalizations in to the HCK and 17.8% of **the reasons of all** neonatal mortality. This rate was higher than the 10.94% found in India [10], but close to the 18% found in Cameroon [11].

Mortality related to perinatal asphyxia (table 2) was higher than 10% found in India [10], **18.5%** in Nigeria [7] and 21% in Ghana [9]. It was lower than 32% in Liberia [8], 30% in Nepal [16], 25% in Kenya [6] and Malawi [17].

Permanence of physicians and nurses at any moment, availability of resuscitation kit (suction device, ambu bag, masks) and oxygen for neonatal resuscitation contributed for many to these results. Nevertheless, the prevalence remains high and can be due to an elevated proportion of newborns referred from other hospitals for lack of resuscitation devices. (Rewrite) They often arrived too late [18] and found no cooling device available in HCK.

Low birthweight, reference from another hospital, age **more than superior to** 24 hours, severe asphyxia, **elevated higher** Sarnat grade (Table 3) were factors associated **to with** mortality related to peripartum asphyxia. These data agree with those of the other previous studies: the low birthweight newborn is more fragile and at high risk of developing hypoglycemia.


The Sarnat grade describes severity of HIE: higher grades had more severe HIE. Therefore, in this study, like in many others, mortality **was** associated to the **more elevated higher** grades [10-12, 18, 20].

Reference from other **motherhoods hospital** and age at admission were evident risk factors [7]. For the best moment to avoid asphyxia is the period of the **4** first minutes **(Golden minute)**. If asphyxia could not be avoided, the first 6 hours of life were the most important time to use **cooling device therapeutic hypotheria** in order to prevent HIE and neurologic damage [19]. This shows how bad can be the prognosis of newborn arriving later than 24 first hours. No cooling device were available in HCK. So, only children born there could be immediately resuscitated and that explains the significant difference between them and those referred from other hospitals.

4. CONCLUSION

The mortality rate of peripartum asphyxia was high in HCK and associated to avoidable and preventable factors. Networking newborn referral, improving skills of nurses in the motherhood staffs of hospitals in Kisangani about emergency obstetric and neonatal care, and providing to them devices **for neonatal resuscitation are necessary interventions to reduce mortality due to peripartum asphyxia.**

Formatted: Underline



ETHICAL APPROVAL

This study received the agreement of Research authorities of Faculty of Medicine and Pharmacy of University of Kisangani.

REFERENCES

1. Li Liu, Hope I J, Cousens S, Perin J, Scott S, Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000, *Lancet* 2012; 379: 2151–61
2. WHO, world health statistics 2010, Geneva, 2010
3. Ministère du plan et suivi de la mise en œuvre de la révolution de la modernité (mpsmrm), ministère de la santé publique (msp) et ICF international, 2014. *Enquête démographique et de santé en république démocratique du Congo 2013-2014*. Rockville, maryland, usa : mpsmrm, msp et ICF international
4. Brantuo M N, Cristofalo E, Mehe M, Ameh J, Okai B N, Evidence-based training and mentorship combined with enhanced outcomes surveillance to address the leading causes of neonatal mortality at the district hospital level in Ghana, *tropical medicine and international health*. 2014, 19:4, p 417–426,
5. Ashish KC, Målqvist M, Wrammert J, Verma S, Raj arya I D, Implementing a simplified neonatal resuscitation protocol-helping babies breathe at birth (hbb) at a tertiary level hospital in Nepal for an increased perinatal survival, *bmc pediatrics* 2012, 12:159
6. Yego F, Stewart W J, Byles J, Nyongesa P, Aruasa W and D'este D, A retrospective analysis of maternal and neonatal mortality at a teaching and referral hospital in Kenya, *Reproductive health* 2013, 10:13
7. Ekwochi U, Ndu IC, Nwokoye, Ezenwosu, of Amadi, Osuorah D, Pattern of morbidity and mortality of newborns admitted into the sick and special care baby unit of Enugu state university teaching hospital, Enugu state, *Nigerian journal of clinical practice*, 2014, 17:3,
8. Jody R. L, Facnm, Ffaan, a case series study of perinatal deaths at one referral center in rural post-conflict Liberia, *matern child health j*. 2014 January; 18(1)
9. Welaga P, Moyer C, Aborigo R, Adongo P, Williams J et al. Why are babies dying in the first month after birth? A 7-year study of neonatal mortality in northern Ghana. *Plos one*, 2013, 8(3): e58924
10. Kumar A, Suruchi S, Vimalk, Apeksha P, Deba P B, Clinical profile and short-term outcome of hypoxic ischemic encephalopathy among birth asphyxiated babies in Katihar medical college hospital, *journal of clinical neonatology*, 2014; 3(4)
11. Monebenimp F, Tietche F, Eteki N, Asphyxie néonatale au centre hospitalier et universitaire de Yaoundé, Cameroon, *clinics in mother and child health*, 2005; 2(2): 335-338
12. Biselele T, Naulaers G, Bunga Muntu P, Nkidiaka E, Kapepela M et al, A descriptive study of perinatal asphyxia at the university hospital of Kinshasa (Democratic Republic of Congo), *J Trop Pediatr*, 2013; 59 (4): 274-279
13. Mohamed M, Aly H, impact of race on male predisposition to birth asphyxia, *j perinatol*. 2014; 34(6): 449-52
14. Eghbalian F, frequency of hypoxic-ischemic encephalopathy among hospitalized neonates in west Iran, IRAN, *JPediatr*, 2010; 20 (2), 244-245,
15. Tina K, Mukuku O, Kamona I, Mukelenge K, Mbula O, Facteurs de risque de mortalité chez les nouveaux-nés transférés au service de néonatalogie de l'hôpital Jason Sendwe de Lubumbashi, République Démocratique du Congo, *pan african medical journal*. 2014 ; 19 :169,
16. Lee C, Mpha M, Mullany L, James M, Tielsch, Katz, Risk factors for neonatal mortality due to birth asphyxia in southern Nepal: a prospective, community-based cohort study, *Pediatrics*. 2008; 121(5): e1381–e1390

- 223 17. Michael K H, Olmsted, K, Kiromera A, Chamberlain, a neonatal resuscitation curriculum in
224 Malawi, Africa: did it change in-hospital mortality? *International journal of pediatrics*,
225 2012, 10:1155
- 226 18. Memon S, Shaikh S, Bibi S. To compare the outcome (early) of neonates with birth
227 asphyxia in-relation to place of delivery and age at time of admission. *J pak med assoc*.
228 2012;62(12):1277-81
- 229 19. Daetwyler K, Brotschib B, Berger T, Wagner P, Feasibility and safety of passive
230 cooling in a cohort of asphyxiated newborn infants, *Swiss med wkly*. 2013;143: 13767
- 231 20. Qureshi AM, Rehman UR, Siddiqi T, Hypoxic ischemic encephalopathy in neonates, *J*
232 *Ayub Med Coll Abbottabad*. 2010;22(4):190-3.