

Measurement of Natural Radionuclides (^{40}K , ^{238}U and ^{232}Th) and Associated Dose Rates in Soil and Commonly Consumed Foods (Vegetables and Tubers) at Okitipupa, Ondo State, Southwestern Nigeria.

ABSTRACT

The naturally occurring radioactive materials (NORMs) are known as the largest sources of exposure to human. Thus, the aim of this study is to collect soil and some commonly consumed food materials in Ondo State, Nigeria such as tubers (cassava, *Manihot esculent* and yam, *Dioscorea alata*) samples and vegetables (waterleaf, *Talinum triangulae* and bitter leaf, *Vernonia amygdalina*) samples at some selected locations in Okitipupa, Ondo state, Southwestern, Nigeria in order to determine following natural radionuclides (^{40}K , ^{238}U and ^{232}Th) levels using a well calibrated NaI(Tl) that is well shielded with a detector couple to a computer resident quantum MCA2100R Multichannel. The transfer factors, annual absorbed dose rate and the annual effective dose in the samples collected were estimated.

The results showed that the measured natural radionuclides were present in the mean concentrations of $323.79 \pm 12.45 \text{ Bqkg}^{-1}$, $81.87 \pm 45.30 \text{ Bqkg}^{-1}$ and $57.62 \pm 18.04 \text{ Bqkg}^{-1}$ for ^{40}K ; $11.76 \pm 36.03 \text{ Bqkg}^{-1}$, $4.67 \pm 10.12 \text{ Bqkg}^{-1}$ and $3.45 \pm 2.10 \text{ Bqkg}^{-1}$ for ^{238}U and $9.66 \pm 0.89 \text{ Bqkg}^{-1}$, $3.07 \pm 2.45 \text{ Bqkg}^{-1}$ and $2.45 \pm 0.92 \text{ Bqkg}^{-1}$ for ^{232}Th for soil, yam and cassava samples respectively. The results also showed that the radionuclides were present in the concentrations of $11.76 \pm 36.03 \text{ Bqkg}^{-1}$ and $9.66 \pm 0.89 \text{ Bqkg}^{-1}$ for ^{40}K ; $9.67 \pm 8.53 \text{ Bqkg}^{-1}$ and $7.87 \pm 1.89 \text{ Bqkg}^{-1}$ for ^{238}U and $8.63 \pm 6.08 \text{ Bqkg}^{-1}$ and $6.58 \pm 0.76 \text{ Bqkg}^{-1}$ for ^{232}Th for water leaf and bitter leaf samples respectively. The soil-to-yam transfer factors were found to be 0.26, 0.40 and 0.32 for ^{40}K , ^{238}U and ^{232}Th and soil-to-cassava yam transfer factors were found to be 0.18, 0.29 and 0.25 for ^{40}K , ^{238}U and ^{232}Th respectively. The soil-to-water leaf transfer factors were found to be 0.37, 0.82 and 0.82 for ^{40}K , ^{238}U and ^{232}Th and the soil-to-bitter leaf transfer factors were found to be 0.32, 0.74 and 0.68 for ^{40}K , ^{238}U and ^{232}Th respectively.

The mean absorbed dose rate was $25.08 \pm 0.57 \text{ nGyh}^{-1}$ and the mean annual outdoor effective dose was $46,166.96 \mu\text{Svy}^{-1}$. The annual effective dose reported for this present study area represents 65.95% of the world average value of $70,000 \mu\text{Svy}^{-1}$ and 47.11% of Nigeria value of $98000 \mu\text{Svy}^{-1}$ 

Keywords: *Natural radionuclides, transfer factor, absorbed dose rate, annual outdoor effective dose, Nigeria.*

1.0 Introduction

The environment is constituted by various materials containing different forms and amount of natural radionuclides and their decayed products. The isotopes of Potassium (^{40}K), Uranium (^{238}U) and Thorium (^{232}Th), and their daughter products can be found abundantly in the outer layers of the Earth ⁽¹⁾. The radioactivity and composition of these natural radionuclides will be studied in soil, tubers and vegetables for this present study. Soils with more than average

level of radiation can be found in many regions due to the exposure to cosmic rays from the outer space ⁽²⁾. Knowing the level of radiation in the environment is important in implementing a suitable control for the sake of radiological protection and radiation exposure to a large population with dose about 1.5Sv increases cancer incidence and mortality ⁽³⁾. Thus, all foods are slightly radioactive and all food sources combined exposes a person to about 0.4 mSv per year ⁽³⁾. Tubers and vegetables are very vital in human diet, the presence of natural radionuclides (⁴⁰K, ²³⁸U and ²³²Th) in them may results in radiological implication not only in the food but also on the populace consuming these foods. Tubers are cultivated in nearly all tropical regions of the world providing a staple starchy food for 80 million people and as important source of income ⁽⁴⁾. Human beings have always been exposed to ionizing radiations of natural origin, namely terrestrial and extraterrestrial radiation. Radiation of extra-terrestrial origin is from high energy cosmic ray particles and at sea level it is about 30 nGyh⁻¹ ⁽³⁾, while that of terrestrial origin is due to the presence of naturally occurring radionuclides; mainly potassium, rubidium and the radionuclides in the decay chains of thorium and uranium ⁽⁵⁾. These radionuclides have half-life which is comparable to the age of the earth ⁽⁶⁾. The mandate to control nuclear and radiation generating sources in Nigeria is vested with the Nigerian Nuclear Regulatory Authority which is authorized by law to ensure that radiation protection and safety regulations are adhered to several studies have been conducted around the world to assess natural radioactivity levels in the soil of certain areas ⁽⁷⁾.

Agriculture has been one of the backbones of the economy of many developing countries like Nigeria. Many countries in Africa have laid down policies on the provision of sustainable food security. When people have sufficient food to eat, many of the nutrition-related problems are avoided and healthy citizen are available to work for the growth of respective countries. It is expected that developing countries do not only concentrate on the provision of adequate food for their citizens but also food that is chemically and radiologically safe ⁽³⁾.

An important goal of the United Nations (UN) relating to sustainable food security is to assist members' states in ensuring that people have access to food that is sufficient, nutritionally adequate and above all considerably safe for human consumption ⁽⁸⁾. The presence of radionuclide in soil above a certain threshold leads to contamination of food products since these food products derived their nutrients for growth from the top soil on which they are grown. Radionuclides in solution can then be incorporated through root hair and then to the root of vegetables for onward transfer to the leave system of the vegetables that human feed on. In most cases, this is facilitated by their chemical similarity with other element that the grasses usually depend on for growth ⁽⁸⁾.

The absorption of radionuclides associated with particles can be in a form of direct dietary intake or an indirect process following particle weathering and soil-grass-animal transfer. This is dependent on soil water pH, organic matter, microbial activities and the vegetation present ⁽⁸⁾. It is therefore very important to assess radiological safety of these tubers and vegetables consumed by human, this is because the ingestion of foods contaminated with radionuclides has the potential of exposing human beings to high level of radiation dose. Furthermore, radionuclides with relatively long half-lives are considered human health risk as they can get into the human system through the food chain and thereby increase the radiation burden for many years ⁽²⁾.

The natural radioactivity levels in soils of some locations in Nigeria, with Benin formation underlay, have been reported to be significantly higher than the world average ⁽⁷⁾. The radioactivity levels in vegetables and tubers grown in such soils may be enhanced through the plant-root uptake of radionuclides and such, humans feeding on the food products may have high retention of natural radionuclides in their body causing the concentrations of several pollutants, radioactive or otherwise, including Uranium, Thorium and Potassium, have


increased over the last few decades ⁽⁸⁾. These elemental pollutants do exist in the environment and are a threat to many organisms. Compared to other physical and biological end points, the effects of pollutants such as radionuclides on human have been the focus of only a few studies. Therefore, it is necessary to measure the activity concentrations of the natural radionuclides in the soil samples and food products grown in the study area. Thus, the aim of this study is to collect soil samples and commonly consumed foods (waterleaf, bitter leaf, cassava and yam) at some locations in Okitipupa, Ondo state, Nigeria in order to determine the natural radioactivity levels in the samples with the objectives to measure the activity concentrations of ⁴⁰K, ²³⁸U and ²³²Th radionuclides in all the samples collected, determine the transfer factors (TFs), estimate the annual absorbed dose rate in the samples collected and estimate the annual effective dose.

2.0 Materials and Methods

2.1 Geographical description of the study area

Okitipupa is a Local Government Area in Ondo State, Nigeria. It is native to the Ikales, who are a sub-set of the larger Yoruba tribe. It has always been the central town for inhabitants of the Ondo South senatorial district of Ondo state comprising okitipupa, Irele, Ilaje, Ese-Odo, Odigbo and Ile-oluji/Oke-Igbo local governments due to the presence of several amenities. It was a district in the colonial days before Nigeria's independence in 1960. It has a university, a Specialist hospital, several private hospitals, a Magistrate court, a High court, a Police division, an Army Base, Commercial banks, a Telephone exchange, numerous primary and secondary schools.

The Ikale natives are predominantly farmers. The major cash crop being cultivated in the area are oil-palm, rubber and cassava. They also cultivate yam, beans, okro, pepper, melon and vegetables. Staple food includes but is not limited to baked cassava popularly known as pupuru, yam, rice, yam flour and cassava flakes (Garri) among others. Major industries located in the town include the Okitipupa Oil Palm Plc and Oluwa Glass Factory. Palm-oil and rubber plantations litter the landscape. It has an area of 803 km², coordinates 6°30'0" North, 4°48'0" East and a population of 233,565 ⁽⁹⁾.



Map 1: Geological map of the study area (Okitipupa)

2.2 Samples collection

10 samples were collected each for soil, cassava, yam, water leaf and bitter leaf respectively for the present study, these points of collection were marked out using a Global Positioning System (GPS). Soil samples were collected from depths 5cm the different locations within the sample site while all food samples were collected directly from farm lands in the study area. At the collection point, all samples were wrapped in separate black plastic bags and were well labelled with a paper masking tape. The samples were then transported to the laboratory for preparation.

2.3 Samples preparation

Soil samples were well mixed after removing extraneous materials such as roots, pieces of stones and gravels. Samples were then weighted and dried into an electric oven at 110°C for 4 days until a constant dry weight was obtained. The dried samples were then crushed using mortar and pestle. After crushing and mixing thoroughly, soil samples were shaken in a sieve shaker using a 2mm mesh size. The samples were later scaled in 200g each in a radon tight container for a minimum of 28days so as to reach secular equilibrium between radon and its daughter nuclides before radiometric counting ⁽¹⁾.

The vegetable samples were washed thoroughly in order to remove dust and surface contaminations. They were then dried in an oven at 110°C for 24 hours and placed in a furnace. The samples were allowed to ash at this temperature for 24 hours. The ash samples were cooled to room temperature. The cooled ash samples were then weighed ⁽⁴⁾. The samples were later scaled into 200 g each in radon tight containers for a minimum of 28days so as to reach secular equilibrium between radon and its daughter nuclides before radiometric counting ⁽¹⁰⁾.

Cassava and yam samples were rinsed to remove sands and dusts so as to avoid contaminations, they were then sliced into bits respectively, sun dried and then oven dried at 80°C until a constant weight is reached. Each sample were crushed and sieved using sieve

mesh (2mm) and kept in air-tight containers and sealed. The crushing and sieving were crucial for achieving homogeneous state of the sample. The samples were later scaled in 200g each in a radon tight container for a minimum of 28days so as to reach secular equilibrium between radon and its daughter nuclides before radiometric counting.

2.4 Activity determination

The activity concentrations of the soil samples were measured using a well calibrated NaI(Tl) and well shielded detector couple to a computer resident quantum MCA2100R Multichannel analyzer for 36,000s. An empty container under identical geometry was also counted for the same time. The 1460KeV gamma-radiation of ^{40}K was used to determine the concentration of ^{40}K in the sample. The gamma transition energy of 1764.5KeV ^{214}Bi was used to determine the concentration of ^{238}U while the gamma transition energy of 2614KeV ^{208}Tl was used to determine the concentration of ^{232}Th while ^{137}Cs was detected by its 661.6KeV gamma transition.

$$C_s = \frac{C_\alpha}{P_\gamma \left(\frac{M_s}{V_s}\right) \varepsilon_\gamma t_c} (BqKg^{-1}) \quad (1)$$

Where C_s is the sample concentration, C_α is the net peak energy, ε_γ is the efficiency of the detector for a γ -energy of interest, M_s/V_s is the sample mass per volume of soil, t_c is the total counting time and P_γ is abundance of the γ -line in a radionuclide.

The efficiency calibration of the detector was done using a reference standard mixed source traceable to Analytical Quality Control Service (AQCS, USA), which has certified activities of the selected radionuclide and has a geometrical configuration identical to sample container. The standard sources contained ten known radionuclides. The energy calibration was also performed by using the peaks of the radionuclide present in the standard sources. The channel number is proportional to energy; the channel scale was then converted to an energy scale. This produces an energy calibration curve, i.e. energy versus channel.

The detection limit (DL) of a measuring system describes its operating capability without the influence of the sample. The DL is given in $Bqkg^{-1}$ which is required to estimate the minimum detectable activity in a sample. This was obtained using the equation:

$$DL (Bqkg^{-1}) = \frac{1.96 \left(\frac{B}{T} + SD^2\right)^{1/2}}{k \times \varepsilon \times m} \quad (2)$$

Where SD is the estimated standard error of the net background count rate in the peak, T is the counting time (s), ε is the counting efficiency (cps/Bq), m is the mass of the sample, k is the factor that converts cps (counting per seconds) to Bq and 1.96 represents the 95% confidence level.

2.5 Transfer factor

The soil-to-crop transfer factors (TFs) measured the transfer of radionuclides from the soil to crops. In soil, each radioactive element follows complex dynamics in which a part of its concentration is transported into the soil solution, while another part gradually becomes strongly bound to the particles of the soil. The portion of these radionuclides, which is in the soil solution, can be incorporated via the root into the grass⁽¹¹⁾.

195 From observed activity concentrations of the radionuclides in the food crops and in the
196 corresponding soil, the transfer factors (TFs) values were calculated according to the
197 equation:

$$198 \quad TF = \frac{A_g}{A_s} \quad (3)$$

199 Where: A_g = Activity of radionuclides in crops ($BqKg^{-1}$ dry weight)

200 A_s = Activity of radionuclides in soil ($BqKg^{-1}$ dry weight)

201 The dry weight was preferred because the amount of radioactivity per kilogram dry weight is
202 much less variable than the amount per unit fresh weight. The soil-to-crop TF can be used as
203 an index for the accumulation of trace elements by crops or the transfer of elements from the
204 soil to the crop.

205

206 **2.6 Absorbed dose rate**

207 The estimated absorbed dose rate in $nGyh^{-1}$ due to the radionuclide concentration was done
208 according to the equation ⁽¹²⁾:

$$209 \quad D = 0.042S_K + 0.429S_U + 0.666S_{Th} \quad (4)$$

210 Where D is the absorbed dose rate in $nGyh^{-1}$ due to the specific radionuclide concentration
211 S_K , S_{Th} and S_U for ^{40}K , ^{232}Th and ^{238}U respectively in $Bqkg^{-1}$ at 1m above the ground.

212

213 **2.7 Annual outdoor effective dose**

214 Effective dose is a measure of the stochastic effects on health risk due to radiation dose either
215 internal or external to whole or part of the body. The annual outdoor effective dose equivalent
216 was estimated using outdoor occupancy factor of 0.3 and conversion factor of $0.7 SvGy^{-1}$ ⁽³⁾.
217 The 0.7 conversion factor shows that an average person stays about 7 hours outside daily.

$$218 \quad E_x = D_x \times N(h) \times O_f \times K_f \quad (5)$$

219 E_x is the annual outdoor effective dose ($\mu Sv y^{-1}$), $N(h)$ is the number of hours in a year (24
220 hours x 365.24 days), D_x is absorbed dose rate in the air ($nGyh^{-1}$), O_f is outdoor occupancy
221 factor and K_f is conversion factor ($SvGy^{-1}$).

222 **3.0 Results and discussions**

223 **3.1 Activity concentrations in the soil samples**

224 In the study, the radioactivity level in the soil samples ranged from 290.09 to 395.22 $Bqkg^{-1}$
225 with a mean value of $323.79 \pm 12.45 Bqkg^{-1}$ for ^{40}K ; 10.19 to 12.35 $Bqkg^{-1}$ with a mean value
226 of $11.76 \pm 36.03 Bqkg^{-1}$ for ^{238}U and 8.42 to 10.93 $Bqkg^{-1}$ with a mean value of 9.66 ± 0.89
227 $Bqkg^{-1}$ for ^{232}Th . The activity concentrations of ^{40}K , ^{238}U and ^{232}Th radionuclides in the soil
228 samples were higher than the corresponding values in food samples, this affirmed the general
229 assertion as that only a fractional part of the radionuclides in the soil is transferable to the
230 plants ⁽¹³⁾. The activity concentrations of ^{40}K , ^{238}U and ^{232}Th were lower than the world
231 average values of 410.0 $Bqkg^{-1}$ for ^{40}K ; 35.0 $Bqkg^{-1}$ for ^{238}U and 28.0 $Bqkg^{-1}$ for ^{232}Th ⁽³⁾.

The values were also lower than the mean activity concentration values of $39.24 \pm 1.12 \text{ Bqkg}^{-1}$, $52.86 \pm 1.40 \text{ Bqkg}^{-1}$ and $445.02 \pm 12.24 \text{ Bqkg}^{-1}$ for ^{238}U , ^{232}Th and ^{40}K respectively earlier reported for some food crop producing region in Ondo State ⁽¹⁴⁾.

235

Table 1: Activity concentrations of ^{40}K , ^{238}U and ^{232}Th in soil samples from the study area

Samples	GPS LOCATIONS		SOIL		
	North	East	^{40}K (Bqkg^{-1})	^{238}U (Bqkg^{-1})	^{232}Th (Bqkg^{-1})
S1	06°27'21.2"	004°27'28.4"	321.16±13.08	10.19±33.11	9.13±0.93
S2	06°27'28.4"	004°45'46.5"	290.09±25.04	10.90±35.14	10.03±1.25
S3	06°26'26.2"	004°46'11.8"	312.38±13.91	11.61±30.65	10.93±2.45
S4	06°26'28.6"	004°46'11.0"	305.38±3.28	11.82±42.17	9.68±0.80
S5	06°26'27.9"	004°46'11.2"	298.37±8.41	12.03±38.01	8.42±1.20
S6	06°26'27.1"	004°48'11.1"	322.58±13.21	12.11±29.98	8.82±3.69
S7	06°26'25.0"	004°47'11.8"	346.80±4.39	12.19±33.09	9.23±2.20
S8	06°26'27.2"	004°42'31.1"	395.22±2.11	12.35±30.13	10.03±1.98
S9	06°27'30.1"	004°41'27.3"	347.06±14.31	12.26±28.20	10.13±0.91
S10	06°26'27.5"	004°46'62.6"	298.90±8.29	12.17±43.41	10.22±1.19
Range			290.09-395.22	10.19-12.35	8.42-10.93
Mean			323.79±12.45	11.76±36.03	9.66±0.89

237

3.2 Activity concentrations in water leaf (*Talinum triangulae*) and bitter leaf (*Vernonia amygdalina*) samples

In the study, the radioactivity level in the water leaf samples ranged from 104.60 to 135.40 Bqkg^{-1} with a mean value of $118.96 \pm 22.43 \text{ Bqkg}^{-1}$ for ^{40}K ; 8.43 to 10.40 Bqkg^{-1} with a mean value of $9.67 \pm 8.53 \text{ Bqkg}^{-1}$ for ^{238}U and 6.98 to 8.92 Bqkg^{-1} with a mean value of $7.87 \pm 1.89 \text{ Bqkg}^{-1}$ for ^{232}Th . ^{40}K activity concentration exhibited highest values among other radionuclides measured for all samples. The radioactivity level in the bitterleaf samples ranged from 97.34 to 110.70 Bqkg^{-1} with a mean value of $103.26 \pm 6.08 \text{ Bqkg}^{-1}$ for ^{40}K , 7.80 to 9.50 Bqkg^{-1} with a mean value of $8.63 \pm 3.45 \text{ Bqkg}^{-1}$ for ^{238}U and 5.97 to 7.03 Bqkg^{-1} with a mean value of $6.58 \pm 0.28 \text{ Bqkg}^{-1}$ for ^{232}Th . In the present study, the value of ^{238}U was higher than the value of 1.30 Bqkg^{-1} reported for vegetable samples (tomatoes) in Cameron Highlands and Penang ⁽¹⁵⁾. The values for ^{40}K and ^{232}Th were lower than the values of 132.79 Bqkg^{-1} and 47.33 Bqkg^{-1} reported for carrot samples in Barkin Ladi LGA., Plateau State-Nigeria ⁽¹⁶⁾.

In the present study, the values of ^{238}U and ^{232}Th were higher than the values of 6.25 Bqkg^{-1} and 0.41 Bqkg^{-1} respectively reported for vegetable samples in Cameron Highlands and Penang ⁽¹⁵⁾. The mean value of ^{40}K is lower than the mean value of 125.17 Bqkg^{-1} reported for

vegetable samples in Barkin Ladi LGA., Plateau State-Nigeria ⁽¹⁶⁾. The activity concentrations in waterleaf samples were higher than the values in bitter leaf samples reported in the present study.

Table 2: Activity concentrations of ⁴⁰K, ²³⁸U and ²³²Th in waterleaf and bitter leaf samples

Samples	waterleaf			bitter leaf		
	⁴⁰ K (Bqkg ⁻¹)	²³⁸ U (Bqkg ⁻¹)	²³² Th (Bqkg ⁻¹)	⁴⁰ K (Bqkg ⁻¹)	²³⁸ U (Bqkg ⁻¹)	²³² Th (Bqkg ⁻¹)
S1	120.34±20.12	10.00±8.30	7.50±1.80	110.70±6.00	9.50±3.40	6.70±0.28
S2	119.07±18.37	9.22±4.11	8.21±2.11	108.15±6.99	8.91±3.59	6.87±0.22
S3	117.80±25.22	8.43±7.44	8.92±3.21	105.60±7.20	8.32±3.10	7.03±1.45
S4	126.60±24.18	9.42±2.62	8.29±0.92	101.47±5.46	8.41±3.09	6.77±2.03
S5	135.40±23.42	10.40±3.00	7.65±0.99	97.34±12.29	8.50±4.28	6.50±0.98
S6	127.70±14.99	10.21±3.50	7.48±1.70	98.63±4.01	8.63±2.98	6.37±0.86
S7	120.00±18.91	10.01±4.50	7.32±1.68	99.92±3.99	8.76±3.45	6.24±1.40
S8	104.60±21.25	9.62±3.90	6.98±1.99	102.50±6.22	9.02±3.07	5.97±1.28
S9	107.55±27.00	9.66±2.11	7.79±1.86	103.60±6.81	8.41±3.00	6.43±0.85
S10	110.50±24.10	9.70±1.69	8.60±1.33	104.70±6.01	7.80±2.45	6.89±2.04
Range	104.60-135.40	8.43-10.40	6.98-8.92	97.34-110.70	7.80-9.50	5.97-7.03
Mean	118.96±22.43	9.67±8.53	7.87±1.89	103.26±6.08	8.63±3.45	6.58±0.76

3.3 Activity concentrations in yam (*Dioscorea alata*) and cassava (*Manihot esculenta*) samples

In the study, the radioactivity level in the yam samples ranged from 65.20 to 97.50 Bqkg⁻¹ with a mean value of 81.87±45.30 Bqkg⁻¹ for ⁴⁰K; 3.92 to 5.22 Bqkg⁻¹ with a mean value of 4.67±10.12 Bqkg⁻¹ for ²³⁸U and 2.51 to 3.60 Bqkg⁻¹ with a mean value of 3.07±2.45 Bqkg⁻¹ for ²³²Th. ⁴⁰K activity concentration exhibited highest values among other radionuclides measured for all samples. In the present study, the values of ⁴⁰K and ²³²Th were higher than the values of 41.59 Bqkg⁻¹ and 52.69 Bqkg⁻¹ while the values of ²³⁸U is lower than the value of 14.02 Bqkg⁻¹ reported for yam samples in Barkin Ladi LGA., Plateau State-Nigeria ⁽¹⁶⁾.

The radioactivity level in the cassava samples ranged from 45.34 to 69.42 Bqkg⁻¹ with a mean value of 57.62±18.04 Bqkg⁻¹ for ⁴⁰K; 2.50 to 3.95 Bqkg⁻¹ with a mean value of 3.45±2.10 Bqkg⁻¹ for ²³⁸U and 2.06 to 3.06 Bqkg⁻¹ with a mean value of 2.45±0.92 Bqkg⁻¹ for ²³²Th. The activity concentrations in the cassava samples were lower when compared with the concentrations in yam samples. The high concentrations of ⁴⁰K in food samples may be due to the use of fertilizers at the farm lands. The activity concentrations in yam samples were higher than the values in cassava samples reported in the present study.

Table 3: Activity concentrations of ⁴⁰K, ²³⁸U and ²³²Th in yam and cassava samples

Samples	Yam			Cassava		
	⁴⁰ K (Bqkg ⁻¹)	²³⁸ U (Bqkg ⁻¹)	²³² Th (Bqkg ⁻¹)	⁴⁰ K (Bqkg ⁻¹)	²³⁸ U (Bqkg ⁻¹)	²³² Th (Bqkg ⁻¹)

S1	88.70±46.83	4.59±9.95	3.60±2.34	55.42±19.46	2.50±2.10	2.54±0.96
S2	93.10±41.76	4.82±8.39	3.59±3.00	59.66±17.92	3.11±1.99	2.80±0.83
S3	97.50±40.23	5.04±12.28	3.57±2.89	63.90±12.66	3.72±1.29	3.05±0.73
S4	86.42±39.77	5.13±11.62	3.26±3.01	66.66±19.73	3.84±2.01	2.56±0.98
S5	75.34±43.83	5.22±10.71	2.94±2.99	69.42±30.53	3.95±1.4 3	2.06±0.94
S6	77.13±40.92	4.97±9.98	2.91±3.22	63.40±12.71	3.70±2.11	2.17±0.98
S7	78.92±39.91	4.72±10.22	2.87±3.77	57.38±13.84	3.45±2.58	2.28±0.98
S8	82.50±41.94	4.21±12.14	2.80±3.11	45.34±15.45	2.95±2.34	2.49±0.89
S9	73.85±41.38	4.07±9.23	2.66±2.59	46.77±18.94	3.41±1.88	2.37±0.83
S10	65.20±43.39	3.92±11.71	2.51±3.89	48.20±22.84	3.86±1.98	2.24±0.88
Range	65.20-97.50	3.92-5.22	2.51-3.60	45.34-69.42	2.50-3.95	2.06-3.06
Mean	81.87	4.67	3.07	57.62	3.45	2.45

279

280 3.6 Transfer factors (TFs)

281 Generally, transfer of radionuclides depends on soil type, pH, solid/liquid distribution
 282 coefficient, exchangeable K⁺ and organic matter contents. The transfer factors (TFs) for ⁴⁰K,
 283 ²³⁸U and ²³²Th were determined and the results were presented in Table 4 below.

284 The mean value of the transfer factors for water leaf samples were 0.37±0.32, 0.82±0.10 and
 285 0.82±0.18 for ⁴⁰K, ²³⁸U and ²³²Th respectively while the mean values of the transfer factors
 286 for bitter leaf samples were 0.32±0.12, 0.74±0.09 and 0.68±0.94 for ⁴⁰K, ²³⁸U and ²³²Th
 287 respectively. The TFs values for ⁴⁰K in the samples were low compared to that of ²³⁸U and
 288 ²³²Th. The transfer values in water leaf samples were higher than that of bitter leaf samples.

289 The mean value of the transfer factors for yam samples were 0.26±0.45, 0.40±0.15 and
 290 0.32±0.88 for ⁴⁰K, ²³⁸U and ²³²Th respectively while the mean value of the transfer factors for
 291 cassava samples were 0.18±0.22, 0.29±0.13 and 0.25±0.81 for ⁴⁰K, ²³⁸U and ²³²Th
 292 respectively. The TFs values for ⁴⁰K in the samples were low compared to that of ²³⁸U and
 293 ²³²Th. The transfer values in yam samples were higher than that of cassava samples.

294 Table 4: Soil-to-yam, soil-to-cassava, soil-to-waterleaf and soil-to-bitter leaf transfer factors
 295 (TFs)

Samples	TF (yam)			TF (cassava)			TF (waterleaf)			TF (bitter leaf)		
	⁴⁰ K	²³⁸ U	²³² Th	⁴⁰ K	²³⁸ U	²³² Th	⁴⁰ K	²³⁸ U	²³² Th	⁴⁰ K	²³⁸ U	²³² Th
S1	0.28	0.45	0.39	0.17	0.25	0.28	0.37	0.98	0.82	0.34	0.93	0.73
S2	0.32	0.44	0.36	0.21	0.29	0.28	0.41	0.85	0.82	0.37	0.82	0.68
S3	0.31	0.43	0.33	0.20	0.32	0.28	0.38	0.73	0.82	0.34	0.72	0.64
S4	0.28	0.43	0.34	0.22	0.32	0.26	0.41	0.8	0.86	0.33	0.71	0.7
S5	0.25	0.43	0.35	0.23	0.33	0.24	0.45	0.86	0.91	0.33	0.71	0.77
S6	0.24	0.41	0.33	0.20	0.31	0.25	0.4	0.84	0.85	0.31	0.71	0.72
S7	0.23	0.39	0.31	0.17	0.28	0.25	0.35	0.82	0.79	0.29	0.72	0.68
S8	0.21	0.34	0.28	0.11	0.24	0.25	0.26	0.78	0.7	0.26	0.73	0.6
S9	0.21	0.33	0.26	0.13	0.28	0.23	0.31	0.79	0.77	0.3	0.69	0.64
S10	0.22	0.32	0.25	0.16	0.32	0.22	0.37	0.8	0.84	0.35	0.64	0.67

Mean	0.26	0.40	0.32	0.18	0.29	0.25	0.37	0.82	0.82	0.32	0.74	0.68
------	------	------	------	------	------	------	------	------	------	------	------	------

3.7 Absorbed dose rate and annual outdoor effective dose

The estimated absorbed dose rate in $nGyh^{-1}$ due to the radionuclide concentration in the soil was calculated for the present study. This was done to estimate the health risk associated with exposure to radioactivity in the soil from the study area. The result is shown in Tables below, the values ranged from 23.30 ± 0.67 to 28.58 ± 2.9 $nGyh^{-1}$ and mean value of 25.08 ± 0.57 $nGyh^{-1}$ for Okitipupa. The absorbed dose rate value reported in this present study is higher than the value of 12.35 ± 0.65 $nGyh^{-1}$ reported for Owo but lower than the value of 179.59 ± 4.16 $nGyh^{-1}$ reported for Oba, Ondo State ⁽¹⁴⁾.

The annual outdoor effective dose reported in this study ranged from 42891.10 to 52605.50 μSvy^{-1} with a mean value of 46166.96 μSvy^{-1} . The value reported is lower to the value of 124273.83 μSvy^{-1} reported for Oba ⁽¹⁴⁾. The value reported for annual effective dose represents 65.95% of the world average value of 70,000 μSvy^{-1} ⁽³⁾ and 47.11% of Nigeria value of 98000 μSvy^{-1} ⁽¹⁷⁾.

Table 5: Absorbed dose rate and annual outdoor effective dose of the study area

Samples	Absorbed Dose Rate	Annual Outdoor Effective Dose
	D ($nGyh^{-1}$)	E (μSvy^{-1})
S1	23.94 ± 1.31	44,070.47
S2	23.54 ± 0.32	43,332.40
S3	25.38 ± 1.89	46,719.80
S4	24.34 ± 2.90	44,805.45
S5	23.30 ± 0.75	42,891.10
S6	24.62 ± 2.25	45,319.70
S7	25.94 ± 0.92	47,748.30
S8	28.58 ± 0.71	52,605.50
S9	26.58 ± 0.63	48,927.45
S10	24.58 ± 0.33	45,249.40
Mean	25.08 ± 0.57	46,166.96




Fig 1: Soil-to-samples Transfer Factors (TFs)

Table 6: Mean Distribution of Natural Radionuclides in Soil

S/N	Location	^{40}K (Bqkg ⁻¹)	^{238}U (Bqkg ⁻¹)	^{232}Th (Bqkg ⁻¹)
1	Okitipupa, Ondo State, Nigeria (present study)	323.79	11.76	9.66
2	Akoko, Ondo State, Nigeria (Ojo and Gbadegesin, 2013)	445.02	39.24	52.86
3	World Average (UNSCEAR, 2000)	410.0	35.0	28.0

Table 7: Annual Outdoor Effective Dose (E)

S/N	Location	E (μSvy ⁻¹)
1	Okitipupa, Ondo State, Nigeria (present study)	46,166.96
2	Nigeria (Jibiri and Farai, 2005)	98,000
3	World Average (UNSCEAR, 2000)	70,000

4.0 Conclusion

The activity concentration of ^{40}K , ^{238}U and ^{232}Th in soil, waterleaf, bitter leaf, yam and cassava samples collected at Okitipupa, Ondo State Southwestern Nigeria were measured using a well calibrated NaI(Tl) and well shielded detector coupled to a computer resident quantum multichannel analyzer. The radioactivity distribution of ^{40}K , ^{238}U and ^{232}Th in all

samples were reported, the transfer factors, absorbed dose rate and annual outdoor effective doses were reported and compared with values from other literatures and the world average value.

The results showed that the concentration in the water leaf samples were higher than that in the bitter leaf samples. The results also showed that the concentration in the yam samples were higher than those in the cassava samples. The values reported were higher than values reported for some areas in various articles and lower for values in other areas.

5.0 Recommendation

It is recommended that activity concentrations should be measured in other food crops to evaluate the likely effects of consuming such crops in the area of study.

6.0 References

- [1] Abbady, E.G.A., Uosif, M.A.M. and El-Taher, A. (2005). Natural radioactivity and dose assessment for phosphate rocks from Wadi El-Mashash and El-Mahamid mines, Egypt. J. Environ. Radiat., 84, 65-78.
- [2] Abu-khadral, S.A., Abdel-sabour, M.F., Abdel-Fattah, A.T. and Eissa, H.S. (2008). Transfer factor of natural radionuclides from Egyptian soils to roots and leaves of wheat plant, 15-19, Cairo.
- [3] UNSCEAR. (2000). Sources and Effects of Ionizing Radiation. United Nations Scientific Committee on the Effects of Atomic Radiation, *Report to the General Assembly with Scientific Annexes*. United Nations: New York.
- [4] IAEA. (1989). Measurement of radionuclides in food and the environment. *Tech. Rep. Series* (No. 295), Vienna.
- [5] UNSCEAR (2008). United Nation Scientific Committee on Sources and Effects of Ionizing Radiation, Report to the Gen. Assembly, United Nations (Vol. 1, 223). New York.
- [6] NCRPM (1975). Natural Background Radiation in the United States by National Council on Radiation Protection and Measurement. NCRP Report No. 45, Washington, D.C.
- [7] Farai, I. P. and Jibiri, N. N. (2000), Baseline Studies of Terrestrial Outdoor Gamma Dose Rate Levels in Nigeria. *Radiation Protection Dosimetry*, 88(3) :247- 254.
- [8] Jibiri, N.N., Farai, I.P. and Alausa, S.K. (2006). Activity concentration of ^{226}Ra , ^{228}Th and ^{40}K in different food crops from a high background radiation area in Bitsichi, Jos Plateau, Nigeria. *Journal of Environmental Radioactivity*, 90(1), pp. 29 – 36.
- [9] FRN (2006). National Population Census Report, Federal Republic of Nigeria. web.archive.org. Retrieved 15 October 2017.
- [10] Tchokossa, P, Olomo, J.B, Balogun, F. A and Adesanmi,C.A (2013). Assessment of Radioactivity Contents of Food in the Oil and Gas Producing Areas in Delta State, Nigeria. *International Journal of Science and Technology*, 3(4):245-250.
- [11] Alharbi, A. and El - Taher, A. (2013). A study on transfer factors of radionuclides from soil to plant. *Life Science Journal* (2). <http://www.lifesciencesite.com>. Retrieved 12 September 2017.

- 366 [12] Beck, H. L., De Campo, J. A., and Gogolak, C. V. (1972). "In situ Ge(Li) and NaI (Tl)
367 Gamma Ray Spectrometry", Report HASL-258 U. S. Atomic Energy Commission, New
368 York.
- 369 [13] Chen, S.B., Zhu, Y.G. and Hu, Q.H. (2005). Soil to plant transfer of ^{238}U , ^{226}Ra and
370 ^{232}Th on a uranium mining impacted soil from southeastern China. J. Environ Rad. 78, 101-
371 111.
- 372 [14] Ojo, T.J. and Gbadegesin, K.A.J. (2013). Radioactivity in the Soils of Akoko Area of
373 Ondo State Southwestern Nigeria. Advances in Physics Theories and Applications, vol. 13.
- 374 [15] Murtadha, S.A, Mohamad, S.J. and Sabar, B. (2013). Assessment of Radionuclide
375 Transfer from Soil to Vegetables in Farms from Cameron Highlands and Penang, (Malaysia)
376 using Neutron Activation Analysis. Applied Physics Research, vol. 5, No. 5. Published by
377 Canadian Center of Science and Education.
- 378 [16] Jwanbot, D.I., Izam, M.M., Nyam, G.G. and John, H.N. (2013). Radionuclides Analysis
379 of some Soils and Food Crops in Barkin Ladi LGA, Plateau State-Nigeria. Journal of
380 Environmental and Earth Sciences, vol. 3. No.3.
- 381 [17] Jibiri, N.N. and Farai, I.P. (2005). Application of in-situ gamma-ray spectrometry in the
382 determination of activity concentrations of ^{40}K , ^{238}U and ^{232}Th and mean annual effective
383 dose rate levels in Southeastern cities in Nigeria. Radioprotection, vol. 40, No. 4, 489 – 501.