

Original Research Article

Entropic Uncertainty Relations, Entanglement and the Generalized Uncertainty Principle

Abstract

We examine quantum gravity effects by applying the generalized uncertainty principle (GUP) to entropic uncertainty relation conditions on quantum entanglement. In particular, we study the GUP corrections to the Shannon entropic uncertainty condition for entanglement. We combine previous work on the Shannon entropy entanglement criterion for bipartite systems and the GUP corrections to the Shannon entropy for a single system to calculate the GUP correction for an entangled bipartite system. As in an earlier paper of the second author, which dealt with variance relations, it is shown that there is an increase in the upper bound for the entanglement condition upon the application of the generalized uncertainty principle. Necessary fundamental concepts of the generalized uncertainty principle, entanglement and the entropic uncertainty relations are also discussed. This paper puts together the concepts of entanglement, entropic uncertainty relations and the generalized uncertainty principle all of which have been separately discussed in pedagogical papers by Schroeder, Majernik et al., Blado et al. and Sprenger.

Key words: generalized uncertainty principle, entanglement, entropic uncertainty conditions, minimal length

23 1. Introduction

24

25 The idea of a minimal length scale which limits the distance at which one can probe
 26 nature has been motivated by thought experiments involving gravitational effects in quantum
 27 mechanics and various attempts to formulate a quantum theory of gravity (such as string
 28 theory, loop quantum gravity, non-commutative geometry, etc.). In particular, string theory
 29 and certain thought experiments have been shown to give rise to a generalized uncertainty
 30 principle (GUP) from which models are developed which result in a minimal length scale [1].
 31 In this paper, we will use the most widely used model which is through the modified
 32 commutation relations.

33 The Heisenberg Uncertainty Principle (HUP) of ordinary quantum mechanics is given
 34 by (variance form)

$$35 \Delta x \Delta p \geq \frac{1}{2} (\hbar = 1).$$

$$36 (1)$$

37 This is derivable from the commutator of x and p , $[x, p] = i$, using $(\Delta A)^2 (\Delta B)^2 \geq$
 38 $\left(\frac{1}{2i} \langle [\hat{A}, \hat{B}] \rangle\right)^2$ from [2], with $A = x$ and $B = p$. String scattering has been shown to give rise
 39 to a modification of the HUP to a GUP [1, 3-5]. In this paper, we will use

$$40 \Delta x \Delta k \geq \frac{1}{2} (1 + \beta (\Delta k)^2)$$

$$41 (2)$$

42 as the form of the GUP which results from the modified commutation relation $[x, k] =$
 43 $i(1 + \beta k^2)$. k here is the momentum operator representation in terms of a GUP and β is the
 44 small positive GUP correction parameter. It is related to the HUP momentum operator by [6]

$$45 k = \frac{\tan(\sqrt{\beta} p)}{\sqrt{\beta}}.$$

$$46 (3)$$

47 Eq. (2) exhibits a non-zero minimal length [7]. There is a rich literature which discuss
 48 the GUP to which we refer the reader [1, 7-10]. Our short introduction above is meant to
 49 establish notation and discuss the most relevant equation for the discussions below.

50 Similar to [11], we study in the present paper, entanglement in continuous variable
 51 systems but instead of the variance form in Eq. (1), we will make use of entanglement
 52 conditions using the Shannon entropic uncertainty relations. The paper is organized as
 53 follows. In section 2 we give an elementary discussion of entanglement in terms of a finite

dimensional bipartite system of qubits and briefly state how entanglement and the HUP can be related for a continuous variable system. Section 3 involves an account of the entropic uncertainty relations (Shannon entropy in particular). We work out in detail the Shannon entropic quantum entanglement criterion for a bipartite system involving the Einstein-Podolsky-Rosen-type (EPR-type) operators in section 4 to make the derivation of the GUP correction apparent in section 5. Some conclusions are given in section 6.

2. Entanglement and the HUP

Particles are said to be entangled if their quantum states are somehow linked such that knowing the state of one particle determines the states of the others. Let us give an example using a qubit bipartite system to make this more explicit¹.

Suppose we have a system of two electrons (1 and 2). Let the state of electron 1 be

$$|\psi_1\rangle = \alpha_1|0\rangle + \beta_1|1\rangle \quad (4)$$

and electron 2 be

$$|\psi_2\rangle = \alpha_2|0\rangle + \beta_2|1\rangle. \quad (5)$$

The composite (pure) state of the system can be written as a tensor product (indicated by the operator $\otimes$), $|\psi\rangle = |\psi_1\rangle \otimes |\psi_2\rangle = (\alpha_1|0\rangle + \beta_1|1\rangle) \otimes (\alpha_2|0\rangle + \beta_2|1\rangle) = \alpha_1\alpha_2|0\rangle \otimes |0\rangle + \alpha_1\beta_2|0\rangle \otimes |1\rangle + \alpha_2\beta_1|1\rangle \otimes |0\rangle + \beta_1\beta_2|1\rangle \otimes |1\rangle$. A convenient notation is to let $|A\rangle \otimes |B\rangle = |AB\rangle$. So, we can write the preceding equation as

$$|\psi\rangle = \alpha_1\alpha_2|00\rangle + \alpha_1\beta_2|01\rangle + \alpha_2\beta_1|10\rangle + \beta_1\beta_2|11\rangle. \quad (6)$$

Note that the expansion coefficients are such that $|\alpha_1\alpha_2|^2 + |\alpha_1\beta_2|^2 + |\alpha_2\beta_1|^2 + |\beta_1\beta_2|^2 = 1$.

Next, consider a state

$$|\psi\rangle = \frac{1}{2\sqrt{2}}|00\rangle - \frac{\sqrt{3}}{2\sqrt{2}}|01\rangle + \frac{1}{2\sqrt{2}}|10\rangle - \frac{\sqrt{3}}{2\sqrt{2}}|11\rangle \quad (7)$$

which is known to be the composite system made up of 2 states $|\psi_1\rangle$ and $|\psi_2\rangle$ of Eq. (4) and Eq. (5). We can factorize this into 2 states,

¹ We will follow the discussion in [12].

$$\begin{aligned} |\psi\rangle &= \frac{1}{\sqrt{2}}|0\rangle\left(\frac{1}{2}|0\rangle - \frac{\sqrt{3}}{2}|1\rangle\right) + \frac{1}{\sqrt{2}}|1\rangle\left(\frac{1}{2}|0\rangle - \frac{\sqrt{3}}{2}|1\rangle\right) = \left(\frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle\right) \otimes \left(\frac{1}{2}|0\rangle - \frac{\sqrt{3}}{2}|1\rangle\right) \\ &= |\psi_1\rangle \otimes |\psi_2\rangle \end{aligned} \quad (8)$$

with $|\psi_1\rangle = \left(\frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle\right)$ and $|\psi_2\rangle = \left(\frac{1}{2}|0\rangle - \frac{\sqrt{3}}{2}|1\rangle\right)$. Factorizable states are separable states [13]. Hence, we say that $|\psi\rangle$ in Eq. (7) is a separable state.

However, given a composite state, it is not always possible to factorize it into individual states! For example, consider the state,

$$|\psi'\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle. \quad (9)$$

If this is to factorize, then the right-hand side of Eq. (9) must be equal to the tensor product of two states. Let these two general states be as in Eq. (4) and Eq. (5). From Eq. (9) and Eq. (6), $\frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle = \alpha_1\alpha_2|00\rangle + \alpha_1\beta_2|01\rangle + \alpha_2\beta_1|10\rangle + \beta_1\beta_2|11\rangle$. Comparing coefficients, we get $\alpha_1\alpha_2 = \frac{1}{\sqrt{2}}$ so NEITHER α_1 NOR α_2 CAN BE ZERO. We also get $\beta_1\beta_2 = \frac{1}{\sqrt{2}}$ so NEITHER β_1 NOR β_2 CAN BE ZERO. However, we also get $\alpha_1\beta_2 = 0$ which means that either $\alpha_1 = 0$ or $\beta_2 = 0$! We have a contradiction! Hence, there is no solution and we cannot write $|\psi\rangle$ of Eq. (9) as a tensor product of the two states in Eq. (4) and Eq. (5). $|\psi'\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$ is an example of an entangled state. We can say that for the entangled state $|\psi'\rangle \neq |\psi_1\rangle \otimes |\psi_2\rangle$.

$$(10)$$

In the discussion above we can consider the state $|0\rangle$ to describe a particle with spin UP and $|1\rangle$ a particle with spin DOWN. So, for composite systems, the state $|10\rangle$ will describe the composite state in which particle 1 is spin down and particle 2 is spin up, etc. If we measure the spin of particle 1 in Eq. (9) to be up then we are sure that the second particle will have spin up also because of the composite state $|00\rangle$. This, however is not true for Eq. (7). Due to the presence of the two terms $|00\rangle$ and $|01\rangle$, if particle 1 is spin up $|0\rangle$, there is a probability that the second particle is also spin down $|1\rangle$! Hence, Eq. (9) is called an entangled state because if we know the state of one particle, we surely know the state of the other unlike in Eq. (7).

The above example which makes use of a finite dimensional system, is a standard starting point in discussing the concept of entanglement. To explore the connection of the HUP and entanglement, we need to look at continuous variable systems which will involve

the position and momentum operators. An excellent introductory discussion on entanglement beyond spin systems can be found in [14]. A more advanced treatment is given in [15].

Similar to Eq. (10) an entangled composite state represented by $\Psi(x_1, x_2)$ in coordinate space or $\mathcal{P}(p_1, p_2)$ in momentum space is such that $\Psi(x_1, x_2) \neq \psi_1(x_1)\psi_2(x_2)$ or $\mathcal{P}(p_1, p_2) \neq \varphi_1(p_1)\varphi_2(p_2)$ where $\psi_i(x_i)$ ($\varphi_i(p_i)$) could represent the coordinate (momentum) space wave function of particle i . With the states dependent on the position and momentum (which are continuous variables), and with the uncertainty principle expressed in terms of the position and momentum operators in Eq. (1), one could draw connections between the entanglement and the HUP [16-19] and subsequently with the GUP [11].

3. Entropic Uncertainty Relations

For two observables that do not commute, such as position and momentum, Eq. (1) addresses a fundamental limit to the extent with which one can know about one of the particle's properties if the other has already been measured. For years, the favored approach towards uncertainty in quantum mechanics was the use of standard deviations in Eq. (1) primarily because in probability and statistics, the dispersion is the most well-known gauge of uncertainty in measurements [20].

However, as pointed out by [20], it was not the ideal measure, indicated by two major flaws that appeared in the Robertson relation [21], the expression from which Eq. (1) can be derived when some non-commuting observables A and B have been specified to be position and momentum,

$$\Delta A \Delta B \geq \frac{1}{2} |\langle \Psi | [\hat{A}, \hat{B}] | \Psi \rangle|.$$

(11)

The first weakness is that in a finite, N-dimensional Hilbert space, the lower bound in the right-hand side becomes dependent upon the state of the particle, which becomes especially problematic when the state is an eigenstate that gives an eigenvalue of the observable. Eq. (11) yields zero which basically eliminates the restriction on the variances of A and B . The second weakness is that for conjugate variables with continuous probability distributions, the dispersion may not be the useful measure of uncertainty especially if the distributions have several peaks. This is especially true for applications in information theory where standard deviation does not play an important role [22].

To overcome these issues, alternative uncertainty relations were sought out by physicists. Recently, an approach utilizing entropic uncertainty relations has been gaining traction due to the relations' clear validity as mathematical inequalities [23]. Consequently, it has been postulated that such relations may bear physical significance in measurements and so the search of a useful relation, if any exist, for analysis of quantum mechanical systems has been the topic of debate [24-25]. The desirable features of the entropic uncertainty relations over the HUP have been discussed in [22].

In favor of the view that entropic uncertainty relations can be useful in experiments, many physicists attempt to explain what entropy can mean in a quantum context. In one particular definition, entropy can be seen as the amount of information one lacks when one witnesses an event that could have many outcomes [26]. This information can be used to characterize its source based not on what the information is about, but on how much of it is transmitted compared to how much of it can be transmitted.

One way to look at it is similar to Shannon's own approach about compressing and decompressing messages in which some information that cannot be compressed is not sent, so it is lost. Recall the difference between one's thoughts and one's speech. A person can think about something with all their knowledge about it at their disposal, but expressing what they think into words may not convey everything that they know to a listener. The lack of a complete view about what the speaker knows about something, would be reflected in information entropy just like the measurement of some particle's wavefunction.

To understand the definition of the particular entropic function we will be using, let us first go back to the Boltzmann entropy $S = k_B \ln W$, of classical thermodynamics and statistical mechanics [27-28] where k_B is the Boltzmann's constant and W is the number of microscopic configurations that a system can be in, assuming equiprobable configurations. Hence $W = 1/P$ where P is the probability of each configuration. Thus $S/k_B = -\ln(P)$. However, if the different configurations are not equiprobable, then we need to add the weight-factor of them occurring as we sum over the different probabilities P_i ,

$$S/k_B = -\sum_i P_i \ln(P_i).$$

(12)

From Eq. (12), Shannon [26] defined the (dimensionless) average information entropy content of a random variable X (called the Shannon entropy) as

$$H \equiv \sum_x P_X(x) \ln(1/P_X(x)) = -\sum_x P_X(x) \ln(P_X(x)),$$

(13)

where $P_X(x)$ is the probability distribution of the random variable X to have the measured value x . Note that if the probability of measuring X as x is 1 then there is no information entropy or no uncertainty. The lower the probability of measuring x , the higher the information entropy or uncertainty. Hence information entropy can be considered a measure of uncertainty.

Extending Eq. (13) to continuous variables, we get the differential Shannon entropy (or we will refer to this as simply “Shannon entropy” in the succeeding sections) as

$$H = - \int_{x_1}^{x_2} P(x) \ln(P(x)) dx$$

(14)

where $P(x)$ is the probability distribution of the continuous variable x .

It is apparent that we can define the Shannon entropy involving position and momentum probability distributions from Eq. (14) as

$$H[w] = - \int_{-\infty}^{\infty} dx w(x) \ln w(x); H[v] = - \int_{-\infty}^{\infty} dp v(p) \ln v(p)$$

(15)

where $w(x) \equiv |\psi(x)|^2$ and $v(p) \equiv |\varphi(p)|^2$. Bialynicki-Birula and Mycielski [29] derived an entropic uncertainty relation (“BBM inequality”) involving the Shannon entropy similar to the HUP

$$H[w] + H[v] \geq \ln(\pi e).$$

(16)

The above expression owes its validity to a fact that Heisenberg himself acknowledged—to gain information about the position of a particle, one must lose information about the momentum of that same particle and vice versa due to their nature as conjugate variables [30]. Essentially, as the entropy of one property decreases, the entropy of the other must increase to maintain the total uncertainty of the particle above the limit shown in the right-hand side of Eq. (16). In addition, although the HUP describes uncertainty due to a lack of the simultaneous existence of precise values for both position and momentum, it ignores the uncertainty induced by the measurement of an actual instrument, which Eq. (16) can account for [23]. Coles, et al. [22] discussed how Eq. (16) overcomes the two weaknesses of the HUP discussed above.

Since the BBM inequality is a reliable representation of the uncertainty inherent in knowing the properties of one particle, it will also be useful when considering two particles which apparently involves more information. This forms the basis of an entropic uncertainty relation that can be used to detect a specific characteristic of the two-particle system, namely

entanglement, due to the fact that the measurement of one correlated particle gives rise to information on the other particle as well, resulting to information about both particles instead of only that which was measured. This mechanism decreases the total uncertainty of the system beyond its expected limit. Identification of entangled particles has always proven inconvenient with factorization since the process can sometimes be tedious and it is difficult to be certain that some composite states are definitively inseparable. A lack of factorizability can indicate that particles are entangled, but it can also mean that the composite state is simply very complex, so an entropic criterion that can guarantee entanglement would be very useful indeed.

4. Shannon Entropy Quantum Entanglement Criterion

The following discussion will follow closely Walborn, et al. [31]. Considering a bipartite system, we define operators similar to the annihilation operators defined in [32], which are linear combinations of the position and momentum operators x_j and p_j with $j=1, 2$, pertaining to subsystems 1 and 2, and the usual commutation relation, $[x_j, p_k] = i\delta_{jk}$

$$r_j = \cos(\theta_j)x_j + \sin(\theta_j)p_j, \quad s_j = \cos(\theta_j)p_j - \sin(\theta_j)x_j.$$

(17)

From Eq. (17), we define further the operators, $r_{\pm} = r_1 \pm r_2$ and $s_{\pm} = s_1 \pm s_2$. As in [33], we study EPR-type operators by letting $\theta_1 = \theta_2 = 0$ in Eq. (17) which give, similar to the general $r_{\pm}$ and $s_{\pm}$ operators,

$$x_{\pm} = x_1 \pm x_2; \quad p_{\pm} = p_1 \pm p_2.$$

(18)

Consider a separable pure state $|\Psi\rangle = |\psi_1\rangle \otimes |\psi_2\rangle$ with the corresponding wavefunction in coordinate space, $\Psi(x_1, x_2) = \psi_1(x_1)\psi_2(x_2)$ and in momentum space $\mathcal{P}(p_1, p_2) = \varphi_1(p_1)\varphi_2(p_2)$. A change in variables in Eq. (18), yields

$$\frac{1}{\sqrt{2}}\psi_1\left(\frac{x_++x_-}{2}\right)\psi_2\left(\frac{x_+-x_-}{2}\right) = \Psi(x_+, x_-); \quad \frac{1}{\sqrt{2}}\varphi_1\left(\frac{p_++p_-}{2}\right)\varphi_2\left(\frac{p_+-p_-}{2}\right) = \mathcal{P}(p_+, p_-).$$

(19)

Let us compute the Shannon entropy associated with the measurement of $x_{\pm}$ and $p_{\pm}$. From the definition of the Shannon entropy in Eq. (15),

$$H[w_{\pm}] = -\int_{-\infty}^{\infty} dx_{\pm} w_{\pm}(x_{\pm}) \ln(w_{\pm}(x_{\pm})); \quad H[v_{\pm}] = -\int_{-\infty}^{\infty} dp_{\pm} v_{\pm}(p_{\pm}) \ln(v_{\pm}(p_{\pm})), \text{ where}$$

245 $w_{\pm}(x_{\pm}) = w_{\pm}(x_1, x_2)$ and $v_{\pm}(p_{\pm}) = v_{\pm}(p_1, p_2)$ are the probability distributions of $x_{\pm}$ and
 246 $p_{\pm}$ respectively. From Eq. (19), these are given by

$$247 \quad w_{\pm}(x_{\pm}) = \int_{-\infty}^{\infty} dx_{\mp} |\Psi(x_+, x_-)|^2 = \frac{1}{2} \int_{-\infty}^{\infty} dx_{\mp} |\psi_1|^2 |\psi_2|^2 ;$$

$$248 \quad v_{\pm}(p_{\pm}) = \int_{-\infty}^{\infty} dp_{\mp} |\varphi(p_+, p_-)|^2 = \frac{1}{2} \int_{-\infty}^{\infty} dp_{\mp} |\varphi_1|^2 |\varphi_2|^2.$$

$$249 \quad (20)$$

250 Let us define $w_i(x_i) \equiv |\psi_i(x_i)|^2$ and $v_i(p_i) \equiv |\varphi_i(p_i)|^2$. Hence Eq. (20) becomes

$$251 \quad w_{\pm}(x_{\pm}) = \frac{1}{2} \int_{-\infty}^{\infty} dx_{\mp} w_1\left(\frac{x_+ + x_-}{2}\right) w_2\left(\frac{x_+ - x_-}{2}\right) \text{ and } v_{\pm}(p_{\pm}) = \frac{1}{2} \int_{-\infty}^{\infty} dp_{\mp} v_1\left(\frac{p_+ + p_-}{2}\right) v_2\left(\frac{p_+ - p_-}{2}\right).$$

$$252 \quad (21)$$

253 For convenience, as will be apparent later, we make a further change in variables in

254 terms of $\frac{x_+ + x_-}{2} = x_1 = x$ and so $\frac{x_+ - x_-}{2} = \mp x \pm x_{\pm}$ and similarly $\frac{p_+ + p_-}{2} = p_1 = p$ and so

$$255 \quad \frac{p_+ - p_-}{2} = \mp p \pm p_{\pm}. \quad \text{Eq. (21) becomes} \quad w_{\pm} = \int_{-\infty}^{\infty} dx w_1(x) w_2(\mp x \pm x_{\pm});$$

256 $v_{\pm} = \int_{-\infty}^{\infty} dp v_1(p) v_2(\mp p \pm p_{\pm})$ which in terms of the convolution operation “*” can be
 257 written simply as

$$258 \quad w_{\pm} = w_1 * w_2^{\pm}, \quad v_{\pm} = v_1 * v_2^{\pm} \text{ with } w_2^{\pm} = w_2(\pm x) \text{ and } v_2^{\pm} = v_2(\pm p).$$

$$259 \quad (22)$$

260 From the inequality [26, 34],

$$261 \quad e^{2H[A*B]} \geq e^{2H[A]} + e^{2H[B]}$$

$$262 \quad (23)$$

263 and with [26]

$$264 \quad H[w_2^+] = H[w_2^-] \text{ and } H[v_2^+] = H[v_2^-]$$

$$265 \quad (24)$$

266 we get from Eq. (22)

$$267 \quad H[w_{\pm}] = H[w_1 * w_2^{\pm}] \geq \frac{1}{2} \ln\{e^{2H[w_1]} + e^{2H[w_2]}\}; \quad H[v_{\mp}] = H[v_1 * v_2^{\mp}] \geq \frac{1}{2} \ln\{e^{2H[v_1]} +$$

$$268 \quad e^{2H[v_2]}\}. \quad (25)$$

269 The preceding equation leads to state-dependent inequalities satisfied by separable
 270 pure states.

$$271 \quad H[w_{\pm}] + H[v_{\mp}] \geq \frac{1}{2} \ln\{(e^{2H[w_1]} + e^{2H[w_2]})(e^{2H[v_1]} + e^{2H[v_2]})\} \equiv h[w_i, v_j]$$

$$272 \quad (26)$$

273 Hence if

$$H[w_{\pm}] + H[v_{\mp}] < \frac{1}{2} \ln\{(e^{2H[w_1]} + e^{2H[w_2]})(e^{2H[v_1]} + e^{2H[v_2]})\} \equiv h[w_i, v_j]$$

(27)

then we have entangled states.

From the BBM inequality [29], we can write

$$H[w_j] + H[v_j] \geq \ln(\pi e) \text{ with } j = 1, 2.$$

(28)

This yields from Eq. (26),

$$H[w_{\pm}] + H[v_{\mp}] \geq \frac{1}{2} \ln\{2(\pi e)^2 + e^{2H[w_1]+2H[v_2]} + e^{2H[w_2]+2H[v_1]}\}.$$

(29)

Using Eq. (28) once again, we can write Eq. (29) in terms of only w_j (or v_j) as $H[w_{\pm}] +$

$$H[v_{\mp}] \geq \frac{1}{2} \ln\{2(\pi e)^2 + (\pi e)^2 [e^{2(H[v_2]-H[v_1])} + e^{-2(H[v_2]-H[v_1])}]\}$$
 or simply

$$H[w_{\pm}] + H[v_{\mp}] \geq \frac{1}{2} \ln\{2(\pi e)^2 (1 + \cosh \Delta H)\} \geq \ln(2\pi e)$$

(30)

where $\Delta H \equiv H[v_2] - H[v_1]$. One can compute that the minimum value of the argument of the natural logarithm function occurs when $\Delta H = 0$ which then yields the minimum value $4(\pi e)^2 = (2\pi e)^2$. Substituting this value, gives the last inequality in the preceding equation. Although Eq. (30) is a weaker set of state-independent inequalities, states with

$$H[w_{\pm}] + H[v_{\mp}] < \ln(2\pi e)$$

(31)

are still entangled states. However entangled states can still exist for $\ln(2\pi e) \leq H[w_{\pm}] + H[v_{\mp}] < h[w_i, v_j]$ owing to the stronger inequalities in Eq. (26) and Eq. (27).

Let us next consider mixed states. Mixed states are more commonly encountered experimentally since a system can be in any of a number of states in varying probabilities which can occur when we prepare a similar system several times. Note that a mixed state is different from a state formed by a superposition of states (which can be a pure state). A bipartite separable mixed state is more conveniently described in terms of the density matrix ρ .

$$\rho = \sum_m \lambda_m \rho_{1m} \otimes \rho_{2m}, \text{ where } \rho_{im} = |\psi_{im}\rangle\langle\psi_{im}|, i = 1, 2, \lambda_m \geq 0, \sum_m \lambda_m = 1.$$

(32)

Note that $\psi_{1m} = (\psi_1(r_1))_m$ is the m th pure state for subsystem 1 and $\psi_{2m} = (\psi_2(r_2))_m$ is the m th pure state for subsystem 2. We can write the probability distribution for $x_{\pm}$ and $p_{\pm}$ as

$$w_{\pm} = \sum_m \lambda_m w_{m\pm}, v_{\pm} = \sum_m \lambda_m v_{m\pm} \quad (33)$$

where $w_{m\pm}$ ($v_{m\pm}$) is the probability distribution to detect $x_{\pm}$ ($p_{\pm}$) for each pure state ψ_{im} .

Since the Shannon entropy is a concave function [34], we have

$$H[w_{\pm}] = H[\sum_m \lambda_m w_{m\pm}] \geq \sum_m \lambda_m H[w_{m\pm}]; \quad H[v_{\mp}] = H[\sum_m \lambda_m v_{m\mp}] \geq \sum_m \lambda_m H[v_{m\mp}]. \quad (34)$$

Similarly, from Eq. (25), we get

$$H[w_{\pm}] \geq H[\sum_m \lambda_m w_{m\pm}] \geq \sum_m \lambda_m \frac{1}{2} \ln\{e^{2H[w_{1m}]} + e^{2H[w_{2m}]}\}; \quad H[v_{\mp}] \geq H[\sum_m \lambda_m v_{m\mp}] \geq \sum_m \lambda_m \frac{1}{2} \ln\{e^{2H[v_{1m}]} + e^{2H[v_{2m}]}\}. \quad (35)$$

Hence, adding the preceding equations,

$$H[w_{\pm}] + H[v_{\mp}] \geq \sum_m \lambda_m \frac{1}{2} \ln[(e^{2H[w_{1m}]} + e^{2H[w_{2m}]})(e^{2H[v_{1m}]} + e^{2H[v_{2m}]})] \equiv h[w_{im}, v_{jm}] \quad (36)$$

for separable states where $i, j = 1, 2$. Hence for entangled mixed states we have,

$$H[w_{\pm}] + H[v_{\mp}] < h[w_{im}, v_{jm}]. \quad (37)$$

Going back to Eq. (34), we can write $H[w_{\pm}] + H[v_{\mp}] \geq \sum_m \lambda_m (H[w_{m\pm}] + H[v_{m\mp}])$ and from Eq. (30), we get $H[w_{\pm}] + H[v_{\mp}] \geq \sum_m \lambda_m (\ln(2\pi e)) = (\ln(2\pi e)) \sum_m \lambda_m = (\ln(2\pi e)) \cdot 1$ from Eq. (32). Hence, $H[w_{\pm}] + H[v_{\mp}] \geq (\ln(2\pi e))$ even for separable mixed states and similarly we get the same weaker conditions for entanglement in Eq. (31) for mixed states

$$H[w_{\pm}] + H[v_{\mp}] < (\ln(2\pi e)). \quad (38)$$

5. GUP-correction to the Shannon Entropy Quantum Entanglement Criterion

Having set the notation and discussed the Shannon Entropy Quantum Entanglement Criterion in section 4, we use the results of [35, 6] to get the GUP correction. As discussed in section 1, the GUP-modified commutation relation is now given by $[x_i, k_j] = i\delta_{ij}(1 + \beta k_i^2)$ with i , and j as in section 4 above. From Eq. (3), p_i and k_i are related by

$$k_i = \frac{1}{\sqrt{\beta}} \tan(\sqrt{\beta} p_i).$$

$$(39)$$

We now have for the k momentum operator, analogous to $p(p_+, p_-)$,

$$\Phi(k_+, k_-) = \frac{1}{\sqrt{2}} \phi_1\left(\frac{k_+ + k_-}{2}\right) \phi_2\left(\frac{k_+ - k_-}{2}\right)$$

$$(40)$$

with

$$k_{\pm} = k_1 \pm k_2 \text{ and } \phi_1 = \phi_1(k_1), \phi_2 = \phi_2(k_2)$$

$$(41)$$

and the corresponding Shannon entropy $H[u_{\pm}] = - \int_{-\infty}^{\infty} dk_{\pm} u_{\pm}(k_{\pm}) \ln(u_{\pm}(k_{\pm}))$ where

$u_{\pm}(k_{\pm}) = u_{\pm}(k_1, k_2)$ is the probability distribution of $k_{\pm}$. Hence from Eq. (40),

$$u_{\pm}(k_{\pm}) = \int_{-\infty}^{\infty} dk_{\mp} |\Phi(k_+, k_-)|^2 = \frac{1}{2} \int_{-\infty}^{\infty} dk_{\mp} |\phi_1|^2 |\phi_2|^2.$$

$$(42)$$

Defining $u_i(k_i) \equiv |\phi_i(k_i)|^2$, Eq. (42) becomes

$$u_{\pm}(k_{\pm}) = \frac{1}{2} \int_{-\infty}^{\infty} dk_{\mp} u_1\left(\frac{k_+ + k_-}{2}\right) u_2\left(\frac{k_+ - k_-}{2}\right).$$

$$(43)$$

From Eq. (41), we define $\frac{k_+ + k_-}{2} = k_1 \equiv k$, and so we get $\frac{k_+ - k_-}{2} = \mp k \pm k_{\pm}$ then Eq.

(43) becomes $u_{\pm}(k_{\pm}) = \int_{-\infty}^{\infty} dk u_1(k) u_2(\mp k \pm k_{\pm})$ which in terms of the convolution

operation “*” becomes $u_{\pm}(k_{\pm}) = u_1 * u_2^{\pm}$ with $u_2^{\pm} \equiv u_2(\pm k)$. Using Eq. (23) and Eq. (24)

(with v replaced by u), we get similar to the second part of Eq. (25),

$$H[u_{\mp}] = H[u_1 * u_2^{\mp}] \geq \frac{1}{2} \ln\{e^{2H[u_1]} + e^{2H[u_2]}\}.$$

$$(44)$$

The probability distributions of the operators k and p are such that $u_i(k_i) dk_i =$

$v_i(p_i) dp_i$ or $u_i(k_i) = v_i(p_i) \frac{dp_i}{dk_i}$ with $i = 1, 2$. Using Eq. (39), we get

$$u_i(k_i) = \frac{v_i(p_i)}{1 + \beta k_i^2}.$$

$$(45)$$

From the definition of the Shannon entropy Eq. (15) and Eq. (45),

$$H[u_i] = - \int_{-\infty}^{\infty} u_i(k_i) \ln u_i(k_i) dk_i = - \int_{-\infty}^{\infty} \frac{v_i(p_i)}{1 + \beta k_i^2} \ln\left(\frac{v_i(p_i)}{1 + \beta k_i^2}\right) dk_i =$$

$$- \int_{-\infty}^{\infty} \frac{v_i(p_i)}{1 + \beta k_i^2} \ln v_i(p_i) dk_i + \int_{-\infty}^{\infty} \frac{v_i(p_i)}{1 + \beta k_i^2} \ln(1 + \beta k_i^2) dk_i = - \int_{-\infty}^{\infty} \frac{v_i(p_i)}{1 + \beta k_i^2} \ln v_i(p_i) dk_i +$$

364 $\int_{-\infty}^{\infty} u_i(k_i) \ln(1 + \beta k_i^2) dk_i$. Finally, using again Eq. (39) for a change in the integrating
 365 variable in the first integral yields $H[u_i] = -\int_{-p_0}^{p_0} v_i(p_i) \ln(v_i(p_i)) dp_i + \int_{-\infty}^{\infty} u_i(k_i) \ln(1 +$
 366 $\beta k_i^2) dk_i$ with $p_0 \equiv \frac{\pi}{2\sqrt{\beta}}$. Hence, we can write

$$367 \quad H[u_i] = H[v_i] + \eta_i(\beta) \text{ with } \eta_i(\beta) \equiv \int_{-\infty}^{\infty} u_i(k_i) \ln(1 + \beta k_i^2) dk_i > 0 .$$

368 (46)

369 $\eta_i(\beta)$ basically gives the GUP correction. From Eq. (46) and Eq. (44), we get

$$370 \quad H[u_{\mp}] \geq \frac{1}{2} \ln\{e^{2H[v_1]+2\eta_1(\beta)} + e^{2H[v_2]+2\eta_2(\beta)}\}.$$

371 (47)

372 Combining the preceding equation with the inequality for $w_{\pm}$ in Eq. (25), we get $H[w_{\pm}] +$

$$373 \quad H[u_{\mp}] \geq \frac{1}{2} \ln\{e^{2\eta_1(\beta)} e^{2H[v_1]} + e^{2\eta_2(\beta)} e^{2H[v_2]}\} + \frac{1}{2} \ln\{e^{2H[w_1]} + e^{2H[w_2]}\} \text{ or}$$

$$374 \quad H[w_{\pm}] + H[u_{\mp}] \geq \frac{1}{2} \ln\{(e^{2H[w_1]} + e^{2H[w_2]})(e^{2\eta_1(\beta)} e^{2H[v_1]} + e^{2\eta_2(\beta)} e^{2H[v_2]})\} \equiv$$

375 $h_{GUP}(w_i, v_j)$ (48)

376 for separable pure states. Since from Eq. (46) $\eta_i(\beta = 0) = 0$ for the non-GUP case, Eq. (48)
 377 reduces to Eq. (26) as expected. Hence, if $H[w_{\pm}] + H[u_{\mp}] < h_{GUP}(w_i, v_j)$ then we have
 378 entangled states. Clearly since $\eta_i(\beta) > 0$ in Eq. (46), we see from Eq. (26) and Eq. (48) that
 379 $h_{GUP}(w_i, v_j) > h[w_i, v_j]$. Hence with Eq. (27) and Eq. (48) we have a higher upper bound for
 380 the GUP-corrected entanglement condition. It is interesting to note that a similar increase in
 381 the upper bound for the GUP-corrected entanglement condition was found when using
 382 variance relations [11].

383 We now derive the corresponding GUP-corrected expression for Eq. (30). Since the
 384 BBM inequality has been shown to still hold in the GUP framework [6], we can combine Eq.
 385 (46) and Eq. (28) to yield

$$386 \quad H[u_i] + H[w_i] \geq \ln(\pi e) + \eta_i(\beta)$$

387 (49)

388 which yields $e^{H[u_i]+H[w_i]} \geq (\pi e) e^{\eta_i(\beta)}$ or

$$389 \quad e^{2(H[u_i]+H[w_i])} \geq (\pi e)^2 e^{2\eta_i(\beta)}.$$

390 (50)

391 From Eq. (44) and Eq. (25), we get $H[w_{\pm}] + H[u_{\mp}] \geq \frac{1}{2} \ln(e^{2H[w_1]} + e^{2H[w_2]}) +$
 392 $\frac{1}{2} \ln\{e^{2H[u_1]} + e^{2H[u_2]}\} = \frac{1}{2} \ln\{(e^{2H[w_1]} + e^{2H[w_2]})(e^{2H[u_1]} + e^{2H[u_2]})\}$ or

$$H[w_{\pm}] + H[u_{\mp}] \geq \frac{1}{2} \ln(e^{2H[w_1]+2H[u_1]} + e^{2H[w_2]+2H[u_2]} + e^{2H[w_1]+2H[u_2]} + e^{2H[w_2]+2H[u_1]}).$$

(51)

Because of Eq. (50), we can replace the first two terms of Eq. (51) to yield

$$H[w_{\pm}] + H[u_{\mp}] \geq \frac{1}{2} \ln\{(\pi e)^2 e^{2\eta_1(\beta)} + (\pi e)^2 e^{2\eta_2(\beta)} + e^{2H[w_1]+2H[u_2]} + e^{2H[w_2]+2H[u_1]}\}.$$

(52)

Let us look at the third term of the preceding equation. Using Eq. (49) to eliminate $H[w_1]$, we have $e^{2H[w_1]+2H[u_2]} \geq e^{2(\ln(\pi e)+\eta_1(\beta)-H[u_1])+2H[u_2]} = (\pi e)^2 e^{2\eta_1(\beta)} e^{2(H[u_2]-H[u_1])}$. Similarly, we have $e^{2H[w_2]+2H[u_1]} \geq (\pi e)^2 e^{2\eta_2(\beta)} e^{-2(H[u_2]-H[u_1])}$. We can then replace the last two terms in Eq. (52) to give $H[w_{\pm}] + H[u_{\mp}] \geq \frac{1}{2} \ln\{(\pi e)^2 e^{2\eta_1(\beta)} + (\pi e)^2 e^{2\eta_2(\beta)} + (\pi e)^2 e^{2\eta_1(\beta)} e^{2(H[u_2]-H[u_1])} + (\pi e)^2 e^{2\eta_2(\beta)} e^{-2(H[u_2]-H[u_1])}\}$. The right-hand side readily simplifies to

$$H[w_{\pm}] + H[u_{\mp}] \geq \ln(\pi e) + \frac{1}{2} \ln\{(e^{2\eta_1(\beta)} + e^{2\eta_2(\beta)}) + (e^{2\eta_1(\beta)} e^{2\Delta H_{GUP}} + e^{2\eta_2(\beta)} e^{-2\Delta H_{GUP}})\} \quad (53)$$

with $\Delta H_{GUP} \equiv H[u_2] - H[u_1]$. As in section 4, Eq. (30), let us attempt to rewrite the inequality independent of the Shannon entropy difference ΔH_{GUP} . We can solve for the minimum of $e^{2\eta_1(\beta)} + e^{2\eta_2(\beta)} + e^{2\eta_1(\beta)} e^{2\Delta H_{GUP}} + e^{2\eta_2(\beta)} e^{-2\Delta H_{GUP}}$. It occurs at $\Delta H_{GUP} = \frac{1}{2} \ln\left(\frac{e^{2\eta_2(\beta)}}{e^{2\eta_1(\beta)}}\right)$. This yields a minimum value of $e^{2\eta_1(\beta)} + e^{2\eta_2(\beta)} + 2e^{\eta_1(\beta)} e^{\eta_2(\beta)} = (e^{\eta_1(\beta)} + e^{\eta_2(\beta)})^2$. We can substitute this into the argument of the natural logarithm function in Eq. (53) to get $H[w_{\pm}] + H[u_{\mp}] \geq \ln(\pi e) + \ln(e^{\eta_1(\beta)} + e^{\eta_2(\beta)})$ or

$$H[w_{\pm}] + H[u_{\mp}] \geq \ln(2\pi e) + \ln\left(\frac{e^{\eta_1(\beta)} + e^{\eta_2(\beta)}}{2}\right).$$

(54)

Eq. (54) is the GUP-corrected version of Eq. (30). Unlike Eq. (30), the preceding equation is not state-independent. The GUP-corrected version of Eq. (31) for entangled states is then

$$H[w_{\pm}] + H[u_{\mp}] < \ln(2\pi e) + \ln\left(\frac{e^{\eta_1(\beta)} + e^{\eta_2(\beta)}}{2}\right).$$

(55)

As before the upper bound for entangled states is increased when compared to Eq. (31).

The GUP correction for mixed states expressed in Eq. (32) follows naturally. Similar to Eq. (33),

$$w_{\pm} = \sum_m \lambda_m w_{m\pm}, u_{\pm} = \sum_m \lambda_m u_{m\pm}$$

(56)

where $u_{m\pm}$ is the probability distribution to detect $k_{\pm}$ for each pure state ψ_{im} .

Similar to Eq. (46),

$$H[u_{im}] = H[v_{im}] + \eta_{im}(\beta) \text{ with } \eta_{im}(\beta) \equiv \int_{-\infty}^{\infty} u_{im}(k_i) \ln(1 + \beta k_i^2) dk_i > 0, \eta_{im}(\beta),$$

$$i = 1, 2 \text{ (57)}$$

giving the GUP correction. Using the concavity of the Shannon entropy function, Eq. (56)

gives

$$H[u_{\mp}] = H[\sum_m \lambda_m u_{m\mp}] \geq \sum_m \lambda_m H[u_{m\mp}].$$

$$(58)$$

Similar to Eq. (47), $H[u_{m\mp}] \geq \frac{1}{2} \ln\{e^{2H[v_{1m}]+2\eta_{1m}(\beta)} + e^{2H[v_{2m}]+2\eta_{2m}(\beta)}\}$. Hence Eq. (58)

becomes $H[u_{\mp}] \geq \sum_m \lambda_m \frac{1}{2} \ln\{e^{2H[v_{1m}]+2\eta_{1m}(\beta)} + e^{2H[v_{2m}]+2\eta_{2m}(\beta)}\}$. Adding the preceding

equation to Eq. (35),

$$H[w_{\pm}] + H[u_{\mp}] \geq$$

$$\sum_m \lambda_m \frac{1}{2} \ln\{(e^{2H[w_{1m}]} + e^{2H[w_{2m}]}) (e^{2H[v_{1m}]+2\eta_{1m}(\beta)} + e^{2H[v_{2m}]+2\eta_{2m}(\beta)})\} \equiv$$

$$\hbar_{GUP}(w_{im}, v_{jm})$$

$$(59)$$

for separable states. Hence for entangled mixed states, we have $H[w_{\pm}] + H[u_{\mp}] <$

$$\hbar_{GUP}(w_{im}, v_{jm}).$$

Hence, similar to the conclusion from Eq. (48), we get a higher upper bound for the GUP-

corrected entanglement condition as compared to Eq. (37) since $\hbar_{GUP}(w_{im}, v_{jm}) >$

$\hbar_{GUP}(w_{im}, v_{jm})$ as in the pure state case.

Returning to Eq. (34) and Eq. (58), and similar to Eq. (54),

$$H[w_{\pm}] + H[u_{\mp}] \geq \sum_m \lambda_m (H[w_{m\pm}] + H[u_{m\mp}]) \geq$$

$$\sum_m \lambda_m \left(\ln(2\pi e) + \ln \left[\frac{e^{\eta_{1m}(\beta)} + e^{\eta_{2m}(\beta)}}{2} \right] \right) = \sum_m \lambda_m (\ln(2\pi e)) + \sum_m \lambda_m \ln \left[\frac{e^{\eta_{1m}(\beta)} + e^{\eta_{2m}(\beta)}}{2} \right]$$

$$\text{and from Eq. (32), } H[w_{\pm}] + H[u_{\mp}] \geq \ln(2\pi e) + \sum_m \lambda_m \ln \left[\frac{e^{\eta_{1m}(\beta)} + e^{\eta_{2m}(\beta)}}{2} \right]. \text{ The}$$

entanglement criterion is then

$$H[w_{\pm}] + H[u_{\mp}] < \ln(2\pi e) + \sum_m \lambda_m \ln \left[\frac{e^{\eta_{1m}(\beta)} + e^{\eta_{2m}(\beta)}}{2} \right].$$

$$(60)$$

Eq. (60) is the GUP-corrected expression for Eq. (38) for mixed states (generalized for both

pure states and mixed states) which shows an increase in the upper bound of Eq. (38). It is

apparent that Eq. (60) reduces to Eq. (55) for a pure state with a weight-factor of 1.

6. Conclusions

The study of the effects of the minimal length and the generalized uncertainty principle on entanglement involving continuous variable systems has been confined to the variance form of the HUP [11]. Since the more modern approach of expressing the uncertainty principle is through the use of entropic measures [36], as discussed in section 3 above, we conducted a study of how the entanglement criterion using the Shannon entropic uncertainty relation is modified by the generalized uncertainty principle. As expected, the upper bounds of the entanglement criteria for pure and mixed states in bipartite systems are increased just like in the variance form [11] from which we can infer as in [11] that the increase in the bounds due to GUP effects enhances entanglement making the quantum effects more pronounced. Intuitively, we expect the introduction of gravitational effects to increase the entropy and hence the uncertainty leading to a higher upper bound for entanglement. It should also be noted that the main difference between the weaker inequality entanglement criteria using the Shannon entropy for pure and mixed states without the GUP (Eq. (31) and Eq. (38)) and with the GUP (Eq. (55) and Eq. (60)) is the state-dependence due to the GUP correction $\eta_i(\beta)$ and $\eta_{im}(\beta)$ defined in Eq. (46) and Eq. (57). However, a state-independent entanglement criterion can apparently be derived using the Rényi and Tsallis entropies since state-independent entropic uncertainty relations with GUP have been obtained in [35]. Since the entropic bounds are increased we expect also an increase in the entanglement criteria upper bounds.

The connection of gravity and entanglement via the entropic uncertainty relations may not be surprising after all. Recently the idea of gravity as an entropic force [37] has been proposed. Gravity can then be thought of as resulting from the tendency of physical systems to increase their entropy. Furthermore, it has been proposed [38] that the entanglement (measured in terms of entanglement von Neumann entropy) of degrees of freedom in quantum systems can give rise to the emergence of spacetime in the gravity picture from field theory with the gravity- gauge theory correspondence (the so called anti-deSitter space-conformal field theory or AdS-CFT correspondence) [39-41]. Whereas these studies indicate the effect of quantum entanglement on gravity, our work opens up yet another avenue of the effect of gravity on quantum entanglement. Gravity and quantum entanglement seem to be mutually related.

As mentioned above, with the GUP correction obtained for the Rényi and Tsallis entropic uncertainty relations in [35], it will be interesting to study carefully the GUP effect on the entanglement criteria based on the Rényi entropy as a generalization of the Shannon entropy. Preliminary work on the GUP effects on the Rényi entropic uncertainty relations has indeed shown an increase in the entanglement criteria upper bound [Gadea O. Derivation of the Rényi Entropic Uncertainty Relation and Entanglement Criterion. 2018; unpublished].

References:

1. Hossenfelder S. Minimal Length Scale Scenarios for Quantum Gravity. *Living Rev. Relativ.* 2013; 16: 2-90.
2. Griffiths D J. *Introduction to Quantum Mechanics*. 2nd ed. Englewood Cliffs, NJ: Prentice-Hall; 2005.
3. Gross D J, Mende P F. String theory beyond the Planck scale. *Nucl. Phys. B.* 1988; 303: 407-454.
4. Amati D, Ciafaloni M, Veneziano G. Can Space-Time Be Probed Below the String Size?. *Phys. Lett. B.* 1989; 216: 41-47.
5. Mende P F, Ooguri H. Borel summation of string theory for Planck scale scattering. *Nucl. Phys. B.* 1990; 339: 641–662.
6. Pedram P. The Minimal Length and the Shannon Entropic Uncertainty Relation. *Adv. High Energy Phys.* 2016; 2016: 5101389.
7. Kempf A, Mangano G, Mann R B. Hilbert space representation of the minimal length uncertainty relation. *Phys. Rev. D.* 1995; 52: 1108–1118.
8. Blado G, Owens C, Meyers V. Quantum Wells and the Generalized Uncertainty Principle. *Eur. J. Phys.* 2014; 35: 065011.
9. Sprenger M, Nicolini P, Bleicher M. An Introduction to Minimal Length Phenomenology. *Eur. J. Phys.* 2012; 33: 853-862.
10. Ali A F, Das S, Vagenas E C. Discreteness of space from the generalized uncertainty principle. *Phys. Lett. B.* 2009; 678: 497–499. ; Ali A F, Das S, Vagenas E C. A proposal for testing quantum gravity in the lab. *Phys. Rev. D.* 2011; 84: 044013–22.; Das S, Vagenas E C. Phenomenological implications of the generalized uncertainty principle. *Can. J. Phys.* 2009; 87: 233–240.; Nozari K, Azizi T. Some aspects of minimal length

- quantum mechanics. *Gen. Relativ. Gravit.* 2006; 38: 735–742.; Vahedi J, Nozari K, Pedram P. Generalized uncertainty principle and the Ramsauer–Townsend effect. *Gravit. Cosmol.* 2012; 18: 211–215.; Blado G, Prescott T, Jennings J, Ceyanes J, Sepulveda R. Effects of the Generalised Uncertainty Principle on Quantum Tunnelling. *Eur. J. Phys.* 2016; 37: 025401.
11. Blado G, Herrera F, Erwin J. Entanglement and the Generalized Uncertainty Principle, *Phys. Essays.* 2018; 31: 397-402.
12. Vijayan J. Quantum Entanglement: A Comprehensive Introduction. IISERM Summer Project Report; 2013. Accessed 30 January 2018. Available: https://www.academia.edu/4195691/Quantum_entanglement_for_the_uninitiated.
13. de la Torre A C, Catuogno P, Ferrando S. Uncertainty and Nonseparability Foundation of *Physics Letters.* 1989; 2: 235-244.
14. Schroeder D V. Entanglement isn't just for spin. *Am. J. Phys.* 2017; 85: 812-820.
15. Eisert J, Plenio M B. Introduction to the basics of entanglement theory in continuous-variable systems. *Int. J. Quantum Inform.* 2003; 01: 479-506.
16. Rigolin G. Uncertainty Relations for Entangled States. *Found. Phys. Lett.* 2002; 15: 293-298.
17. Rigolin G. Entanglement, identical particles and the uncertainty principle. *Commun. Theor. Phys.* 2016; 66: 201-206.
18. Duan L, Giedke G, Cirac J I, Zoller P. Inseparability Criterion for Continuous Variable Systems. *Phys. Rev. Lett.* 2000; 84: 2722-2725.
19. Van Loock P, Furusawa A. Detecting genuine multipartite continuous-variable entanglement. *Phys. Rev. A.* 2003; 67: 052315.
20. Majernik V, Richterek L. Entropic uncertainty relations. *Eur. J. Phys.* 1997; 18: 79-89.
21. Robertson H P. The Uncertainty Principle. *Phys. Rev.* 1929; 34: 163.
22. Coles P J, Berta M, Tomamichel M, Wehner S. Entropic Uncertainty Relations and their Applications. *Rev. Mod. Phys.* 2017; 89: 015002.
23. Bialynicki-Birula I, Rudnicki Ł. Entropic Uncertainty Relations in Quantum Physics. In: Sen K D, editor. *Statistical Complexity*. Heidelberg, Germany: Springer; 2011.
24. Brukner Č, Zeilinger A. Conceptual inadequacy of the Shannon information in quantum measurements. *Phys. Rev. A.* 2001; 63: 022113.
25. Timpson C G. On a supposed conceptual inadequacy of the Shannon information in quantum mechanics. *Stud. Hist. Philos. Sci. B.* 2003; 34: 441-468.

- 552 26. Shannon C E, Weaver W. The Mathematical Theory of Communication. Urbana and
553 Chicago: University of Illinois Press; 1949.
- 554 27. Sharp K, Matschinsky F. Translation of Ludwig Boltzmann's Paper "On the Relationship
555 between the Second Fundamental Theorem of the Mechanical Theory of Heat and
556 Probability Calculations Regarding the Conditions for Thermal Equilibrium". Entropy.
557 2015; 17: 1971-2009.
- 558 28. Planck M. On the Law of Distribution of Energy in the Normal Spectrum. Ann. der Phys.
559 1901; 4: 553.
- 560 29. Bialynicki-Birula I, Mycielski J. Uncertainty Relations for Information Entropy in Wave
561 Mechanics. Commun. Math. Phys. 1975; 44: 129-132.
- 562 30. Heisenberg W. The Physical Principles of the Quantum Theory. New York: Dover; 1930.
- 563 31. Walborn S P, Taketani B G, Salles A, Toscano F, de Matos Filho R L. Entropic
564 Entanglement Criteria for Continuous Variables. Phys. Rev. Lett. 2009; 103: 160505.
- 565 32. Simon R. Peres-Horodecki Separability Criterion for Continuous Variable Systems. Phys.
566 Rev. Lett. 2000; 84: 2726-2729.
- 567 33. Einstein A, Podolsky B, Rosen R. Can Quantum-Mechanical Description of Physical
568 Reality Be Considered Complete?. Phys. Rev. 1935; 47: 777-780.
- 569 34. Cover T A, Thomas J A. Elements of Information Theory. Hoboken, NJ: John Wiley and
570 Sons Inc.; 2006.
- 571 35. Rastegin A E. On entropic uncertainty relations in the presence of a minimal length. Ann.
572 Phys. 2017; 382: 170-180.
- 573 36. Oppenheim J, Wehner S. The Uncertainty Principle Determines the Nonlocality of
574 Quantum Mechanics. Science. 2010; 330: 1072-1074.
- 575 37. Verlinde E. On the Origin of Gravity and the Laws of Newton. J. High Energy Phys.
576 2011; 04: 029.
- 577 38. Van Raamsdonk M. Building up spacetime with quantum entanglement. Gen. Relativ.
578 Gravit. 2010; 42: 2323-2329.
- 579 39. Banks T, Fischler W, Shenker S H, Susskind L. M theory as a matrix model: A
580 conjecture. Phys. Rev. D. 1997; 55: 5112-5128.
- 581 40. Maldacena J M. The large N limit of superconformal field theories and supergravity. Adv.
582 Theor. Math. Phys. 1998; 2: 231-252.
- 583 41. Aharony O, Gubser S S, Maldacena J M, Ooguri H, Oz Y. Large N field theories, string
584 theory and gravity. Phys. Rept. 2000; 323: 183-186.