

# Development of triploid callus of *Hevea brasiliensis* using endosperm

## ABSTRACT

**Aim:** To generate a pathway for development of *Hevea* triploids using endosperm tissue as an explant.

**Study Design:** Standardization and optimisation of various parameters for isolation and culture of endosperm tissue and protoplast. Completely randomized design for data from different treatment. Ploidy of the obtained culture was determined.

**Place and Duration of Study:** Department of Biotechnology, Rubber Research Institute of India and duration of the study 12 months.

**Methodology:** Endosperm tissues were collected from *Hevea* seeds of different developmental stages. Somatic embryogenesis from endosperm tissue using callus mediated embryogenesis and direct method of embryogenesis were carried. Endosperm tissue from both immature and mature seeds was cut into thin slices and subjected to enzymatic digestion for the release of protoplasts. Different concentrations and combinations of cell wall digestion enzymes and osmotic agents experimented. The callus obtained from endosperm tissue was subjected to cytological analysis and flow cytometric analysis.

**Results:** Endosperm tissue from immature fruits was found to be ideal one week between (8-10 weeks) both for somatic embryogenesis and for the release of large amount of protoplasts. Of the two basal media tried, Nitsch medium favoured callus induction, 6 % callus induction from mature endosperm tissue in presence of 2,4-D (6.3  $\mu\text{M}$ ) and Kin (12.1  $\mu\text{M}$ ) and 10 % callus induction from immature endosperm tissue in presence of BA (4.4  $\mu\text{M}$ ) and NAA (2.2  $\mu\text{M}$ ). Direct embryogenesis (2 %) has been obtained from immature endosperm in MS basal medium along with GA<sub>3</sub> (2.0  $\mu\text{M}$ ) and BA (11.1  $\mu\text{M}$ ). A few of the endosperm protoplasts showed division when cultured over K&M medium with NAA (0.1  $\mu\text{M}$ ) 2,4-D (0.2  $\mu\text{M}$ ) and BA (0.4  $\mu\text{M}$ ).

**Conclusion:** Endosperm can be used for the development of triploids of *Hevea brasiliensis*. The ploidy variants i.e. triploids, developed through these *in vitro* techniques can be further used in *Hevea brasiliensis* breeding.

**Keywords:** Endosperm, Triploid, *Hevea* Seeds

## 1. INTRODUCTION

Triploidy, a genomic condition that is favourable for vigor and vegetative productivity have been proved to be beneficial in several crops. Development of triploids with increased biomass is highly

desirable in *Hevea* since it may lead to a reduction in the immaturity period as well as an increase in yield. Triploids are generally sterile. Hence triploid development is suitable for plants where seed is not the economic part. The advantage of triploids can be well exploited in *Hevea* since latex, not seed, is the product of commercial value in this crop. Moreover, development of triploids will lead to seedlessness, which can combat Phytophthora disease thereby making the tree resistant to this disease (Jude et al.,2015)

Traditionally triploids are produced by hybridization between tetraploids and diploids (Xiling et al., 2016). *In vitro* regeneration of plants from the endosperm, the sole naturally occurring triploid plant tissue, offers a direct single step approach for triploid production. Parenchymatous nature of the endosperm and the absence of vascular tissues make it a unique and excellent experimental system for *in vitro* culture studies (Hoshino et al., 2011). The endosperm in angiosperms is formed via double fertilization and triple fusion (i.e., fusion between 3 different haploid nuclei, 1 from the paternal and 2 from the maternal side), which is a unique process in higher plants and is present in all angiosperm families except Orchidaceae, Podostemaceae, and Trapaceae. Endosperm functions as a nutritive tissue for the growing embryo, as the growth and development of the embryo depends on the presence of the endosperm (Paolo and Brian,2009). Moreover, the endosperm exists as a reserve food in some seeds like cereals. The endosperm represents about 60% of the world's food supply. Failure of the endosperm to develop properly leads to the abortion of the embryo. Endosperm may be fully utilized by the developing embryo (non-endospermous), or it may persist in mature seeds (endospermous). Attempts were made from 1930 by the scientist Lampe and Mills to grow young corn endosperm using *in vitro* techniques. Different developmental stages of endosperm from immature to mature stages were used by different workers for the *in vitro* development of triploids. Culture of endosperm protoplasts is yet another option for the production of triploids. Moreover, triploid protoplasts once isolated are useful for the production of aneuploids through fusion with haploid or diploid protoplasts. Also, plant protoplasts provide a unique single cell system to underpin several aspects of modern biotechnology. Hence experiments such as somatic embryogenesis from endosperm tissue using direct and indirect method and Isolation and culture of protoplast from endosperm tissue.

## 2. MATERIALS AND METHODS

### 2.1Explant

Endosperm tissue which is the explant used in this study were collected from seeds of different developmental stages. Broadly these source seeds can be divided into two categories mature and immature seeds, depending on their stage of development.

#### 2.1.1 Mature seeds


Seeds (Fig. 1b) were collected from mature fruits (Fig. 1a) from the field grown trees of *Hevea* clone RR11 105, on the day of dispersal. The hard seed coat was removed mechanically and the seeds were sterilized using 0.1 % mercuric chloride solution with a few drops of Tween 20 for 5 min, followed by

several washes in sterile distilled water. **Washing was done uniformly throughout the experiment.** The inner integuments, as well as the embryos, were removed and the remaining tissue, the endosperm (Fig. 1c), was cut into small pieces and used for the further process.

### **2.1.2 Immature seeds**

Young fruits at different maturity stages of 1-10 weeks (Fig. 1d) were collected from field grown trees of *Hevea* and were surface sterilized using alcohol for 15 min. The developing seeds were separated and cut transversely into two halves (Fig. 1e), of which the half with the micropylar end was cultured in Nitsch medium fortified with growth regulators 2, 4-D (9.0  $\mu\text{M}$ ) and Kin (13.9  $\mu\text{M}$ ) for inducing endosperm development (Fig. 1f).

Endosperm tissue obtained from both mature and immature seeds were used as the source for protoplast isolation and culture and also for callus mediated and direct embryogenesis.



**Fig. 1: Explants used for endosperm culture**

**a) Mature fruit, b) Mature seeds, c) Mature endosperm tissue, d) Immature fruits & corresponding ovules at different developmental stages (1 to 10 weeks), e) Cross section of young seed for half ovulo culture, f) Endosperm developing from cultured immature seeds**

## **2.2 Somatic embryogenesis from endosperm tissue**

### **2.2.1 Callus mediated embryogenesis**

The endosperm tissue from both the sources were cut into small segments and cultured for callus induction in two basal media, MS and Nitsch, fortified with different combinations of the growth regulators 2, 4-D (4.5 - 9.0  $\mu\text{M}$ ), Kinetin (9.3 - 13.9  $\mu\text{M}$ ), BA (2.2 - 8.8  $\mu\text{M}$ ) and NAA (1.1 - 5.3  $\mu\text{M}$ ). Best combination for callus induction was worked out and the calli induced in this medium were

transferred to embryo induction medium after proliferation. Two basal media viz. MS and WPM fortified with GA<sub>3</sub> (0.87- 2.9 µM) and BA (4.4 - 13.3 µM) were used for embryo induction.

### **2.2.2 Direct embryogenesis**

For direct embryogenesis, the endosperm tissue from both the sources were kept in two different embryo induction media, MS and WPM fortified with different levels of GA<sub>3</sub> (0.87- 2.9 µM) and BA (4.4 - 13.3 µM). Cultures were maintained in dark.

## **2.3 Protoplast culture**

### **2.3.1 Isolation of protoplasts**

Endosperm tissue from both immature and mature seeds were cut into thin slices and subjected to enzymatic digestion for the release of protoplasts. Different concentrations and combinations of cell wall digestion enzymes and osmotic agents were experimented with a view to enhance the protoplast yield.

### **2.3.2 Osmoticum**

Experiments were carried out for identifying the suitable osmoticum for protoplast release. Cell and protoplast washing medium (CPW medium\*) was used as the basal medium. Different concentrations (0.2 - 1.0 M) of the sugars and sugar alcohols viz: sucrose, glucose, mannitol and sorbitol were tried for identifying the right osmoticum. Also different combinations of the two sugar alcohols were tried for optimization of osmoticum. In addition, the osmotic stabilizer MES (5mM) was incorporated in all the above solutions and autoclaved at 120°C and 15lb/sq pressure for 15 min and stored at 25°C.

{\*CPW medium- KH<sub>2</sub>PO<sub>4</sub> (27.4 mg l<sup>-1</sup>), KNO<sub>3</sub> (101mg l<sup>-1</sup>), CaCl<sub>2</sub>.2H<sub>2</sub>O (1480 mg l<sup>-1</sup>), MgCl<sub>2</sub>.7H<sub>2</sub>O (276 mg l<sup>-1</sup>), KI (0.16 mg l<sup>-1</sup>), CuSO<sub>4</sub>.5H<sub>2</sub>O (0.025 mg l<sup>-1</sup>), pH- 5.8}

### **2.3.3 Cell wall digestion enzymes**

In order to identify the type and concentration of digestion enzymes for protoplast release, different concentrations and combinations of the enzymes cellulase onozuka RS (0.5, 1.0, 2.0 %) and pectinase Y23 (0.05, 0.1, 0.2, 0.5 %) were tried. The enzymes were dissolved in the osmoticum and the pH was adjusted to 5.7. The solution was filter sterilized using a membrane filter (Millipore 0.22µm), stored at 4°C and thawed to room temperature before use. Thin slices of endosperm tissue were incubated in different enzyme solutions and incubated for different time intervals (2 - 6 h).

### **2.3.4 Protoplast purification**

After enzyme incubation, protoplast suspensions were first filtered through nylon sieves of different mesh sizes (30, 64, 71, 85 and 100 µm) to remove the debris and undigested tissue. The filtrate was then transferred to centrifuge tube and the protoplasts were pelleted by centrifugation at 500 rpm for 2 min.

The supernatant was removed using a pasteur pipette and the pellet was re-suspended in the osmoticum, mixed well and again centrifuged. This was repeated three times to remove traces of enzymes. Finally the purified pellet was re-suspended in 1ml of osmoticum and the suspension was

used for culturing. The protoplast yield per gram fresh weight was determined with the help of a haemocytometer.

$$\text{Yield/gFW} = \frac{\text{Total number of protoplasts from four 1mm squares of haemocytometer}}{\text{Sample volume in 4 squares}} \times \text{Total volume of protoplast suspension}$$

### **2.3.5 Culturing of protoplasts**

Three different basal media viz. MS (Murashige and Skoog, 1962), WPM (Lloyd and McCown, 1980) and KM (Kao and Michayluk, 1975) were used for plating the isolated protoplasts. Sucrose and glucose were provided as carbon source at concentrations of 20 g l<sup>-1</sup> and 10 g l<sup>-1</sup> respectively in all these three media. Different growth regulators combinations of 2, 4-D (0.1 - 1.0 µM), NAA (0.1 - 1.0 µM), Kinetin (0.1 - 1.0 µM) and BA (0.1 - 1.0 µM) were also added. Aliquots of purified protoplasts were plated over these different media combinations and all the cultures were incubated in the dark at 28°C until growth of callus was obtained. They were examined routinely to detect cell division, multiplication, growth and callusing.

### **2.4 Statistical analysis**

All experiments were conducted in completely randomized design (CRD) and analyzed using SPSS 16.0 software. The data was subjected to square root/arc sine transformation and analyzed using ANOVA with a significance of p≤0.05.

### **2.5 Confirmation of ploidy**

#### **2.5.1 Cytological analysis**

The proliferating callus obtained from the endosperm tissue was subjected to cytological analysis (Rekha et al., 1993). Callus with actively dividing cells were pre-treated with 0.2 mM 8-hydroxyquinoline for 5 h at 4°C. After this pretreatment the solution was drained off, the callus was washed with distilled water and transferred to cold freshly prepared fixative, Carnoys fluid II (3:1 ethanolacetic acid), for 48 h at room temperature. Afterwards the fixative was drained off and the callus was washed thoroughly to remove traces of fixative. Then the samples were stained with 1 % Snows carmine for 4 hrs. The samples were smeared in 45% acetic acid with a glass rod and mounted on slides as per standard protocol. The slides were observed under a light trinocular microscope (Leica).

#### **2.5.2 Flow cytometric analysis**

The ploidy analyzer I (Partec GmbH, Germany) was used to determine the ploidy level of the endosperm derived callus. For sample preparation, the callus was crushed in galbriath's\* buffer and kept for 5 min incubation. The suspension containing the nuclei was mixed by pipetting up and down several times and then filtered through a 50 µm nylon mesh. The filtrate containing the nuclear suspension was stained with 50 µg/ml propidium iodide and incubated at room temperature for 5 min. 50 µg/ml RNase was then added and mixed and this mixture was used for ploidy analysis (Rashmi

and Rakhi, 2013). The position of peak G1 nuclei of the control (Diploid callus derived from immature anther) was established at channel 400 on a 1024-channel scale, after which the instrument setting was kept constant and the test samples were run under the same parameters.

### 3. RESULT

#### 3.1 Somatic embryogenesis from endosperm tissue

##### 3.1.1 Callus mediated embryogenesis

Of the two basal media tried, Nitsch medium favored callus induction. Among the different growth regulators, a combination of 2, 4-D and Kin responded towards callus induction from mature endosperm. 6 % callus induction was obtained in a combination of Nitsch medium supplemented with 2, 4-D (6.3  $\mu\text{M}$ ) and Kin (12.1  $\mu\text{M}$ ) (Fig. 2a&b). At higher concentrations of 2, 4-D and Kin there was no callus induction (Table 1). When the mature endosperm tissue was kept in the callus induction medium for longer periods, root organogenesis was observed in a combination of 6.6  $\mu\text{M}$  BA and 4.3  $\mu\text{M}$  NAA (Fig. 2c).

**Table 1. Effect of 2,4-D and Kin on callus induction from mature endosperm tissue**


2,4-D ( $\mu\text{M}$ ) → Kin ( $\mu\text{M}$ ) ↓	4.5	6.3	8.1	9.0
9.3	0.0(0.7)	0.0(0.7)	0.0(0.7)	0.0(0.7) *
11.2	1.0 (1.2)	4.5(2.2)	3.0(1.8)	2.5(1.7)
12.1	2.0(1.5)	6.0(2.4)	4.0(2.1)	3.0(1.8)
13.0	1.0(1.2)	3.5(1.9)	3.0(1.8)	2.5(1.7)
13.9	1.5(1.4)	3.0(1.8)	3.0(1.8)	1.5(1.4)

\*CD=0.31

**\* CD-Coefficient of determination**

**\*callus induction (%) from mature endosperm tissue**

*Data were subjected to square root transformation and transformed means are given in Parenthesis*



**Fig. 2 Culture of mature endosperm**

**a & b) Cultured mature endosperm isolated from mature seeds**

**and induction of callus from this cultured endosperm {2, 4-D (6.3  $\mu$ M) Kin (12.1  $\mu$ M)}**

**c) Root induction from mature endosperm (6.6  $\mu$ M BA and 4.3  $\mu$ M NAA)**

In the case of immature endosperm tissue, a combination of BA and NAA responded towards callus induction whereas with the combination of 2,4-D and Kin no callus initiation occurred, only enlargement of endosperm tissue was observed. 10 % callus induction was obtained in a combination of BA (4.4  $\mu$ M) and NAA (2.2  $\mu$ M) **growth regulators** (Fig. 3a&b). At higher concentrations of these growth regulators, especially BA, the callus induction was nil (Table 2).

**Table 2. Effect of BA and NAA on callus induction from immature endosperm tissue**


BA ( $\mu$ M) $\rightarrow$ NAA ( $\mu$ M) $\downarrow$	2.2	4.4	6.6	8.8
1.1	1.0 (1.2)	6.5 (2.6)	4.5 (2.2)	0.0 (0.7)*
2.2	7.0 (2.7)	10 (3.1)	7.0 (2.7)	0.0 (0.7)
3.2	5.5 (2.4)	9.0 (3.08)	3.0 (1.8)	0.0 (0.7)
4.3	2.5(1.7)	4.0 (2.1)	1.5 (1.4)	0.0 (0.7)
5.3	2.0 (1.5)	0.0 (0.7)	0.0 (0.7)	0.0 (0.7)

CD= 0.23

**\* percentage of callus induction from immature endosperm tissue**

*Data were subjected to square root transformation and transformed means are given in parenthesis*






**Fig: 3 Culture of immature endosperm**

**a &b Cultured immature endosperm and induction**

**of callus from this cultured endosperm {BA (4.4  $\mu$ M) and NAA (2.2  $\mu$ M)}**

### **3.1.2 Direct embryogenesis**

Endosperm tissue isolated from 8 week old fruits was found to be suitable for direct embryogenesis. Embryo induction at a low frequency (2 %) was obtained directly from immature endosperm, when cultured over MS medium fortified with GA<sub>3</sub> (2.0  $\mu$ M) and BA (11.1  $\mu$ M) (Fig.4a&b). These embryos are kept in the same medium for further development. From the mature endosperm tissue, no embryo development could be obtained and later the tissue got dried up.



**Fig.4: Direct embryogenesis**

**a Cultured immature endosperm tissue (8 week old)**

**b Direct embryo induction {GA<sub>3</sub> (2.0  $\mu$ M) and BA (11.1  $\mu$ M)}**



## 3.2 Protoplast isolation and culture

### 3.2.1 Protoplast isolation

Among the different stages of immature fruits cultured, endosperm development was observed in the 8 - 10 week old fruits (Fig. 1d). Protoplasts could be isolated from both mature and immature endosperm tissue, but the yield from immature endosperm was more compared to the mature endosperm tissue.

### 3.2.2 Osmoticum


Among the different osmotic agents tried, mannitol has been found to be the most efficient, followed by sorbitol for maintaining osmotic stability. The optimum concentration of mannitol and sorbitol was in the range 0.6 - 0.8 M and 0.4 - 0.6 M respectively (Table 3). Protoplast yield was more in immature endosperm compared to mature endosperm in both mannitol and sorbitol. In presence of mannitol a protoplast yield of  $17 \times 10^4$  /gfw from mature endosperm and  $28 \times 10^4$  /gfw from immature endosperm were obtained. With sorbitol  $12 \times 10^4$  /gfw from mature endosperm and  $16 \times 10^4$  /gfw from immature endosperm was obtained. No protoplast release occurred with sucrose and glucose.

**Table 3 Effect of sugar alcohols on protoplast release from mature and immature endosperm tissue**

SOURCE	Mannitol (M)					Sorbitol (M)				
	0.2	0.4	0.6	0.8	1	0.2	0.4	0.6	0.8	1
Mature fruit endosperm	4	7	15	17	7	6	10	12	6	3*
Immature fruit endosperm	10	12	26	28	20	10	15	16	12	9

\* Protoplast yield  $\times 10^4$ /gfw


It was observed that a combination of the two sugar alcohols, mannitol and sorbitol, yielded more number of protoplasts when compared to the number of protoplast released when this osmotic agents were added separately. 0.7M mannitol with 0.5M sorbitol was found optimum in both the case of immature and mature endosperm tissue (Fig. 5). The highest yield of protoplast of about  $20 \times 10^4$  /gfw from mature endosperm tissue and in case of immature endosperm tissue,  $34 \times 10^4$  /gfw was obtained.



**Fig.5: Effect of osmoticum on protoplast release**

### **3.2.3 Cell wall digestion enzymes**


Out of the five enzyme combinations tested, 2.0 % cellulase mixed with 0.2 % pectinase was found to be optimum for the release of protoplasts from mature endosperm tissue. A combination of 1.0 % cellulase and 0.2 % pectinase gave highest protoplast yield from immature endosperm. Protoplast yield from other combinations of these enzymes was quite low in both the explants (Fig. 6). A protoplast yield of  $24 \times 10^4/\text{gfw}$  was obtained from mature endosperm when subjected to enzymatic digestion in the enzyme combination E4 whereas protoplast release from immature endosperm tissue was  $36 \times 10^4/\text{gfw}$  in the combination E3.



**Fig.6: Effect of digestion enzymes on protoplast release**

With the optimized parameters of osmotica and enzymes, endosperm tissue isolated from immature seeds was found to be ideal for isolation of protoplasts (Fig.7b) compared to mature seeds. In the case of mature endosperm, even though a good number of protoplasts were released initially but as

the enzymatic digestion proceeded for about 2hrs, the osmoticum turned turbid, thereby rendering estimation of protoplast yield much difficult (Fig. 7a). Turbidity may be due to the release of oil granules from the mature endosperm tissue.



**Fig. 7 Protoplast release from mature and immature endosperm**

**a Mature endosperm tissue**

**a<sub>1</sub>** Initial stage of digestion, **a<sub>2</sub>** After 1 hr of digestion,

**a<sub>3</sub>** After 2hr of digestion


**b Immature endosperm tissue**

**b<sub>1</sub>** Initial stage of digestion, **b<sub>2</sub>** After 1 hr of digestion,

**b<sub>3</sub>** After 2hr of digestion

**3.2.4 Purification and culture of protoplasts**

Endosperm protoplasts isolated from young developing seeds could be purified easily when compared with the mature tissue due to the lack of oil granules. Protoplasts released from mature endosperm tissue are bigger compared to that from immature tissue (Fig. 8a&b). However, due to the presence of oil granules in case of mature tissue, purification process was difficult and the yield was also less. Purification was carried out using 71 $\mu$  mesh size sieve for protoplasts from immature endosperm and 100 $\mu$  mesh size sieve for protoplasts from mature endosperm. Protoplasts from mature endosperm did not respond in culture. A few of the protoplasts from immature endosperm started division in K&M medium supplemented with 0.1  $\mu$ M NAA, 0.2  $\mu$ M 2,4-D and 0.4 $\mu$ M BA (Fig. 8c&d). Further division and microcolony formation from these protoplasts is awaited.



**Fig.8: Purification and culture of released protoplast**

**a Purified protoplast from mature endosperm tissue**


**b Purified protoplast from immature endosperm tissue**

**c & d Protoplast division.**

### **3.3 Ploidy determination**

#### **3.3.1 Cytological analysis**


Cytological studies showed a chromosome count of  $3n=54$  (Fig. 9) in the callus obtained from endosperm tissue, observed at a magnification of X 400. Hence it can be confirmed that the endosperm derived callus is triploid in nature.



**Fig.9: Chromosome count from endosperm derived callus (3n=54)**

### **3.3.2 Flow cytometer**

Using flow cytometer, ploidy of the endosperm derived callus was determined (Fig. 10a&b). The histogram showed fluorescence intensity of nuclei from the endosperm callus at the highest peak at channel 560 which is almost 1.5 times the value of the control diploid callus (400), thus proving this callus to be triploid. This confirms the presence of an extra set of chromosomes in the endosperm derived callus, thereby rendering it to be triploid.



**Fig.10 Histogram showing peaks for**  
**a Diploid (Immature anther derived callus)**  
**b Triploid (Endosperm derived callus)**

## 4. DISCUSSION

Endosperm tissue shows a unique characteristic when compared with other explants in terms of origin, development and ploidy level. In our study both mature and immature endosperm tissue were taken as explants for protoplast culture and for direct and callus mediated embryogenesis.

### 4.1 Somatic embryogenesis from endosperm tissue

#### 4.1.1 Callus mediated embryogenesis

In triploids chromosome pairing during meiosis is disrupted and the unequal segregation of the chromosomes produces aneuploid gametes, thereby significantly decreasing fertility. The most direct route to generate triploid plants is to regenerate shoots or somatic embryos from mature or immature endosperm tissue. This has been achieved in a range of plants (Garg *et al.*, 1996; Bhojwani, 2004).

In the present study callus induction, both from immature and mature endosperm tissue could be obtained at frequencies 10% and 6% respectively. Basal medium was the same, namely Nitsch medium. The growth regulator combinations favoring callus induction were quite different, 2, 4-D (6.3  $\mu\text{M}$ ) + Kin (12.1  $\mu\text{M}$ ) for mature endosperm and BA (4.4  $\mu\text{M}$ ) + NAA (2.2  $\mu\text{M}$ ) for immature endosperm. This may be due to the difference in requirement of the endosperm tissue during the early stage of development and the late/mature stage. Sehgal & Khurana, (1985) had reported callus induction from endosperm of mature fruits in MS medium with IAA (1  $\text{mg l}^{-1}$ ) + BAP (1  $\text{mg l}^{-1}$ ) and also in 2,4-D (1  $\text{mg l}^{-1}$ ) + Kin (1  $\text{mg l}^{-1}$ ). Callus induction from mature endosperm of *Actinidia deliciosa* was obtained in MS basal medium with 2, 4-D (2  $\text{mg l}^{-1}$ ) + Kin (5  $\text{mg l}^{-1}$ ) (Goralski *et al.*, 2005). Also different growth regulator requirements for callus induction and subsequent callus proliferation from immature endosperm of neem (*Azadirachta indica*) have been reported by Thomas and Chaturvedi, (2008). According to them, best callusing (53 %) was obtained when cultured in MS + NAA (5  $\mu\text{M}$ ) + BA (2  $\mu\text{M}$ ) + CH (500  $\text{mg l}^{-1}$ ) whereas the percentage of callus proliferation was maximum (45 %) in MS + 2, 4-D (5  $\mu\text{M}$ ). In a few cultures, root induction was observed from the mature endosperm tissue upon prolonged culture in the callus induction medium fortified with 6.6  $\mu\text{M}$  BA and 7.5  $\mu\text{M}$  NAA. However, shoot induction hasn't so far been obtained in those cultures. Endosperm exhibits the property of organogenesis, as reported by Johri and Bhojwani, (1965) in the case of *Exocarpus cupressiformis*.

Earlier attempts were made by Rekha *et al.*, (2007) for the development of triploids in *Hevea brasiliensis* using mature endosperm tissue, but due to low plant regeneration frequency, no further work was attempted. In 1947 La Rue, for the first time, reported the possibility of obtaining continuously growing tissues from the cultured immature maize endosperm. In general it has been found that mature endosperm requires the initial association of embryo to form callus but immature endosperm proliferates independent of the embryo. Similar observation for mulberry was reported by Thomas *et al.*, (2000). The stages of immature endosperm at the time of culture were normally expressed as days after pollination (Thomas *et al.*, 2000). However, some researchers estimated the endosperm stage in relation to the stage of developing embryo (Walia *et al.*, 2007). When the callus

was transferred to a medium containing BA or Kin, shoot buds differentiated from all over the callus. Maximum regeneration in terms of number of cultures showing shoot buds and number of buds per callus occurred in the presence of 5 M BA (Chaturvedi *et al.*, 2003).

#### **4.1.2 Direct embryogenesis**

Endosperm tissue, being the storage tissue for the developing zygotic embryo, is expected to be amenable to direct embryogenesis under in vitro condition. In our experiment, direct embryo induction from cultured immature endosperm could be obtained in MS medium supplemented with GA<sub>3</sub> (2.0 µM) and BA (11.1 µM). However the frequency was quite low. Hence further standardization needs to be carried out in this direction to perfect a system for direct embryogenesis from the triploid endosperm tissue. Parameters like exact age of the young fruit for endosperm isolation, various media components and other additives need to be standardized. It is technically demanding but the rate of success is generally very low.

#### **4.1.3 Protoplast isolation and culture**

Successful isolation of protoplasts from both immature and mature endosperm tissues could be achieved. Protoplasts obtained from immature endosperm showed division when cultured for callus induction. Isolation procedures that yield highly purified and functional protoplasts have been described for many species. The isolation of plant protoplasts was first reported more than 50 yr ago (Cocking, 1960). Isolated protoplasts allow the study of various metabolic processes. Freshly isolated protoplasts have been proved to be versatile cell systems for studying a broad spectrum of plant physiology, plant cell biology, plant gene engineering, biomechanics, stress responses and cell death controls (Bethke and Jones, 2001, Tena *et al.*, 2001).

Enzymatic isolation of protoplasts using cellulase was first reported in tomato from root tips (Cocking, 1960). The easy availability of commercial, purified enzymes such as cellulase, cellulysin, pectinase, macerozyme, driselase, rhozyme and hemicellulase has now resulted in an increase in the yield and viability of protoplasts and their subsequent response in the culture medium. Commonly a combination of pectinase and cellulase is used to digest the cell walls and to liberate protoplasts (Power and Cocking, 1970). In our experiment using a combination of 1 % cellulase and 0.2 % pectinase, large number of protoplasts could be released from immature endosperm tissue. Higher concentration of the enzyme cellulase (2.0 %) along with 0.2 % pectinase was found effective in the release of protoplasts from mature endosperm tissue. The concentration and combination of enzymes for the isolation depend upon age, genotype and stage of differentiation of the tissue from which the protoplasts are to be isolated (Mukhtar *et al.*, 2012). Protoplasts can be isolated from a variety of tissues, young in vitro-grown plants (Bajaj, 1972), tissues like callus, cell suspension (Sushamakumari *et al.*, 2000a) and explants such as root tips (Xu *et al.*, 1982), hypocotyl, cotyledons (Hammatt *et al.*, 1987) and shoots (Russell and McCown, 1986) and leaves from old or mature plants (Sheen, 2001). In our experiment protoplast isolation was tried from both mature and immature endosperm tissue. Similar combinations of enzymes were used by Takahashi *et al.*, (2004) to isolate endosperm



protoplast from dwarf rice variety. In legumes the most frequently used cellulase is Onozuka R-10. This enzyme proved to be suitable for efficient protoplast release from primary explants of alfalfa (Zafer *et al.*, 1995; Mizukami *et al.*, 2006). Pectinases interact with cells being in different phases of the cell cycle and act like biochemical sorters (Sinha & Caligari, 2004). The stability, viability and further growth of the isolated protoplasts are closely related to the maintenance of a proper osmoticum during isolation and subsequent culture. In general, osmotic potential is adjusted by adding D-Mannitol, sorbitol, glucose or sucrose to the enzyme mixture (Navratilova, 2004).

Generally protoplast burst in hypotonic solution and collapse in hypertonic solution (Ohshima and Tyama, 1989). The use of metabolically active osmotic stabilizers like glucose, sucrose along with metabolically inert mannitol is advantageous for protoplast culture. Such substances will be utilized by the protoplasts for growth and cell wall regeneration (Vasil, 1976). At optimum sugar alcohol and enzyme combination the protoplast yield from mature endosperm tissue was less compared to the immature endosperm tissue. A decrease in yield and viability of protoplasts obtained from later stage endosperm due to starch increase has been reported in maize (Schwall and Feix, 1988) and wheat (Keeling *et al.*, 1989).

Purification of protoplasts from mature endosperm tissue also became difficult due to the presence of oil granules. Thus endosperm tissue from young developing seeds was identified as the suitable explant for protoplast isolation and culture. Helle *et al.*, (2010) studied the characterization of oil and starch accumulation in tubers of *Cyperus esculentus* Var. *Sativus* (*Cyperaceae*) where he reported that at the initial stage the starch starts accumulating and along with the development, sugar and protein levels decrease and oil starts accumulating and lipid and fatty acid composition begins to reflect a storage character. Similar observations were obtained in our experiment showing that in *Hevea*, as maturation proceeds there is accumulation of oil granules. Protoplast yield of  $36 \times 10^4$  gfw was obtained from immature endosperm tissue of *Hevea* and showed division when cultured in the K&M medium.

In *Hevea*, successful protoplast release could be achieved using embryogenic cell suspension derived from immature inflorescence and inner integument of immature fruits (Sushamakumari *et al.*, 2000a). They also reported that the protoplasts, when cultured in KPR liquid medium, underwent division leading to callus formation and embryogenesis. Similar results were obtained by Pongchawee, (2006) who reported first cell division of protoplast when cultured in KM8P medium. The isolated protoplasts from endosperm are generally metabolically active as discussed by Isabel, (1992) who reported stable transformation of protoplast from maize endosperm. Many useful genes expressing in barley are endosperm specific.

#### **4.1.4 Ploidy determination**

The callus obtained from endosperm tissue was triploid in nature as confirmed through both cytology and flow cytometric analyses. Cytology and flow cytometer are two techniques generally used to count the chromosome and the DNA content in many species like coconut (Abraham and Mathew,

2011), kiwifruit (Goralski *et al.*, 2005). In our result a chromosome count of 54 was obtained from endosperm tissue of *Hevea*. *Hevea* is a diploid species with a chromosome count of 36. Hence it can be assumed that the calli obtained from endosperm tissue are triploid in nature and can be used for the in vitro development of triploid plants of *Hevea* through endosperm culture. The result was also supported by the flow cytometer data, showing peak at 560 which is 1.5 times more than the peak value of control.

## 5. CONCLUSION

Callus induction could be achieved from both mature and immature endosperm tissue. Direct embryogenesis has been obtained from the immature endosperm.

Different parameters for isolation and purification of endosperm protoplasts have been standardized. Protoplast release could be obtained from both immature and mature endosperm tissues. Protoplast division could be induced in a few cultures derived from the immature endosperm. Isolation, culture and division of protoplasts from endosperm tissue of *Hevea brasiliensis* is reported for the first time. Triploid nature of endosperm callus ( $3n = 54$ ) was confirmed through cytology as well as flow cytometry.

## REFERENCE

1. Abraham A, Mathew PM. Cytology of coconut endosperm. aob.oxfordjournals.org.2011.
2. Bajaj YPS. Protoplast culture and regeneration of haploid tobacco plants. Am J Bot. 1972; 59:647.
3. Bethke PC, Jones RL. Cell death of barley aleurone protoplasts is mediated by reactive oxygen species. Plant J.2001; 25:19-29.
4. Bhojwani SS. *In vitro* production of triploid plants. In: Goodman RM (ed) Encyclopedia of plant and crop science. Marcel Dekker, New York, 2004; pp 590–593.
5. Chaturvedi R, Razdan MK, Bhojwani SS. An efficient protocol for the production of triploid plants from endosperm callus of neem, *Azadirachta indica* A. Juss. J Plant Physiol.2003; 160:557–564.
6. Cocking EC.A method for the isolation of plant protoplasts and vacuoles. Nature.1960; 187:927- 9.
7. Garg L, Bhandari NN, Rani V, Bhojwani SS. Somatic embryos and regeneration of triploid plants in endosperm cultures of *Acacia nilotica*. Plant Cell Rep.1996; 15:855–858.
8. Goralski G, Popielarska M, Slesak H, Siwinska D, Batycka M. Organogenesis in endosperm of *Actinidia deliciosa* cv. Hayward cultured *in vitro*. Acta Biol Crac Ser Bot.2005; 47:121–128.
9. Hammatt N, Kim HI, Davey MR, Nelson RS, Cocking EC. Plant regeneration from cotyledon protoplasts of *Glycine canascens* and *G. glandestina*. Plant Sci.1987; 48:129-35.
10. Helle T, Salla M, Karl-Erik G, Per H, Marie EO, Leif B, Sten S, Anders S. Characterization of oil and starch accumulation in tubers of *Cyperus esculentus* var. Sativus: A novel model

- 476 system to study oil reserves in non-seed tissues. American Journal of Botany.2010; 97(11):  
477 1884–1893.
- 478 11. Hoshino Y, Miyashita T, Thomas D. *In vitro* culture of endosperm and its application in plant  
479 breeding: approaches to polyploidy breeding. Sci Hortic. 2011; 130(1):1-8.
- 480 12. Isabel D, Pilar C. Isolation of protoplast from developing barley endosperm: a tool for transient  
481 expression studies. Plant Cell Rep.1992; 10:595-598.
- 482 13. Johri BM, Bhojwani SS Growth response of mature endosperm in cultures. Nature.1965;  
483 298:1345–1347
- 484 14. Jude W. Grosser, Frederick G. Gmitter Jr., Manjul Dutt, Milica Calovic, Paul Ling, Bill Castle.  
485 Highlights of the University of Florida, citrus research and education center's comprehensive  
486 citrus breeding and genetics program. ISHS Acta Horticulturae 1065:  
487 10.17660/ActaHortic.2015;1065.50
- 488 15. Kao KN, Michayluk MR. Nutritional requirements for growth of *Vicia hajastana* cells and  
489 protoplasts at a very low population density in liquid media. Planta. 1975; 126:105-110.
- 490 16. Keeling PL, Baird S, Tyson RH. Isolation and properties of protoplasts from endosperm of  
491 developing wheat grain. Plant Sci.1989; 65:55-62.
- 492 17. La Rue CD (1947). Growth and regeneration of the endosperm of maize in culture. Am.J. Bot.  
493 1947; 34:585–586.
- 494 18. Lampel L, Mills CO. Growth and development of isolated endosperms of *Carthamus*  
495 *tinctorius*. Biol. Plant. 1930; 51: 749–753.
- 496 19. Lloyd C, McCown. B (1980) commercially feasible micropropagation of *Mountain laurel*,  
497 *Kalmia latifolia* by use of shoot tip culture. Int. Plant Prop. Soc. Proc.1980; 30:421–427.
- 498 20. Mizukami Y, Kato M, Takamizo T, Kanbe M, Inami S, Hattori K. Interspecific hybrids between  
499 *Medicago sativa* L. and annual *Medicago* containing alfalfa weevil resistance. Plant Cell Tiss.  
500 Org. Cult. 2006; 84:79-88.
- 501 21. Mukhtar I, Bajwa R, Nasim .G Isolation of Mesophyll Protoplasts from Leaves of *Dalbergia*  
502 *sissoo* Roxb. J. Appl. Sci. Environ.2012; 16 (1): 11 – 15.
- 503 22. Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue  
504 cultures. Physiol Plant.1962; 15: 473-497.
- 505 23.Navratilova B. Protoplast Culture and Protoplast fusion focused on Brassicaceae. Hort. Sci  
506 (Prague). 2004; 31:140-157.
- 507 24.Ohshima M, Toyama S. Studies on culture of cells and tissue of crop plants, I. survey on  
508 isolation and culture of protoplast from rice leaf sheath. Jap. J. Crop Sci.1989; 58: 103-110.
- 509 25. Paolo A. Sabelli and Brian A. Larkins. The Development of Endosperm in Grasses1 Plant  
510 Physiol. 2009; 149(1): 14–26.doi: 10.1104/pp.108.129437
- 511 26. Pongchawee K, Nakorn U, Lamseejan S, Poompuang S, Phansiri S. Factors affecting the  
512 protoplast isolation and culture of *Anubias nana* Engler. International Journal of Botany. 2006;  
513 2: 193-200.
- 514 27. Power JB, Cocking EC. Isolation of leaf protoplast: Macromolecule uptake and growth  
515 substance response. J Exp Bot.1970; 21:64-70.
- 516 28. Rashmi RH, Rakhi Chaturvedi. Establishment of dedifferentiated callus of haploid origin from  
517 unfertilized ovaries of tea (*Camellia sinensis* (L) O. Kuntze) as a potential source of total  
518 phenolics and antioxidant activity. In Vitro Cell. Dev.Biol.-Plant.2013; 49:60-69.
- 519 29. Rekha K .Cytogenetic analysis in Kacholam (*Kaempferia galanga*). MSc thesis. 1993;KAU.

- 520 30. Rekha K, Jayashree R, Sushamakumari S, Sankariammal L, Thulaseedharan A. Endosperm  
521 culture in *Hevea brasiliensis*. In: Recent Trends in Horticultural Biotechnology  
522 (Keshavachandran R, Nazeem PA, Girija D, John PS and Peter KV Eds.), New India  
523 Publishers, 2007; p.111-116.
- 524 31. Russell JA, McCown BH. Culture and regeneration of *Populus* leaf protoplasts isolated from  
525 non-seedling tissue. Plant Sci.1986; 46:33-142.
- 526 32. Schwall M, Feix G. Zein promoter activity in transiently transformed protoplasts from maize.  
527 Plant Sci.1988; 56 161-166.
- 528 33. Sehgal CB, Khurana S. Morphogenesis and plant regeneration from cultured endosperm of  
529 *Emblica officinalis* Gaertn. Plant Cell Rep.1985; 4:263–266.
- 530 34. Sheen J. Signal transduction in maize and Arabidopsis mesophyll protoplasts. Plant  
531 Physiol.2001; 127: 1466–1475.
- 532 35. Sinha A, Caligari PDS. Aspects of isolation underpinning mitotic behaviour in lupin  
533 protoplasts.Australian Journal of Botany.2004; 52: 669-676.
- 534 36. Sushamakumari S, Sobha S, Rekha K, Jayasree R, Asokan MP. Influence of growth  
535 regulators and sucrose on somatic embryogenesis and plant regeneration from immature  
536 inflorescence of *H. brasiliensis*. Indian Journal of Natural Rubber Research. 2000a; 13: 19-29.
- 537 37. Takahashi H, Shigeto M, Takehiro M, Kunisuke T. Preparation of Endosperm Protoplasts from  
538 a Dwarf Rice Variety and Transient Expression of Green-Fluorescent Protein. Plant  
539 Biotechnology.2004; 21(2):109-112.
- 540 38. Tena G, Asai T, Chiu WL, Sheen J. Plant MAP kinase signaling cascades. Curr Opin Plant  
541 Biol.2001; 4:392-400.
- 542 39. Thomas TD, Bhatnagar AK, Bhojwani SS. Production of triploid plants of mulberry (*Morus*  
543 *alba* L.) by endosperm culture. Plant Cell Rep.2000; 19:395–399.
- 544 40. Thomas TD, Chaturvedi R.Endosperm culture: a novel method for triploid plant Production.  
545 Plant Cell Tiss Organ Cult.2008; 93:1–14.
- 546 41. Vasil IK. The progress, problems and prospects of plant protoplast research. In: Advances in  
547 agronomy.1976; 28: Brady N.C (ed.). New York: Academic press.
- 548 42.Walia N, Kaur A, Babbar SB. Proliferation and differentiation from endosperms of *Carthamus*  
549 *tinctorius*. Biologia Plantarum.2007; 51 (4): 749-753.
- 550 43. Xiling Wang, Zong-Ming Cheng, Shuang ZH, Fengxiang XU. Breeding Triploid Plants: A  
551 Review. Czech J. Genet. Plant Breed., 52, 2016;2: 41–54
- 552 44. Xu ZH, Davey MR, Cocking EC. Organogenesis from root protoplasts of the forage legumes  
553 *Medicago sativa* and *Trigonella foenum-graecum*. Z. Pflanzenphysiol.1982; 107:231-5.
- 554 45. Zafar Y, Nenz E, Damiani F, Pupilli F, Arcioni S. Plant regeneration from explant and  
555 protoplast derived calluses of *Medicago littoralis*. Plant Cell Tissue and Organ Culture.1995;  
556 41:48.

557

558