

THE ROLE OF SOIL SEED BANK IN FOREST REGENERATION

ABSTRACT

Forest degradation and deforestation disrupts the structure and functions of the ecosystem with negative impacts on biodiversity and natural regeneration of the forests. In Nigeria, forests are lost through burning, shifting cultivation and logging of trees, thus reducing the potential of the forest to naturally regenerate and perform ecosystem functions as expected in a forest reserve. There have been insufficient knowledge about the role soil seed bank plays in forest restoration and this will play an essential role in natural regeneration of forest reserves. Seed bank performs tremendous roles in the restoration of degraded forest reserves, which includes donation of propagules, conservation of genetic diversity, composition of the understorey plant community and recovery of biodiversity. These ecosystem functions are dependent on the dynamics of the soil seed bank as well as favourable environmental conditions.

KEYNOTE: Resilience, Biodiversity, dormancy, dispersal

1. INTRODUCTION

The seed bank can be described as the storage of plant seeds in the superficial soil. Soil seed bank can also be defined as the sum of all surviving seeds in the litter and soil (Newmaster *et al.*, 2006). According to Oke and Odebiyi (2007), soil seed banks are the aggregation of viable seeds in the soil potentially capable of replacing adult plant. Soil seed bank describes and depicts the viable seeds present in the soil (Roberts, 1981). This reservoir, corresponds to the seeds which are yet to germinate, having inherent potential of replacing adult plants, which had disappeared by natural death or not, and potential plants that are susceptible to plant diseases, disturbance and animal consumption (Bakker *et al.*, 1996). The soil at various depths is greatly increased momentarily by diverse medium of seed dispersal (animals, wind, water etc.), which is responsible for the addition of seed to the soil. Seeds are responsible for non-vegetative production of seedlings which could give rise to a naturally regenerated forest, assisted natural regeneration and artificial forest regeneration. Regeneration of forest can be described as the process of rejuvenating or reviving degraded forest ecosystems, degradation resulting from wild fire, human impact (burning activities, logging, farming) and natural disasters (flood, volcano, hurricane etc). One of the key roles played by soil seed bank is that, it is an essential component of plant communities (Oke and Odebiyi, 2007) and can be used to predict secondary succession (Oberhauser, 1997). The seed bank houses seeds which are the basic parts of all plants which carries genetic materials from one generation to another generation, hence, their role in forest regeneration cannot be over-emphasized.

The availability or absence of seeds determines the representation or its complete loss in a natural forest ecosystem. However, when the seed bank is replaced deliberately or consciously for restoration purposes by human efforts, it can be referred to as artificial forest regeneration (plantation forests). **Aside** from housing seeds, the seed bank serves as a living history of non-existing plant species, whose dead traces or remnants could not be found. After subjecting results obtained from the seed bank to data analysis, the future of a forest ecosystem can be pre-determined. Understanding the role of seed bank and its uses will not only add to existing knowledge, but will also assist in regeneration of degraded forests in Nigeria. The composition of seed banks is variable, and is classified as temporary or persistent, when modifying the regeneration of the vegetation during different time of the year. Temporary banks are composed of seeds of short life, which do not present dormancy and are dispersed in time for short periods during the year (Mohammed and Hussein, 2008). Species like *Avena fatua*, *Alopecurus myosuroides*, *Galium aparine*, *Lapsana communis*, *Matricaria perforata*, *Centaurea cyanus*, are classified as temporary, having more than one year of age and reserves of seeds remain in the soil year after year, generally buried into the soil. *Chenopodium album*, *sinapis arvensis*, *Aethusa cynapium*, *Papaver rhoeas*, *Viola arvensis*, *kickia spuria*, *Capsela bursa pastoris* and *Amaranthus retroflexus* are examples of persistent soil seed banks (Cabin and Marshal, 2000). There is an urgent need to naturally restore degraded forest reserves in Nigeria, hence it is expedient to understand the more, the role seed bank plays in the process of natural forest regeneration.

2 DISCUSSION

2.1 Characteristics of the soil seed bank

2.1.1 Soil Seed Bank Density and Seed Dormancy

The success of a seed bank depends on the seed density ready to germinate, when replacement of a plant is necessary and when the environmental conditions for establishment are favourable (Cabin and Marshal, 2000). The longevity of seeds represents a major mechanism of survival of certain weed species, and this leads to a continuous source of emergency. The seed longevity of seeds represents a major mechanism of survival of certain weed species, and this leads to a continuous source of emergency. The seed longevity in the soil varies among species characteristics of the seeds, burial depth, and climatic conditions (Oke, 1993). Mayor and Pyott (1996) presented a study with weed species buried and placed to germinate in different times of the year. After 40 years, the species, *Amaranthus retroflexus*, *Ambrosia eliator*, *Lepidium virginicu*, *Plantago major*, *Portulaca aleracea* and *Rumex crispus* were the originated seedlings.

The seed dormancy is another characteristic that affects the seed bank reservoir. The seed populations of several vegetable species behave in different ways with respect to germination; the weeds produce polymorphic seeds, with a certain proportion that is dormant while the others are not (Baskin and Baskin, 1989). Several internal and external factors prevent seeds from germinating.

Among the internal factors are: the presence of a seed coat, which is a barrier to the penetration of water and oxygen; presence of a biochemical inhibitor in the seed; and immature embryo. Among the external factors, the most common are soil water content and temperature (Esperago and Peco, 1993). Baskin and Baskin, (1989) used the term innate dormancy (primary) and induced dormancy (secondary) characterize the development of the dormancy in the mother plant and after the dissemination in space, respectively. The term enforced dormancy, has been used for the inability of the seeds to germinate due to an environmental restriction, like water deficit, low temperature and poor aeration. However, some seed physiologists do not consider the induced dormancy as an actual dormancy but responds only to favourable conditions for germination. This situation is more conveniently referred to as a case of quiescent seeds. The dormancy represents a mechanism of species preservation in the seed bank, distributing the germination through the year. It can guarantee the species survival in the form of seeds, under adverse conditions, even when the population of plants is completely eliminated (Baskin and Baskin, 1989).

2.1.2. Dispersal Of Seedbank

Most species are capable of long-distance seed dispersal, despite morphological dispersal syndromes that would indicate morphological adaptations primarily for short-distance dispersal (Cwyner and MacDonald 1986, Higgins *et al.* 2003). Assessments of mean migration rates found no significant differences between wind and animal dispersed plants (Wilkinson, 1997). Long-distance migration can also be strongly influenced by habitat suitability (Higgins and Richardson, 1999) suggesting that rapid migration may become more frequent and visible with rapid changes in habitat suitability under scenarios of rapid climate change. The discrepancy between estimated and observed migration rates during re-colonization of northern temperate forests following the retreat of glaciers can be accounted for by the underestimation of long-distance dispersal rates and events (Brunet and von Oheimb, 1998). Nevertheless, concerns persist that potential migration and adaptation rates of many tree species may not be able to keep pace with projected global warming (Collingham *et al.*, 1996). However, these models refer to fundamental niches and generally ignore the ecological interactions that also govern species distributions.

2.2 THE NEED FOR FOREST RESTORATION

2.2.1 Deforestation: a major cause for the need of forest restoration

Deforestation can be defined as the conversion of forest ecosystem (natural and secondary re-growths) to a preferred permanent non-forested land use such as agriculture and urban development (Van Kooten and Bulte, 2000). Deforestation is of major concern for the developing countries of the tropics (Myers, 1994) because it causes rapid reduction of the tropical forests (Barracough and Ghimire, 2000) causing loss of biodiversity and enhancing greenhouse effect (Angelsen, 1999). A plantation of trees established primarily for timber production can also be referred to as a forest according to Food and Agricultural Organisation (FAO 2001) and therefore does not classify natural

forest conversion to plantation as deforestation (but still records it as a loss of natural forests). Deforestation has served a major threat to the forest ecosystems from time immemorial, causing a major loss of biodiversity as well as global climate change. These loss can be attributed to illegal logging of trees for fuelwood and charcoal, wildfire, unregulated burning for harvesting logs, urbanization and natural disasters.

Forest degradation occurs when the ecosystem functions of the forest are degraded but where the area remains forested rather than being cleared (FAO 2005b). Laurance (2001) noted that logging opens mature forest area for colonization. Also, new roads are constructed in the forest during logging (Wilkie *et al.*, 2000). This has the potential of increasing spontaneous colonization of logged forest by agricultural colonist. The ability of such tropical forest ecosystems to recover is limited as high and excessive logging has negative effects on the availability of quality seed germplasm for natural regeneration (Vordzogbe *et al.*, 2005; Makana and Thomas, 2006). Professor Norman Myers, one of the foremost authorities on rates of deforestation in tropical forests described rate of world's deforestation thus: "the annual destruction rates seems set to accelerate further and could as well double in another decade" (Myers, 1992). Hence, an urgent need for forest restoration.

Forest restoration or rehabilitation aims to manage ecosystems for conservation and preservation rather than extraction (Jordan and Lubick, 2011). Four possible methods for restoring forest vegetation exist: (i) natural regeneration, (ii) direct seeding, (iii) planting seedlings, and (iv) incorporating restoration goals in plantation programs (Lamb *et al.*, 1997). Natural regeneration is the cheapest approach for rehabilitating degraded forest ecosystems, provided that the previous disturbance has left some residuals (e.g. soil seed banks, mother trees or root shoots) that can serve as "succession primers" (Uasuf *et al.*, 2009). In frequently disturbed habitats, such as arable fields, vegetation composition is expected to be mainly determined by the seed bank composition (Marks and Mohler, 1985; Levassor *et al.*, 1990; Lavorel *et al.*, 1991; Warr *et al.*, 1993). Inorder to achieve restoration of forests, the soil seed bank performs the following roles:

2.2.2. PROPAGULES DONOR

One of the major role the seedbank plays is donating propagules to the soil ecosystem. This propagules are derived from viable seeds present in or on the soil. Species have two main means by which they adapt to change: they can either disperse by seed or vegetative propagules in the direction of a more favourable environment, or they can change their gene frequencies to favour genotypes (genetic constitutions) that are better adapted to the changed environment (Burdon and Thrall 2001, Reusch *et al.*, 2005). In restoring degraded forests, the first step is to quantify the natural regeneration potential of the forest and this is carried out by determining the population dynamics of the accrued seeds in the soil as well as that of the seedling emergence (Uasuf *et al*, 2009), this will help us to know if the regeneration process should be allowed to take place naturally or further assisted. Assessment of the seedlings emergence from the naturally regenerated soil seedbank will bring about an understanding of the species whose seeds are viable in the seedbank and are able to

donate propagules to the soil (Taiwo *et al.*, 2017). This will help in the conservation of germplasm and biodiversity in the degraded forest ecosystem. This process helps the recovery state of the forest reserve back naturally, giving rise to a secondary forest, however, Taiwo *et al.* (2017) recorded few number seeds of woody flora in the soil of Olokemeji Forest reserve, seedlings which emerged from the soil seed bank were predominantly herbaceous flora (Table 1) while a few seeds of *Tectona grandis* and *Azadiractha indica* were found in the soil substrates which represented the woody flora recorded at the period of the research (Table 2). According to Uasuf *et al.* 2009, when low soil seed bank dynamics is observed, natural regeneration process should therefore be assisted through direct seeding, planting seedlings and manipulation of the site to improve environmental conditions for seedling establishment and growth. Natural regeneration is mostly heterogeneous across landscapes, depending on seed dispersal, viability, dormancy, predation, herbivory, rainfall and topographic, edaphic and light conditions (Khurana and Singh, 2001). Determining the spatial patterns of natural regeneration is therefore also essential in designing an appropriate restoration technique.

2.2.3. CONSERVATION OF GENETIC DIVERSITY

Genetic diversity within species of the soil seed bank is crucial for the natural selection of genotypes within populations and species as they respond or adapt to environmental changes (Burdon and Thrall, 2001; Etterson, 2004; Burger and Lynch, 1995). The potential for evolutionary change has been demonstrated in numerous long-term programmes based on artificial selection (Falconer, 1989), and genetic strategies for reforestation in the presence of rapid climate change must focus on maintaining species diversity and genetic diversity within species (Ledig and Kitzmiller, 1992). In the face of rapid environmental change, it is important to understand that the genetic diversity and adaptive capacity of forested ecosystems depends largely on in-situ genetic variation within each population of a species (Bradshaw, 1991). Species may also adapt through phenotypic plasticity, if their genotype entails a range of permissible responses (with respect to the species morphological, physiological, behavioural or life history strategies and traits) that are suited to the new conditions (Nussey *et al.* 2005). Populations exposed to a rate of environmental change exceeding the rate at which populations can adapt, or disperse, may be doomed to extinction (Lynch and Lande, 1993). Genetic diversity determines the range of fundamental eco-physiological tolerances of a species. It governs interspecific competitive interactions, which, together with dispersal mechanisms, constitute the fundamental determinants of potential species responses to change (Pease *et al.*, 1989; Halpin, 1997). Molecular genetic diversity within a species, species diversity within a forested community, and community or ecosystem diversity across a landscape and bioregion represent expressions of biological diversity at different scales. The basis of all expressions of biological diversity is the genotypic variation found in populations. The individuals that comprise populations at each level of ecological organization are subject to natural selection and contribute to the adaptive capacity or resilience of tree species and forest ecosystems (Muller-Starck *et al.*, 2005). Diversity at each of these levels has fostered natural (and artificial) regeneration of forest ecosystems and facilitated their

adaptation to dramatic climate changes that occurred during the quaternary period (DeHayes *et al.*, 2000), this diversity must be maintained in the face of anticipated changes from anthropogenic climate warming.

2.2.4. COMPOSITION OF THE UNDERSTORY PLANT COMMUNITIES

The seed bank plays a crucial role in determining the composition and spatial structure of understory plant communities in degraded forests following soil disturbances such as wildfire, harvesting and/or logging (Taiwo *et al.*, 2017). This could be derived from the species composition and species richness of the seedlings which emerges from the soil seed bank. While some seeds are visible enough to identify because of their physical attributes, other seeds are tiny and will require some advanced techniques for identifying the seed (Degafi and Berhanu, 2014). The knowledge of the understory plant communities with the canopy/tree community helps in the understanding of the diverse kind of plants that existed prior to perturbation. Invasive species among the understory plant communities could be prevented once identified through early observation from the seedling emergence in a naturally regenerating forest or seed specific identification. Preventing the wide-spread of invasive species can be done early enough if observed from the soil seed bank before displacing the indigenous species.

3.5. RECOVERY OF BIODIVERSITY IN TROPICAL FORESTS

As defined by the Convention on Biological Diversity, “biological diversity” means the variability among living organisms from all sources including terrestrial, marine, and other aquatic ecosystems and the ecological complexes of which they are part; this includes diversity within species, among species, and of ecosystems. Allen and Hoekstra (1992) defined biodiversity even more broadly to include the variety of life at multiple scales of ecological organization, including genes, species, ecosystems, landscapes, and biomes. Biodiversity can be referred to as the variety that exists among all levels of life forms. One of the major roles of regeneration from soil seed bank is recovery of biodiversity in tropical forests as observed in Porto Trombetas, Para state, Brazil after a major disturbance as a result of anthropogenic impact. The role of biodiversity in the resilience of forest-carbon dynamics is evidenced by the specialized species that have evolved and characterize the distinctive forest ecosystems found in the major climatic and forest domains tropical, temperate, and boreal. Over time, evolution results in new plant traits, which through the filter of natural selection, and aided by ecological processes such as dispersal, result in forests comprising species that function optimally under the climatic conditions and disturbance regimes prevalent in each domain. The recovery-ability of vegetation after disturbance is believed to lie mainly in the buried seed population (Kalamees and Zobel, 2002). Resilience in ecosystems is related to the biological diversity in the system and the capacity that it confers to maintain ecosystem processes (Walker 1995).

225 **TABLE 1: Relative Importance Values of seedling emergence in sample plots. Of Olokemeji Forest Reserve, Ogun state Nigeria**

226

S/N	Species	Family	HP1(0-15cm) RIV	HP(15-30cm) RIV	UP(0-15cm) RIV	UP(15-30cm) RIV	FT1(0-15cm) RIV	FT1(15-30cm) RIV	FT2(0-15cm) RIV	FT2(15-30cm) RIV	AP1(0-15cm) RIV	AP1(15-30cm) RIV	AP2(0-15cm) RIV	AP2(15-30cm) RIV
<i>Leptochloa</i>														
1	<i>caerulescens</i>	Poaceae	0.54	0	1.08	0.54	8.66	0	7.57	1.62	0.54	0	0	0
2	<i>Talinum fruticosome</i>	Potulaceae	2.21	2.95	0	0	2.21	0.74	0	0	6.64	2.21	2.21	0.74
3	<i>Phyllantus amarus</i>	Euphobiaceae	4.73	0	1.58	0.78	2.37	2.37	0	1.57	0	0	0.78	0
4	<i>Oldelandia corymbosa</i>	Rubiaceae	0	0	12.29	0.77	0.38	0	0	0	0	0	0	0
5	<i>Mariscus alternifolius</i>	Cyperaceae	1.98	0	1.32	6.61	0	0	0	1.98	0.66	0	0	0
6	<i>Desmodium scopiurus</i>	Fabaceae	1.74	0	0.87	0	0	1.74	0	0	0	0	0	0
7	<i>Euphorbia heterophylla</i>	Euphobiaceae	0	0	1.41	0.7	0	0	0	0	2.11	0	0	0
8	<i>Chromolaena odorata</i>	Asteraceae	0	0	1.09	0	0	0	0	1.09	1.09	0	0	0
9	<i>Cynodon dactylon</i>	Poaceae	0	0	0	0	1.09	1.09	0	0	0	1.09	0	0
10	<i>Brachiara deflexa</i>	Poaceae	0	0	1.09	0	0	0	0	0	2.19	0	0	0
11	<i>Ageratum conysoides</i>	Asteraceae	0	0	0	0	0	0	0	0	2.17	0	0	0
12	<i>Tridax procumbens</i>	Asteraceae	0	0	0	0	0	0	0	0	1.09	0	0	0
13	<i>Vernonia galnensis</i>	Asteraceae	0	0	0	0	0	0	0	0	0	0	1.09	0

227

228 Source: (Taiwo et al., 2017)

229 **Table 2: Relative Importance Value of woody species in study plots.of Olokemeji Forest Reserve, Ogun state, Nigeria**

Species	FT1 RIV	FT2 RIV	HP RIV	UP RIV	AP1 RIV	AP2 RIV
<i>Tectona grandis</i>	15.31	17.22	5.73	16.45	0	0
<i>Albizia zygia</i>	0	0	11	0	3.41	2.56
<i>Azadiracta indica</i>	0	0	5.68	2.84	0	0
<i>Agnossius leocarpus</i>	0	0	6.05	0	0	0
<i>Albizia lebeck</i>	0	0	0	4.95	0	0
<i>Delonix regia</i>	0	0	0	4.39	0	0
<i>Gliricidia sepium</i>	0	0	4.39	0	0	0

230

231 CITATION: Taiwo *et al.*, 2017



233
234 **APPENDIX 1:** Natural regeneration of lowland Amazonian rainforest 18 years after clearcutting.
235 Regrowth primarily from soil seed bank and re-sprouting of harvested trees. Porto Trombetas, Brazil.
236

2A



2B



APPENDIX 2A and 2B: Forest resilience as illustrated by the recovery of mixedwood forest in eastern Canada as a result of red pine plantation on a logged site, with natural infilling by deciduous species over a period of about 50-80 years.

CONCLUSION

In order to restore the forest, studying the role of soil seed bank will bring to awareness the species composition of the seedlings, the historical knowledge of the lost forest as well as the biodiversity of the species of seedlings observed from the soil seed bank, all of these important information will help to proffer the type of forest restoration that can be successfully practised in degraded forests ecosystem.

REFERENCES

1. Allen, T.F.H. and T.W. Hoekstra. 1992. Toward a unified ecology. Complexity in ecological systems series. Columbia University Press, NY, USA.
2. Angelsen, A. 1999. Agricultural expansion and deforestation: modelling the impact of population, market forces and property rights. *Journal of Development Economics*. **58**:185-218.
3. Barraclough, S. L., and K. B. Ghimire. 2000. *Agricultural Expansion and Tropical Deforestation: Poverty, International Trade and Land Use*. Earthscan, Sterling, Virginia, USA.
4. Bakker, J. P. 1996. Nature Management by Grazing and Cutting. Dordrecht Kluwer Academic Publishers. Pp 42
5. Baskin, J. M., and Baskin, C. C. 1989. Physiology of Dormancy and Germination in relation to Seedbank Ecology. Academic Press, Inc. San Diego California. Service (GFIS) project. Kumasi, Ghana. Pp 53-66.
6. Bradshaw, A.D. 1991. Genostasis and the limits to evolution. *Phil. Trans. Royal Soc. Lond.* **333**: 289-305.
7. Brunet, J. and G. von Oheimb. 1998. Migration of vascular plants to secondary woodlands in southern Sweden. *Jour. Ecol.* **86**: 429-438.
8. Burdon, J.J. and Thrall P.H.. 2001. The demography and genetics of host-pathogen interactions. Pages 197-217 in J. Silvertown and J. Antonovics (eds.), Integrating ecology and evolution in a spatial context. Blackwell Science Ltd., London.
9. Burger, R. and M. Lynch. M. 1995. Evolution and extinction in a changing environment: a quantitative genetic analysis. *Evolution* **49**: 151-163.
10. Cabin, R. J. and D. L. Marshall, 2000. The Demographic Role of Soil Seedbank I. Spatial and Temporal Comparison of Below and Above Ground Population of Desert Mustard *Lesquerella fendleri*. *Journal of Ecology* **88**: 283-292.
11. Collingham, Y.C., M.O. Hill and B. Huntley. 1996. The migration of sessile organisms: a simulation model with measurable parameters. *Jour. Vege. Sci.* **7**: 831-846.
12. Cwynar, L.C. and G.M. MacDonald. 1986. Geographical variation of lodgepole pine in relation to population history 1987. *Amer. Natur.* **129**:463-469.
13. Degafi S. and Berhanu A. 2014. Assessment of oil seedbank composition of woody species in Hgumbirda National Forestry Priority area, Northeastern Ethiopia. *Journal of science (MEJS)*. **6** (1): 25-44.

14. DeHayes, D.H., Jacobson Jr., G.L. Schaberg, P.G., Bongarten, B Iverson L. and Dieffenbacher-Krall, A.C. 2000. Forest responses to changing climate: lessons from the past and uncertainty for the future. Pages 495-539 in R.A. Mickler, R.A Birdsey and J.L Hom (eds.), Responses of northern U.S. forests to environmental change. Springer-Verlag, NY.
15. Esperagos, T. and B. Peco, 1993. Mediterranean Pasture Dynamics: The Role of Germination. *Journal of Vegetation Sciences* **4**: 189-194.
16. Etterson, J.R. 2004. Evolutionary potential of *Chamaecrista fasciculata* in relation to climate change. I. clinal patterns of selection along an environmental gradient in the Great Plains. *Evolution* **58**: 1446-1458.
17. Falconer, D.S. 1989. Introduction to quantitative genetics. 3rd ed. Longman, New York.
18. FAO (Food and Agriculture Organization of the United States). 2001. Global forest resources assessment 2000. FAO, Rome, Italy. *FAO Forestry Paper No. 140*
19. FAO (Food and Agriculture Organization of the United States). 2005b. Helping forests take cover: on forest protection, increasing forest cover and future approaches to reforesting degraded tropical landscapes in Asia and the Pacific. FAO Regional Office for Asia and the Pacific, Bangkok, Thailand. RAP Publication No. **2005/13**
20. Jordan, W. R. and Lubick, G. M. 2011. *Making nature whole: A history of ecological restoration*. Society For Ecological Restoration. Washington, DC: Island Press.
21. Higgins, S. I. and D.M. Richardson. 1999. Predicting plant migration rates in a changing world: the role of long-distance dispersal. *Amer. Natur.* **153**: 464-475.
22. Higgins, S.I., R. Nathan and M.L. Cain. 2003. Are long-distance dispersal events in plant usually caused by nonstandard means of dispersal? *Ecology* **84**: 1945-1956.
23. Halpin, P.N. 1997. Global climate change and natural-area protection: Management responses and research directions. *Ecol. Appl.* **7**: 828-843.
24. Kalamees, R., and Zobel, M., 2002. The role of seed bank in gap regeneration in a calcareous grassland community. *Ecology* **83**, 1017–1025.
25. Khurana E., and Singh J.S., 2001. Ecology of seed and seedling growth for conservation and restoration of tropical dry forest: a review. *Environmental Conservation*, **28**: 39-52.
26. Lamb D., Parrotta J., Keenan R. and Tucker N., 1997. Rejoining habitat remnants: restoring degraded rainforest lands. *In: Tropical forest remnants: ecology, management, and conservation of fragmented communities*. The University of Chicago Press, USA, 366-385.
27. Laurance, W.F. 2001: Tropical logging and human invasions. – *Conservation Biology* **15**: 4-5.
28. Lavorel, S., Lebreton, J.D., Debussche, M., and Lepart, J., 1991. Nested spatial patterns in seed bank and vegetation of Mediterranean old-fields. *J.Veg. Sci.* **2**, 367–376.
29. Ledig, F.T. and Kitzmiller, J.H. 1992. Genetic strategies for reforestation in the face of global climate change. *For. Ecol. Manage.* **50**: 153-169.
30. Levassor, C., Ortega, M., and Peco, B., 1990. Seed bank dynamics of Mediterranean pastures subjected to mechanical disturbance. *J. Veg. Sci.* **1**, 339–344.

31. Lynch, M. and Lande, R. 1993. Evolution and extinction in response to environmental change. Pages 234-250 in P.M. Kareiva, J.G. Kingsolver and R.B. Huey (eds.), Biotic interactions and global change. Sinauer Assoc., Sunderland, MA, USA.
32. Makana, J., and Thomas, S.C. (2006): Impacts of Selective Logging and Agricultural Clearing on Forest Structure, Floristic Composition and Diversity and Timber Tree Regeneration in the Ituri Forest, Democratic Republic of Congo. *Biodiversity and Conservation* **15**: 1375-1397.
33. Mayor, J. and Pyott, W. T. 1996. Buried viable seeds in two california bunch grass sites and their bearing on the definition of a flora. *Vegetation* **13**,253-282.
34. Mohammed, H. and Hussein, M. 2008. Temporal and Spatial variation in soil seed bank in Elain Natural Forest Reserve North Kordofan, Sudan. *Conference on International Research on Food Security, Natural Resource Management and Rural Development*. 7th -9th October, 2008. University of Honhenheim.
35. Muller-Starck, G., Ziehe, M., and Schubert, R. 2005. Genetic diversity parameters associated with viability selection , reproductive efficiency, and growth in forest tree species. Pages.87-108 in M. Scherer-Lorenzen, C. Korner and E.-D. Schulze, (eds), Forest diversity and function: temperate boreal systems. Springer- Verlag, Berlin.
36. Myers, N. 1992. The Primary Source: Tropical Forests and Our Future. Norton, New York.
37. Myers, N. 1994. Tropical deforestation: rates and patterns. In: The Causes of Tropical of Tropical Deforestation. The economic and statistical analysis of factors giving rise to the loss of the tropical forest. pp 27-40. UCL Press.
38. Newmaster, S. G., Bell, F. W., Roosenboom, C. R., Cole, H. A. and Towill, W. D. 2006. Restoration of floral diversity through plantations on abandoned agricultural land. *Can. J. For.* **36**: 1218-1235.
39. Nussey, D.H., E. Postma, P. Gienapp and M.E. Visser. 2005. Selection on heritable phenotypic plasticity in a wild bird population. *Science*. 310: 304-306.
40. Oberhauser, U. 1997. Secondary forest regeneration beneath pine (*Pinus kesiya*) plantations in the Northern Thai highlands : a chronosequence study. *Journal Forest Ecology and Management* **256**, 114-120.
41. Oke, S. O. 1993. The Effect of Vegetation Physiognomy on Sediment Yield in Ile-Ife Area of Southwestern Nigeria. Unpublished Ph.D Thesis, Obafemi Awolowo University Ile- Ife, Nigeria.
42. Oke, D.O. and Odebiyi, K.A. 2007. Traditional cocoa-based agroforestry and forest species conservation in Ondo State, Nigeria. *Agriculture, Ecosystems and Environment* **122**:305-311.
43. Pease, C.M., Lande R. and J.J. Bull. 1989. A model of population growth, dispersal and evolution in a changing environment. *Ecology* **70**: 1657-1664.
44. Reusch, T.B.H., A. Ehler, A. Hammerli and B. Worm. 2005. Ecosystem recovery after climatic extremes enhanced by genetic diversity. *Proc. Natl. Acad. Sci., USA* **102**: 2826-2831.
45. Roberts, H. A. 1981. Seed banks in soils. *Advances in Applied Biology* **6**: 155

46. Taiwo, D. M., Olubode, O. S. and Oyelowo, O. J. (2017). Assessment of Soil Seedbank Composition, Woody Species Composition and Structure in Olokemeji Forest Reserve, Ogun State, Nigeria. *International Journal of Applied Research and Technology*. 6(12): 23 – 29.
47. Uasuf A, Tigabu M, Oden PC. 2009. Soil seed banks and regeneration of Neotropical dry deciduous and gallery forests in Nicaragua. *Bois et Forets Des Tropiques*. 299: 49-62.
48. Van Kooten G. C. and Bulte E. 2000. *The Economics of Nature: Managing Biological Assets*. Environmental and Resource Economists, vol. 23(4), pages 472-474, December.
49. Vordzogbe, V., Ataquayefio, D. and Gbogbo, F. 2005. The flora and mammals of the moist semi-deciduous forest zone in the Sefwi-Wiawso District of the Western Region, Ghana. *West African Journal of Applied Ecology*. 8(1).
50. Warr, S.J., Thompson, K., and Kent, M., 1993. Seed banks as a neglected area of biogeographic research: a review of literature and sampling techniques. *Prog. Phys. Geogr.* 17, 329–347.
51. Walker, B. 1995. Conserving biological diversity through ecosystem resilience. *Cons. Biol.* 9: 747-752.
52. Wilkie, D.S., Shaw E., Rotberg, F., Morelli, G., and Auzel, P. 2000: Roads, development, and conservation in the Congo basin. – *Conservation Biology* 14: 1614-1622.
53. Wilkinson, D.M. 1997. Plant colonization: Are wind dispersed seeds really dispersed by birds at large spatial and temporal scales? *Jour. Biogeogr.* 24: 61-65.

APPENDIX

1. Natural regeneration of lowland Amazonian rainforest 18 years after clearcutting. Regrowth primarily from soil seed bank and re-sprouting of harvested trees. Porto Trombetas, Brazil.
2. Forest resilience as illustrated by the recovery of mixedwood forest in eastern Canada as a result of red pine plantation on a logged site, with natural infilling by deciduous species over a period of about 50-80 years.