

Tree Species Diversity and Structure of Eda Forest Reserve, Ekiti State, Nigeria

ABSTRACT

Tropical rainforest are continuously threatened by timber exploitation and conversion to other land uses. In this study, tree species diversity and forest structure of Eda Forest Reserve in Ekiti State, Nigeria, were assessed using systematic line transect and purposive sampling techniques for plot demarcation and data collection. Two transects (2000m long) were laid in secondary forest and encroached farmland in the reserve, while the primary forest fragments were sampled purposively. Twenty sample plots (20m×20m) were laid out on each of the vegetation types. All trees >10cm diameter at breast height (dbh) were identified to species level and enumerated for total height and dbh. Data were analyzed using descriptive statistics such as tables, charts, frequency, percentages and diversity index analysis using paleontological statistics software (PAST 2.14). There were 60 species from 22 families, with Sterculiaceae, Caesalpiniaceae and Moraceae being the most abundant families. Individual tree populations were 380 trees/ha, 280 trees/ha and 137 trees/ha in the primary forest, secondary forest and encroached farmland, respectively. Species composition comprised 39, 38 and 19 species in primary forest, secondary forest and encroached farmland, respectively. *Khaya ivorensis* had the highest relative density in the three vegetation types (19.74%, 24.53% and 27.74%) respectively, while *Ceiba pentandra* had the highest height (53.87m). The mean basal area ranged from 0.36m²/ha (encroach farmland) to 3.18m²/ha (primary forest). Shannon-Wiener Indices were 3.22, 3.14 and 2.51 for the primary forest, secondary forest and encroached farmland, respectively. Eda forest reserve is a heterogeneous ecosystem that had variations in tree population due to anthropogenic activities. The secondary forest and encroached farmland have great potential for recovery if conservation efforts are put in place.

KEYWORDS: Eda Forest Reserve, tree species diversity, forest structure, alpha diversity, beta diversity

1.0 INTRODUCTION

Approximately, one-third of the earth's land area is covered with forests and nearly 50% of this ecosystem is found in the tropical environments of the world (FAO, 2015). These rainforests are complex ecosystems mostly dominated by diverse tree species of various sizes. The tropical rainforests also contain a high level of diversity of other flora and fauna which provide a wide variety of food, fodder, fibre and raw materials for people living in and around the forests. They help maintain biological diversity, ameliorate microclimates, influence hydrological processes and nutrient cycling; support soil conservation, as well as improve air and water quality, while serving as habitats for wildlife (FAO, 2015; Parthasarathy, 2001). In Nigeria, 20-25 % of the rainforest zone had been placed under reservation since the late 1920s and '30s. Over the years, the forest reservations have protected natural ecosystems, conserved biodiversity, preserved ecological processes, enhanced scientific research and education, while maintaining genetic resources of flora and fauna (Awotoye and Adebola, 2013; Olajuyigbe and Adaja, 2014). However, increased anthropogenic activities in the primary forests of the reserves have resulted in serious deforestation and degradation. Consequently, timber harvesting, forestland encroachment for farming, and the establishment of tree crop plantations

45 are threatening the continued existence of most rainforests (Oke and Odebiyi, 2007; Olajuyigbe and
46 Adaja, 2014). The situation is further compounded by the paucity of information on tree species
47 composition and diversity in most of these in-situ conservation areas.


48 Eda forest reserve is one of the 10 forest reserves in Ekiti state, Nigeria. It is endowed with an array of
49 renewable natural resources that have been subjected to high levels of exploitation through legal and
50 illegal means (EKFD, 2006). A section of the forest reserve had been converted to farmland, exotic
51 and indigenous tree species plantations, while 57.7% is still covered by primary and secondary
52 forests (Alo *et al.*, 2014). However, there is limited information on the tree species composition of the
53 remaining primary forest as well as the recovering secondary forest in this forest reserve. Therefore,
54 this study assessed the tree species diversity and forest structure of the encroached farmland,
55 secondary and primary forest areas in Eda forest reserve, Ekiti State, Nigeria.

56 2.0 MATERIALS AND METHODS

57 2.1 Study Area

58 Eda forest reserve was gazetted in 1941 (gazette number 37) with the objective of actualizing
59 biological diversity conservation and environmental protection. This tropical humid forest is a high
60 forest located along latitude 7°41'3"N and 7°47'5"N and longitude 5°36'1"E and 5°37'6"E, at an
61 altitude ranging from 497 to 560 m above sea level (Figure 1). The reserve is bordered by four towns:
62 Orin/Ara Ekiti (North), Eda-Ile Ekiti (West), Omuo Ekiti (East) and Isinbode Ekiti (South). This 906ha
63 forest reserve is divided, administratively, into two parts: the 318ha plantation compartment (Eda I),
64 and the 508ha natural forest (Eda II). The natural forest had been initially protected from exploitation
65 but has recently been encroached by subsistence farmers and timber harvesters. The natural forest
66 was highly stocked with many economic tree species and this is evidenced by the level of exploitation
67 that had taken place, resulting in secondary forest regrowth (EKFD, 2012). The forest reserve has an
68 undulating terrain, which is gently sloped in Northeast direction and as ultisol and oxisol soil types.
69 The bedrock material is underlain with basement complex and contains undifferentiated igneous
70 rocks, laterites and white sand. The reserve experiences two seasons with the wet season occurring
71 from April to October while the dry season occurs from November to March. Hence, the average
72 annual temperature ranges from 21°C - 28°C, average precipitation is 1800mm, while the relative
73 humidity ranges from 56% and 85%. The fragmented primary forest is dense with tree species
74 forming continuous multilayered canopies, while the lower canopies contain climbers, shrubs and
75 herbaceous plant (Alo *et al.*, 2014; EKFD, 2006).

76




78

79 **FIGURE 1: MAP OF EDA FOREST RESERVE IN EKITI STATE, NIGERIA**80 **2.2 Sampling Technique and Data Collection**

81 Systematic line transect technique was used to lay sample plots in secondary forest and encroached
 82 farmlands, while purposive sampling was used to lay plots in the primary forest, following the method
 83 of Duran *et al.* (2006). Thus, two transects (2,000m long and 1m wide) were laid in each of the
 84 secondary forest and encroached farmland. While, the fragmented nature of the primary forest,
 85 resulted in the use of purposive sampling technique for selection of plots. Twenty sample plots
 86 (20m×20m) were laid in alternate positions along each transect at an interval of 200m (Figure 2),

87 while the same number of plots were purposively selected and evaluated in the primary forests. All
 88 woody plants with diameter at breast height (dbh) > 10 cm were identified and their total height and
 89 dbh measured following the method of Adekunle *et al.* (2013).

90



91

92 **FIGURE 2: Sampling procedure for identification and enumeration in the study area.**

93 Where: T = Transects, p = Sample plot

94 **2.3 DATA ANALYSIS**

95 **Tree basal area and volume estimation**

96 The basal area (BA m²) of trees was calculated using Eqn. 1:

$$97 \quad BA = \frac{\pi D^2}{4} \quad (1)$$

98 Where D = Diameter at breast height (m)

99 The total basal area for trees in each sample plot was obtained and used to determine the per hectare
 100 equivalents.

101 The volume of individual trees was estimated using the Eqn. 2:

$$102 \quad V = BAHf \quad (2)$$

103 Where V = Volume of tree (m³), H = Total Height of tree (m) and f = Form factor. Total plot volumes
 104 were obtained by adding the volume of individual trees encountered in each plot and then mean plot
 105 volume was calculated. This was also scaled up to per hectare basis.

106 **Tree species classification and diversity indices**

107 All trees were identified insitu by an experienced forest taxonomist where possible and samples were
 108 compared with voucher specimens in Forest Herbarium Ibadan (FHI), Forestry Research Institute of
 109 Nigeria. Tree species were classified into taxonomic families and number of individuals in each family
 110 was used for species diversity classification. The frequency of occurrence was used to determine tree
 111 species abundance/richness. The diversity indices were determined using paleontological statistics
 112 software (PAST 2.14) (Hammer *et al.*, 2001) and some of them were listed as follows:

113 (i.) Shannon-Wiener diversity index (H¹): This determines both the richness and abundance of each
 114 tree species in the vegetation types using Eqn. 3 (Sanwo *et al.*, 2015):

$$H^2 = \sum_{i=1}^S p_i \ln p_i \quad (3)$$

Where S = Number of tree species in each vegetation type; p_i = proportion of each tree species to the total number of trees in each vegetation type; Ln = the natural logarithm.

(ii.) Relative Density (RD): This determines the number of individual per hectare in the forest types using Eqn. 4 (Adekunle *et al.*, 2013):

$$RD = \frac{n_i}{N} \times 100\% \quad (4)$$

Where n_i = number of individuals of each tree species i and N = total number of individuals in the entire tree population

(iii.) Relative Dominance (RD_o): This determines the level of abundance of individual species over other species in the forest types using Eqn. 5 (Adekunle *et al.*, 2013):

$$RD_o = \frac{(BA_i \times 100\%)}{\sum BA_n} \quad (5)$$

Where BA_i = Basal Area of individual trees belonging to a particular species i and BA_n = Total Basal Area.

(iv.) Margalef's index of species richness (M) was determined using Eqn. 6 (Aighe *et al.*, 2014):

$$M = \frac{(S - 1)}{\ln N} \quad (6)$$

3.0 RESULTS AND DISCUSSION

3.1 RESULTS

3.1.1 Tree Species Composition and Distribution in Eda Forest Reserve

A total of 60 tree species were encountered during the study and 41 of them were absent in encroached farmland. The primary forest had the highest number of tree species (39), followed by the secondary forest (38) while encroached farmland had the least (19). There were 380 trees/ha in the primary forest while secondary forest and encroached farmlands had 280 trees/ha and 137 trees/ha, respectively (Table 1). *Khaya ivorensis* had the highest number of trees in each vegetation type with 75, 65 and 38 trees/ha in the primary forest, secondary forest and encroached farmland respectively. Similarly, *Milicia excelsa*, *Sterculia rhinopetala* and *Triplochiton scleroxylon* had high representations in all vegetation types (Table 1).

Table 1: Tree Species Composition and Distribution in Eda Forest Reserve.

S/N	Species	Family	Primary Forest (No./ha)	Secondary Forest (No./ha)	Encroached Farmland (No./ha)
1	<i>Afzelia bipindensis</i> Harms	Caesalpiniaceae	---	3	---
2	<i>Albizia adianthifolia</i> (Schumach) W. Wight	Mimosaceae	5	3	3

3	<i>Alstonia</i> <i>congensis</i> Engl.	Apocynaceae	10	---	---
4	<i>Alstonia boonei</i> De Wild.	Apocynaceae	---	8	3
5	<i>Aningeria robusta</i> (A. Chev.) Aubrev. & Pellegr	Sapotaceae	5	---	5
6	<i>Antiaris toxicaria</i> Lesch.	Moraceae	10	10	8
7	<i>Blighia sapida</i> K. Konig.	Sapindaceae	---	13	15
8	<i>Bombax buonopozense</i> P. Beauv.	Bombacaceae	15	---	3
9	<i>Brachystegia eurycoma</i> Harms	Caesalpiniaceae	5	---	---
10	<i>Brachystegia kennedyi</i> Hoyle	Caesalpiniaceae	---	---	3
11	<i>Bridelia atroviridi</i> Wild.	Euphorbiaceae	---	3	---
12	<i>Ceiba pentandra</i> (L.) Gaertn.	Bombacaceae	15	8	5
13	<i>Celtis zenkeri</i> Engl.	Ulmaceae	10	8	3
14	<i>Chrysophyllum albidum</i> Linn.	Sapotaceae	5	8	---
15	<i>Cola gigantea</i> A. Chev.	Sterculiaceae	---	3	---
16	<i>Cordea millenii</i> Baker	Bignoniaceae	5	---	---
17	<i>Cynometra megalophylla</i> Harms	Caesalpiniaceae	10	---	---
18	<i>Dialium guineense</i> Willd	Caesalpiniaceae	5	3	---
19	<i>Daniella ogea</i> (Harms) Rolfe ex. Holland	Caesalpiniaceae	5	---	---
20	<i>Diospyros mespiliformis</i> Hoshst.	Ebenaceae	5	---	3
21	<i>Distemona bentamianus</i> Baill.	Caesalpiniaceae	5	---	---
22	<i>Enantia chlorantha</i> Oliv.	Annonaceae	---	3	---
23	<i>Entandrophragma angolensis</i> (Welw.) C. DC.	Meliaceae	---	5	3
24	<i>Etandrophragma cylindricum</i> Sprague	Meliaceae	---	5	---
25	<i>Erythrophylum suaveolens</i> (Guill. & Perr.) Brenan	Caesalpiniaceae	---	3	---
26	<i>Ficus exasperata</i> Vahl	Moraceae	---	8	---
27	<i>Ficus mucoso</i> Welw. Ex. Ficalho	Moraceae	5	---	---
28	<i>Funtumia elastica</i> (Preuss) Stapf.	Apocynaceae	5	3	---
29	<i>Gossweilodendron balsamiferum</i> J.	Caesalpiniaceae	---	3	---
30	<i>Hildergardia barteri</i> (Mast) Kosterm.	Sterculiaceae	5	---	---
31	<i>Hollarrhena floribunda</i> (G. Don) Dur & Schinz	Apocynaceae	---	8	---
32	<i>Khaya ivorensis</i> A. Chev.	Meliaceae	75	65	38
33	<i>Kigelia africana</i> (Lam) Benth	Bignoniaceae	5	3	---
34	<i>Lophira alata</i> Banks ex.	Ochnaceae	10	---	---
35	<i>Lovoa trichilioides</i> Harms	Meliaceae	---	3	---
36	<i>Mansonia altissima</i> A. Chev	Sterculiaceae	5	3	---
37	<i>Milicia excelsa</i> (Welw.) C.C. Berg.	Moraceae	25	15	15
38	<i>Milletia aboensis</i> (Hook. F.) Baker	Papilionaceae	5	---	---
39	<i>Mitragyna ciliate</i> Aubrev & Pellegr.	Rubiaceae	---	3	---

40	<i>Monodora myristica</i> (Gaertn) Dunal	Annonaceae	---	---	3
41	<i>Musanga cecropioides</i> R. Br.	Moraceae	5	8	---
42	<i>Nesogordonia papaverifera</i> (A. Chev.) R. Capuron	Sterculiaceae	5	3	---
43	<i>Newbouldia laevis</i> (P. Beauv.) Seem	Bignoniaceae	---	3	---
44	<i>Parinari excelsa</i> Sabine	Chrysobalanaceae	5	---	---
45	<i>Pentaclethra macrophylla</i> Benth	Mimosaceae	---	3	---
46	<i>Piptadeniastrum africanum</i> (Hook F.) Brenan	Mimosaceae	5	---	---
47	<i>Pterocarpus erinaceus</i> Poir	Papilionaceae	---	3	---
48	<i>Pterygota macrocarpa</i> K. Schum	Sterculiaceae	5	---	---
49	<i>Pycnantus angolensis</i> (Welw) Warb.	Myristicaceae	---	3	3
50	<i>Ricinodendron heudelotii</i> (Baill) Pierre	Euphorbiaceae	5	8	---
51	<i>Sterculia rhinopetala</i> K. Schum	Sterculiaceae	45	25	5
52	<i>Sterculia tragacantha</i> Lindl	Sterculiaceae	5	---	---
53	<i>Strombosia pustulata</i> Oliv.	Olacaceae	5	---	---
54	<i>Terminalia ivorensis</i> A. Chev.	Combretaceae	5	5	3
55	<i>Terminalia superba</i> Engl. & Diels	Combretaceae	10	---	---
56	<i>Pterocarpus osun</i> Craib.	Papilionaceae	5	3	3
57	<i>Tetrapleura tetraptera</i> Taub.	Mimosaceae	5	---	---
58	<i>Triplochiton scleroxylon</i> K. Schum.	Sterculiaceae	10	10	13
59	<i>Xylopia aethiopica</i> (Dunal) A. Rich	Annonaceae	---	3	---
60	<i>Zanthoxylum zanthoxyloides</i> (Lam) Zepern	Rutaceae	5	3	---
Total			380	280	137

144

145 3.1.2 Family Distribution of Trees in Eda Forest Reserve

146 There were 22 families represented by tree species enumerated in Eda forest reserve (Table 2).
 147 Sterculiaceae family (53 trees/ha) had the highest population, followed by Caesalpiniaceae family with
 148 38 trees/ha, while Sapindaceae family had the least of 3 trees/ha. The species from
 149 Chrysobalanaceae and Rutaceae families were found only in the primary forest. However, tree
 150 species from the Annonaceae, Myristicaceae, Rubiaceae and Sapindaceae families were absent in
 151 the primary forest.

152 **Table 2: Family composition and distribution of tree species in Eda forest reserve.**

S/N	Family	Primary Forest	Secondary Forest	Encroached Farmland	No of tree species /ha in each family
1	Annonaceae	---	5	3	8
2	Apocynaceae	10	8	3	21

3	Bignoniaceae	10	5	---	15
4	Bombacaceae	10	3	5	18
5	Caesalpiniaceae	25	10	3	38
6	Chrysobalanaceae	5	---	---	5
7	Combretaceae	10	3	3	16
8	Ebenaceae	5	---	3	8
9	Euphorbiaceae	5	5	---	10
10	Meliaceae	5	10	5	20
11	Mimosaceae	15	5	3	23
12	Moraceae	20	10	5	35
13	Myristicaceae	---	3	3	6
14	Ochnaceae	5	---	---	5
15	Olacaceae	5	---	---	5
16	Papilionaceae	10	5	3	18
17	Rubiaceae	---	5	---	5
18	Rutaceae	5	---	---	5
19	Sapotaceae	10	5	5	20
20	Sterculiaceae	35	13	5	53
21	Ulmaceae	5	3	3	11
22	Sapindaceae	---	1	2	3
Total		195	99	54	348

153

154 3.1.3 Relative Abundance and Diversity Indices of Tree Species in the Primary 155 Forest of Eda Forest Reserve

156 *Khaya ivorensis* had the highest relative density (19.74%), relative dominance (2.42%) and Species
157 Importance Value Index (IVI) (22.16%) in the primary forest (Table 3). This was followed by *Sterculia*
158 *rhinopetala* with relative density of 11.84%, relative dominance of 2.30% and IVI of 14.14%. Twenty
159 seven different tree species had the lowest relative density (1.32%). These included *Albizia*
160 *adanthifolia*, *Brachystegia eurycoma*, *Aningeria robusta*, *Cordea millenii* to mention a few. *Strombosia*
161 *pustulata* had the least relative dominance (0.23%) and species importance value index (1.55%),
162 along with *Ricinodendron heudelotii* which also had the least species IVI (1.55%).

163

164 **3.1.4 Alpha and Beta Diversity Indices of Tree Species in Eda Forest Reserve**

165 Simpson index revealed that the primary forest was the most diverse (0.93), while secondary forest
166 and encroached farmland had indices of 0.92 and 0.87, respectively (Table 4). Similarly, the Shannon
167 Wiener index had the highest value for primary forest (3.22) when compared with secondary forest
168 (3.14) and encroached farmland (2.51). The species evenness revealed that primary forest contained
169 more species (0.88) than the other vegetation types (Table 4). Species richness (Margalef's index)
170 revealed that primary forest was more endowed than other vegetation types with 39 species/ha,
171 followed by secondary forest with 38 species/ha, while 19 species/ha occurred in encroached
172 farmland. However, the fisher alpha index revealed that, secondary forest (11.86) was slightly diverse
173 in species composition than other vegetation types, because the values for primary forest (10.89) and
174 encroached farmland (5.99) were lower.

175 **3.1.5 Growth Variables of Tree Species in Eda Forest Reserve**

176 The encroached farmland had the highest mean dbh (83.35 ± 9.04 cm), while secondary forest had
177 the least (34.60 ± 3.22 cm) in the forest reserve (Table 5). On the other hand, mean basal area was
178 $3.18 \text{ m}^2/\text{ha}$, $0.36 \text{ m}^2/\text{ha}$ and $1.68 \text{ m}^2/\text{ha}$ for primary forest, secondary forest and encroached farmland,
179 respectively. The tree volume followed a similar trend with primary forest being the highest
180 ($122.44 \text{ m}^3/\text{ha}$), followed by encroached farmland ($53.02 \text{ m}^3/\text{ha}$) while secondary forest had the lowest
181 ($13.20 \text{ m}^3/\text{ha}$). The mean height varied from 23.87m - 27.93m across the vegetation types (Table 5).
182 Trees with dbh < 20.99cm and 41 - 50.99cm were only present in secondary forest, while all other
183 diameter class distributions were represented in primary forest and encroached farmland (Figure 3).
184 The highest frequency was observed for trees in the > 60cm diameter class which dominated the
185 primary forest.

S/N	SPECIES NAME	FAMILY	MEAN Height (m)	MEAN DBH (cm)	NUMBE R OF TREES (/ha)	RELA TIVE DENS ITY (%)	RELATI VE DOMIN ANCE (%)	SPEC IES IMPO TANC E VALU E (%)	VOLU ME (m ³ / ha)	BASAL AREA (m ² /ha)	SHAN NON WIENE R (H')
1	<i>Albizia adiantifolia</i>	Mimosaceae	16.7	115.8	5	1.32	0.48	1.80	10.03	0.60	0.057
2	<i>Alstonia congensis</i>	Apocynaceae	26.7	56	10	2.63	3.07	5.70	114.45	3.80	0.096
3	<i>Aningerea robusta</i>	Sapotaceae	21.8	31	5	1.32	1.26	2.57	33.98	1.56	0.057
4	<i>Antiaris toxicaria</i>	Moraceae	49	134.6	10	2.63	10.17	12.80	616.90	12.60	0.096
5	<i>Bombax buonopozense</i>	Bombacaceae	46.77	101.2	15	3.95	6.14	10.08	350.25	7.60	0.128
6	<i>Brachystegia eurycoma</i>	Caesalpiniaceae	32.3	40.1	5	1.32	7.28	8.59	291.17	9.01	0.057
7	<i>Ceiba pentandra</i>	Bombacaceae	53.87	38.5	15	3.95	13.89	17.83	936.85	17.20	0.128
8	<i>Celtis zenkerii</i>	Ulmaceae	25.25	27.4	10	2.63	1.98	4.61	61.50	2.45	0.096
9	<i>Chysophylum albidum</i>	Sapotaceae	32	71	5	1.32	1.60	2.91	63.36	1.98	0.057
10	<i>Cordea millenii</i>	Bignoniaceae	23.4	201.7	5	1.32	0.99	2.31	28.82	1.23	0.057
11	<i>Cynometra megalophylla</i>	Caesalpiniaceae	16.2	77	10	2.63	0.61	3.24	12.05	0.75	0.096
12	<i>Dalium guinensis</i>	Caesalpiniaceae	16.8	64.6	5	1.32	0.37	1.68	7.63	0.45	0.057
13	<i>Daniella ogea</i>	Caesalpiniaceae	38.5	65.2	5	1.32	3.13	4.44	149.10	3.87	0.057

14	<i>Diospyros mespiliformis</i>	Ebenaceae	18.8	34.7	5	1.32	1.20	2.51	27.84	1.48	0.057
15	<i>Distemona bentamianus</i>	Caesalpiniaceae	16.1	32.5	5	1.32	0.33	1.65	6.68	0.41	0.057
16	<i>Ficus mucoso</i>	Moraceae	18.4	168	5	1.32	0.38	1.70	8.70	0.47	0.057
17	<i>Funtumia elastic</i>	Apocynaceae	28.6	101.2	5	1.32	1.88	3.20	66.60	2.33	0.057
18	<i>Hildergadia baterii</i>	Sterculiaceae	19.6	64.9	5	1.32	1.06	2.38	25.81	1.32	0.057
19	<i>Khaya ivorensis</i>	Meliaceae	34.03	132.7	75	19.74	2.42	22.16	114.70	3.00	0.320
20	<i>Kigelia Africana</i>	Bignoniaceae	21.3	96.4	5	1.32	1.15	2.46	30.22	1.42	0.057
21	<i>Lophira alata</i>	Ochnaceae	37.35	61.1	10	2.63	3.67	6.31	171.15	4.55	0.096
22	<i>Mansonia altissima</i>	Sterculiaceae	17.6	227.8	5	1.32	0.89	2.21	19.49	1.11	0.057
23	<i>Melicia excels</i>	Moraceae	45.68	57.9	25	6.58	5.69	12.27	331.70	7.05	0.179
24	<i>Milletia aboensis</i>	Papilionaceae	14.4	39.1	5	1.32	0.36	1.68	6.42	0.45	0.057
25	<i>Musanga cecropioides</i>	Moraceae	9.2	80.5	5	1.32	0.38	1.69	4.28	0.46	0.057
26	<i>Nesogodonia papaverifera</i>	Sterculiaceae	27	101.2	5	1.32	3.20	4.52	107.11	3.97	0.057
27	<i>Parinari excels</i>	Chrysobalanaceae	16.1	71	5	1.32	1.51	2.83	30.11	1.87	0.057
28	<i>Piptadeniastrum africanum</i>	Mimosaceae	28.4	34	5	1.32	1.93	3.24	67.86	2.39	0.057
29	<i>Pterocarpus osun</i>	Papilionaceae	21.4	196.9	5	1.32	0.51	1.83	13.52	0.63	0.057
30	<i>Pterygota macrocarpa</i>	Sterculiaceae	29.3	134.6	5	1.32	1.71	3.03	62.17	2.12	0.057
31	<i>Ricinodendron heudelotii</i>	Euphorbiaceae	16.7	121.9	5	1.32	0.24	1.55	4.92	0.29	0.057

32	<i>Steculia rhinopetala</i>	Sterculiaceae	29.88	172.1	45	11.84	2.30	14.14	94.35	2.85	0.253
33	<i>Steculia tragacanta</i>	Sterculiaceae	46.8	108.2	5	1.32	2.95	4.26	170.81	3.65	0.057
34	<i>Strombosia pustulata</i>	Olacaceae	14	92.6	5	1.32	0.23	1.55	4.01	0.29	0.057
35	<i>Terminalia ivorensis</i>	Combretaceae	47.8	33.7	5	1.32	5.53	6.84	327.11	6.84	0.057
36	<i>Terminalia superb</i>	Combretaceae	29	34.4	10	2.63	3.47	6.10	128.80	4.30	0.096
37	<i>Tetrapleura tetraptera</i>	Mimosaceae	17	155.6	5	1.32	1.39	2.71	29.26	1.72	0.057
38	<i>Triplochyton scleroxylon</i>	Sterculiaceae	43.35	103.7	10	2.63	4.36	6.99	240.10	5.40	0.096
39	<i>Zanthoxylum zanthoxyloides</i>	Rutaceae	14.7	132	5	1.32	0.30	1.62	5.55	0.38	0.057

187 **Table 4: Diversity indices of tree species in Eda forest reserve, Ekiti State, Nigeria**

Diversity Indices	Primary Forest	Secondary Forest	Encroached Farmland
Simpson index (D)	0.93	0.93	0.87
Shannon Wiener index (H _I)	3.22	3.14	2.51
Mergalef's index (d)	6.39	6.57	3.66
Evenness index (E)	0.88	0.86	0.85
Menhinck index	2.00	2.27	1.62
Fisher alpha index	10.89	11.86	5.99
Dominance index (C)	0.07	0.08	0.12

188

189

190 **Table 5: Growth characteristics of trees in Eda forest reserve, Ekiti State, Nigeria**

Growth variable	Primary forest	Secondary forest	Encroached farmland
Mean dbh (cm)	78.58 ± 6.93	34.61±3.22	83.35±9.04
Dominant dbh (cm)	82.00	30.00	140.00
Mean height (m)	27.23 ± 1.90	23.87±1.54	27.93±1.59
Dominant Height (m)	16.70	17.40	28.00
Mean Basal Area (m ² /ha)	3.18±0.57	0.36±0.08	1.68±0.39
Total Basal Area (m ² /ha)	123.86	13.68	31.87
Mean Volume (m ³ /ha)	122.44±29.92	13.20±4.06	53.02±14.67
Total Vol./ha (m ³ /ha)	4775.32	501.49	1007.31

191




Figure 3: Diameter (Dbh) Distribution pattern of Eda Forest Reserve

3.2 DISCUSSION

Tree species composition was highest in the primary forest which had a richer and more diverse tree population than other vegetation types. The primary forest was dominated by the light demanding species, characteristic of the emergent layer in a tropical forest. This tall species provide cover for shade tolerant understorey species (Adekunle *et al.*, 2013; Bobo *et al.*, 2006). The primary forest was characterized by an abundance of lianas which entangled the branches and crowns of larger trees. On the other hand, the secondary forest was in the recovery mode with medium size trees, most of which were < 60cm in diameter. This distribution of diameter across the dbh range is an indication of the high level of exploitation that the forest had experienced (Olajuyigbe and Adaja, 2014). In addition,

the large trees scattered in the encroached farmland were economic species retained to provide shade and protection for farm crops (Oke and Odebiyi, 2007). The 60 tree species from 22 families in the forest reserve, represent the high level of complexity in terms of structure and function in rainforest ecosystems. Fabaceae, Moraceae, and Sterculiaceae have been consistently reported as dominant plant families in Nigerian tropical forests (Adekunle *et al.* 2013; Olajuyigbe and Adaja, 2014).

Khaya ivorensis had the highest relative density of 19.74% and could be regarded as the most abundant species in the forest reserve. The dominance of emergent layer species (such as *Khaya ivorensis*, *Millicia excelsa*) highlights the fact that the forest was a climax old growth forest before exploitation and opening of the forest canopy (Hawthorne *et al.*, 2011). The importance value index (IVI), which combines the attributes of relative density, relative frequency and relative dominance; measures the relative importance of a species in a forest (Anning *et al.* 2009). This study revealed that *Khaya ivorensis* had the highest IVI (22.16 %) indicating that this species was the most abundant in the forest reserve and was closely followed by *Ceiba pentandra* (IVI of 17.83%). This species also had the highest relative dominance value of 13.89% which also presented the species as the indicator species in the reserve. This was followed by *Antiaris africana* with 10.17% and the least relative dominance value of 0.23% was contributed by *Strombosia pustulata*.

The highest mean height (53.87m) was recorded for *Ceiba pentandra* which is an indicator species in tropical rainforest ecosystems. On the other hand, the least height (9.2m) was recorded for *Musanga cecropioides* which is a pioneer species that colonizes clearings and abandoned farmlands (Olajuyigbe and Adaja, 2014). The study revealed that despite the high level of exploitation, Eda forest reserve was a repository of many indigenous tropical hardwood species and had high potential for germplasm conservation.

The Shannon diversity index (H') which characterizes the level of diversity in tropical forests ecosystems has a general limit of 1.5 – 3.5 (Kent and Coker 1992). Hawthorne *et al.* (2011) opined that the H' index was an indication of the high species diversity and reflected the dominance of few tree species in the forest. The H' value for the primary forest was slightly lower than other tropical rainforests. For instance, Parthasarathy (2001) reported $H' = 3.89$, while Adekunle and Olagoke (2010) reported $H' = 4.02$, for rainforests in India and Nigeria, respectively. Nevertheless, Alpha diversity index was highest in the primary forests (Simpson index = 0.93 and $H' = 3.22$). The Sorensen's index indicated the species similarities among vegetation types (Ihuma *et al.*, 2011). Primary forest had a lower Sorensen's index (0.19), indicating it was more similar to secondary forest (0.23) than encroached farmland (0.28). This is evidenced by higher tree population (380 trees/ha) in primary forest when compared to encroached farmland (137 trees/ha). This finding agrees with similar studies such as Sanwo *et al.* (2015), who reported 335 trees/ha from 63 species and belonging to 25 families in a tropical rainforest in southern Nigeria. Also, Aigbe *et al.* (2014) documented 323 trees/ha from 68 species in Afi River forest reserve, Nigeria. However, the stand density of Eda forest reserve was lower than that of tropical Amazonia forests with approximately 1720 trees/ha (Campbell *et al.*, 1992).

The dbh class distribution revealed the structure of a degraded forest (encroached farmland), a secondary and old growth forest. The presence of more trees in the lower dbh classes (Figure 3), highlighted the process of recovery of the tree vegetation in the secondary forest (Boubli *et al.*, 2004; Bobo *et al.*, 2006). This implies that, the secondary forest has relatively good regeneration and recruitment potential which are indications of forest health and vigour.

4.0 Conclusion

This study revealed level of exploitation that had influenced the tree species composition in different vegetation types in Eda forest reserve. Human disturbances had influenced the tree species composition and structural complexity of the forest reserve. Hence, the removal of large trees resulted in tree density and volume fluctuations in secondary forest and encroached farmland. Notwithstanding, comparably high floristic composition and diversity were observed in the secondary forest. Thus, the degraded areas have potential for recovery if encroachment and uncontrolled exploitation are curbed. Hence, there is need for a reconciliation of the demands for conservation with social and economic expectations from Eda forest reserve. Furthermore, interventions such as enrichment planting, and regulated resource utilization could aid the restoration of encroached farmlands.

REFERENCES

- Adekunle, V.A.J., Olagoke, A.O. and Akindele, S.O. (2013). Tree species diversity and structure of a Nigerian strict nature reserve. *Tropical Ecology* **54**(3): 275 – 289.
- Adekunle, V.A.J. and Olagoke, A.O. (2010). The impacts of timber harvesting on residual trees and seedlings in a tropical rain forest ecosystem, southwestern Nigeria. *International Journal of Biodiversity Science, Ecosystem Services and Management* **6** (3-4):131–138.
- Aigbe, H.I., Akindele, S.O. and Onyekwelu, J.C. (2014). Tree species diversity and density pattern in Afi river forest reserve, Nigeria. *International Journal of Scientific and Technology Research* **3**(10): 178 – 185.
- Alo, A.A., Akindele, S.O. and Onyekwelu, J.C. (2014). Development of Land Use Map of Eda Forest Reserve in Ekiti State, Nigeria. *Applied Tropical Agriculture* **20**, 19-23.
- Anning, A., Akyeampong, S., Addo-Fordjour, P., Anti, K., Kwarteng, A. and Tettey, Y. (2009). Floristic composition and vegetation structure of the KNUST Botanic Garden, Kumasi, Ghana. *Journal of Science and Technology* **28** (3):103-122
- Awotoye, O. and Adebola, S. (2013). The changing structural face of Little-Ose Forest Reserve South-West Nigeria. *Journal of Biological and Chemical Research* **30** (2):875-886.
- Bobo, K.S., Waltert, M., Sainge, M., Njokagbor, J., Fermon, H. and Mühlenberg, M. (2006). From forest to farmland. II Species richness patterns of trees and understorey plants along a gradient of forest conversion in Southwestern Cameroon, *Biodiversity and Conservation* **15**: 4097- 4117.

280 Boubli, J.P., Eriksson, J., Wich, S., Hohmann, G. and Mesoscale, B. F. (2004). Transect sampling of
 281 trees in the Lomako-Yekokora interfluviumll, Democratic Republic of the Congo. *Biodiversity and*
 282 *Conservation* **13**: 2399- 2417, 2004.

283 Campbell, D. G., Stone, J. L. and Rosas, J. A. (1992). A comparison of the phytosociology and
 284 dynamics of three flood plain forests of known ages, Rio Jurua, western Brazilian Amazon. *Botanical*
 285 *Journal of the Linnaean Society* **108**: 213-237.

286 Duran, E., Meave, J. A., Lott, D. J. and Segura, G. (2006). Structure and tree diversity patterns at
 287 landscape level in a Mexican tropical deciduous forest. *Boletín de Sociedad Botánica de Mexico* **79**:
 288 43-60.

289 EKFD, (2006). Reports on the contribution of forest products to commercial activities and income
 290 generation in Ekiti State. Ekiti State Forestry Department (EKFD), Ado-Ekiti, Nigeria. *Annual Report*
 291 *on Forest Products* **2**(3): 8 – 21.

292 EKFD, (2012). Report on the update of forest reserves statistics and status in Ekiti state. Ekiti State
 293 Forestry Department (EKFD), Ado Ekiti, Nigeria. *Annual Report on Forest Reserves* **4**(1): 17 – 28.

294 Food and Agriculture Organisation, (FAO) (2015). *Global Forest Resources Assessment 2015*. Desk
 295 Reference FAO, Rome, Italy.

296 Hammer, O., Harper, D.A.T. and Ryan, P.D. (2001). PAST: Paleontological Statistics Software
 297 Package for Education and Data analysis. *Palaeontologia Electronica. Geosciences e – journal* **20**
 298 **(2)**: 324 – 336.

299 Hawthorne, W.D., Marshall, C.A.M., Juam, M. A. and Agyeman, V.K. (2011). The impact of logging
 300 damage on tropical rainforest, their recovery and regeneration. *An annotated Bibliography*. Pp. 47-
 301 121.

302 Ihuma, J.O., Chima, V.D and Chapman, H.M. (2011). Tree species diversity in a Nigerian montane
 303 forest ecosystem and adjacent fragmented forest. *Journal of Agricultural and Biological Science* **17**-
 304 **22**.

305 Oke, D.O. and Odebiyi, K.A. (2007). Traditional cocoa-based agroforestry and forest species
 306 conservation in Ondo State, Nigeria. *Agriculture, Ecosystems and Environment* **122**:305-311.

307 Olajuyigbe, S.O. and Adaja, A.A. (2014). Floristic composition, tree canopy structure and regeneration
 308 in a degraded tropical humid rainforest in southwest Nigeria. *Tanzania Journal of Forestry and Nature*
 309 *Conservation* **84**, 6-23.

310 Parthasarathy, N. (2001). Changes in forest composition and structure in three sites of tropical
 311 evergreen forest around Sengaltheri, Western Ghats. *Current Science* **80** (3):389-393.

312 Sanwo, S.K., Ige, P.O., Sosanya, O.S. and Ogunlaye, O.G. (2015). Tree species diversity and forest
313 stand dynamics in a tropical rainforest in southern, Nigeria. *Malaysian Applied Biology* **44**(2): 65 – 73.