

PERFORMANCE OF BROILER CHICKENS FED BAOBAB, AMARANTHUS AND TIGER NUT DURING A HOT-DRY SEASON

ABSTRACT

This study investigated effects of baobab pulp (*Adansonia digitata*), *Amaranthus hybridus* and tiger nut (*Cyperus esculentus*) on growth performance, cost-benefit indices and nutrient retention of heat stressed broilers. The test ingredients were crude sources of vitamins A, C and E functioning as anti-stress for broilers during a hot-dry season. A total of 240 unsexed broiler chicks were used for the study in a completely randomized design (CRD), which lasted for 8 weeks. The test ingredients were included in the starter and finisher diets at 1%. Diet 1 (T₁) served as the control. T₂, T₃, T₄ and T₅ contained synthetic vitamin C, baobab pulp, amaranthus and tiger nut seed respectively. T₆ contained 0.3% each of baobab, amaranthus and tiger nut. Results from the study revealed that there were significant ($P<0.05$) differences in the performance indices and the nutrient retained by the birds. Broiler chickens fed baobab pulp (crude source of vitamin C, T₃) had significantly more profit ($P<0.05$), better final body weight ($P<0.05$) and crude protein retention ($P<0.05$) than those fed amaranthus (T₄), tiger nut (T₅) and the mixture of the test ingredients (T₆). Feed conversion ratio was at the best (1.58) in T₃ while T₅ gave the worst (1.64) at 4th week. Birds fed T₁ gave the best (1.07) feed conversion ratio while T₄ had the worst (1.26) at 8th week. T₂, T₃ and T₅ had the least (10%) mortality compared with those fed control diet (T₁), amaranthus (T₄) and the mixture (T₆). In conclusion, the study showed that baobab pulp could replace synthetic vitamin C as antioxidant based on growth performance of broilers and its accrued profitability during a hot-dry season in the tropics.

Key words: Performance, broiler, baobab pulp, amaranthus, tiger nut seed

1.0 INTRODUCTION

Poultry production is a part of livestock industry and the fastest means of increasing animal protein supply thus, holding the greatest promise of bridging the prevailing wide animal protein gap in Nigeria through its meat and eggs [1]. It's rather unfortunate that this industry faces problems of inadequate and expensive feedstuffs as well as heat stress particularly during dry season. Heat stress is so important because all classes of poultry are susceptible to it. High ambient temperature and high relative humidity prevailing in the micro-environment of the poultry house in the tropics are unfavourable for efficient poultry production [2]. The result is that the full genetic potential of the birds is often not achieved. Therefore, efforts should be geared towards reducing or alleviating the problem.

Many studies had been carried out to ameliorate heat stress in broilers especially in the tropics. Orusebio and Alu [3] supplemented synthetic vitamin C in the diets of broiler chickens and reported better body weight gain, feed intake, feed conversion ratio and lowered percentage mortality. Such beneficial effects were reported by [4] in broiler chickens when vitamins A, C and E were supplemented in the diets. The problem in using synthetic vitamins

in broiler diets is that their costs add to the overall cost of production which eventually increases the costs of broiler products in the market. Also the problem of availability of these vitamins to farmers in rural areas is another concern. Therefore, there is a need to look inward for cheap and locally available crude sources of these vitamins. Promising amongst the natural sources of the vitamins are Baobab pulp (*Adansonia digitata*, a crude source of vitamin C), *Amaranthus hybridus* (a crude source of vitamin A), and Tiger nut (*Cyperus esculentus*, a crude source of vitamin E). Manfredini *et al* [5] reported that Baobab pulp was high in vitamin C, and [6] reported that its content ranges from 169-500 mg/100g. The latter author [6] also reported that the pulp is rich in both calcium and phosphorus. The anti-nutrients present in baobab pulp, *amaranthus hybridus* leaves and tiger nut tubers are phytate, tannin, saponin, oxalate and cynogenic glycosides. Amaranthus was reported to be an important source of vitamins, especially vitamin A [7] and it was also very rich in minerals and an excellent source of amino acids [8]. Although [8] reported that the leaves contained anti-nutrients, but [9] reported that they are within tolerable limits and can be detoxified by soaking, boiling or frying. Tiger nut is rich in minerals like phosphorus and potassium, and in vitamins like E and C [10]. The tubers are also rich in edible oil which can be used for cooking and in making soap [11]. This study therefore investigated the effects of these crude sources of vitamins A, C and E in the diets of broilers vis-à-vis their performance indices, nutrient retention and cost implication of using them during a hot-dry season.

2.0 MATERIALS AND METHODS

2.1 Site of experiment

The study was carried out at the Poultry Unit of the Faculty of Veterinary Medicine, Usmanu Danfodiyo University, Sokoto, Nigeria. The state is situated between latitudes 12° – 13°58'N and longitudes 4°8' – 6°54'E [12]. The temperature is generally high with annual mean of about 38°C, and it shared border with Niger Republic to the North.

2.2 Experimental Diets

A total of 6 experimental diets were formulated in such a way that the control diet (T₁) had no industrial vitamin C. Baobab pulp, amaranthus leaf and tiger nut were included in both the starter and finisher diets (Tables 1 and 2 respectively). Diets in T₂ contained 1% industrial vitamin C. The diets in T₃, T₄ and T₅ contained 1% each of baobab pulp, amaranthus and tiger nut respectively. T₆ contained a mixture of 0.3% each of baobab pulp, amaranthus and tiger nut.

2.3 Management of Experimental Birds

A total of 240 day-old broiler chicks (Anak strain) were randomly allotted into 6 dietary treatment groups in a completely randomized design. Each treatment group contained 40 chicks of four replicates. The chicks were brooded for a period of 2 weeks during a hot-dry season. Normal routine and management practices as outlined for the tropics by Oluyemi and Roberts [13], were put to use during the study. Feed and water were provided *ad libitum*. All necessary vaccinations and medications were given to the experimental birds as at when due.

2.4 Data Collection

Data were collected on performance indices – body weight gain, feed intake, feed conversion ratio and mortality (Table 4). Also, data were collected on nutrient retention and cost benefit indices (Table 5). Cost benefit indices were carried out based on the values of feedstuffs at the time of experiment. Temperature within the pen was recorded thrice (8am, 1pm and 6pm)

85 daily and the means were calculated and recorded (Table 3). Nutrient retention studies were
86 also carried out at the seventh week. Four birds per treatment were taken into a metabolic
87 cage. The birds were allowed a three-day adjustment period and a day fast before the
88 droppings were collected. They were weighed fresh and dried in an oven at 60°C for 48
89 hours. The dried droppings were ground and taken to Laboratory for proximate analysis.

90 2.5 Statistical Analysis

91 Data collected were analyzed by one-way analysis of variance using SAS [14]. Significant
92 means were separated using Duncan option of the same software and a probability of 5% was
93 considered significant.

94 3.0 RESULTS AND DISCUSSION

95 The effects of supplementing the test materials on the birds were shown in Table 4. At 4th
96 week of the study, broiler chicks fed synthetic vitamin C diet in T₂ had the heaviest (660g)
97 final body weight whereas those fed diet containing the mixture of baobab, amaranthus and
98 tiger nut in T₆ had the lowest (572.50g). At 8th week of the study, broiler chickens in control
99 (T₁) had the highest (1750g) final body weight while the least (1450g) was from birds fed diet
100 T₄ and T₆. Significantly, higher daily feed intake ($P<0.0001$) and weight gain were observed
101 in broiler chicks fed control (T₁) and vitamin C (T₂) diets than those fed other diets at 4th week
102 of the study. Similar trend was also observed in feed intake at 8th week of the study. Also,
103 daily weight-gain of broilers fed baobab pulp (T₃) and amaranthus (T₄) were better ($P<0.05$)
104 than those fed tiger nut (T₅) and the mixture of the test ingredients (T₆) at 4 weeks. Mortality
105 was at highest (30%) in the birds fed control diet (T₁), while the least was 10% for those fed
106 synthetic vitamin C (T₂), Baobab (T₃) and Tiger nut (T₅). Feed conversion ratio was at the
107 best (1.58) in T₃ at age four. Birds fed control (T₁) had the best (1.07) at 8th week and this
108 resulted into the highest final weight recorded by the birds in the treatment. Average
109 temperature throughout the period of study ranged from 30.98°C to 34.79°C (Table 3).

110 Birds fed vitamin C (T₂) and control diet (T₁) had similar performance at 4th week. Broiler
111 chickens fed baobab pulp had comparable growth performance with those fed T₁. However,
112 birds fed vit. C, crude sources of vitamins A (amaranthus), E (tiger nut), and combination of
113 the test ingredients, did not perform well when compared to those fed control diet (T₁) as far
114 as the final weight is concerned. The lowest final body weight was obtained in birds fed diet
115 containing mixture of baobab, amaranthus and tiger nut (T₆). This might be attributed to the
116 saponins in the test ingredients which may have caused growth depression. This was in line
117 with the report of [15] that saponins in the diets caused growth depression. There was no
118 beneficial effect of synthetic vitamin C (T₂), amaranthus (T₄), tiger nut (T₅) and the mixture
119 (T₆) on the final body weight of broiler chickens as they exhibited lower body weight gain
120 when compared to the control.

121 Feed intake obtained in this study was lower than the values reported by [4]. It also
122 contradicted the report of [16] that combination of vitamins C and E promoted feed intake of
123 broilers reared in thermoneutral or heat stress environment. The feed intakes of broilers fed
124 control and vitamin C diets were significantly higher than those fed other diets. This finding
125 was in accordance with the observation of [3] that broilers fed synthetic vitamin C showed
126 better feed intake. There were no significant differences ($P>0.05$) in feed conversion ratio at
127 4th and 8th week of the study. Mortality was at the highest among broilers fed control diet (T₁)
128 which agreed with the findings of [3] and [17]. However, the mortality was lower than the
129 findings reported by [18] when Isa Vedette broiler birds were fed *ad libitum* under chronic
130 heat stress. Therefore, it can be inferred from the rate of mortality among the treatments that

the anti-stress and antioxidant properties of synthetic vitamin C, baobab pulp and tiger nut reduced the rate of mortality among these birds during the hot-dry season.

Nutrient retention and cost-benefit analysis of broilers fed vitamin C, baobab, amaranthus and tiger nut showed significant differences among treatments (Table 5). Ether extract retention of birds fed test ingredients compared favourably with control diet in this study. Broiler chickens fed vitamin C and amaranthus had significantly ($P<0.05$) better crude protein (CP) retention than those fed other diets. Furthermore, Broilers fed baobab and control diets had similar crude protein retention. However, those fed tiger nut and the mixture of the test ingredients had lowest crude protein retention. Crude protein retention of broilers in this study was generally low. The low retention of crude protein obtained in the present study might be due to heat stress experienced by experimental birds because when feed intake declines during heat stress, availability of important nutrients may have been reduced. This corroborated the findings of [19] who found that the true digestibility of 12 amino acids were generally depressed in two rape seed- and two soybean meal- diets when fed to broilers subjected to an increasing ambient temperature exposure from 21 to 32°C. It may equally be due to the presence of anti-nutritional factors like tannins and saponins which could affect feed intake as well as digestion in birds as reported by [20 & 21]. Nutrient retentions obtained in the present study were lower than those in the reports of [22 & 23]. Birds fed vitamin C diet (T_2) had the best crude protein and ash retentions. This might be due to the anti-stress and antioxidant properties of synthetic vitamin C. Birds fed tiger nut diet T_5 had the least crude protein retention probably because of higher crude fibre content of tiger nut. The lower crude protein retention obtained for birds fed tiger nut in this study might have resulted from nutrient intake restriction precipitated by lower feed intake or nutrient dilution effect of crude fibre. This agreed with the observation of [24] who reported implications of restricted feed intake on the growth rate and body weight performance of broiler breeders. Cost benefit indices of heat stressed broiler chickens revealed that the control (T_1) produced maximum profit and best economic efficiency of growth. However, more mortality was observed for control broilers than those fed the test ingredients. Furthermore, birds fed baobab yielded ($P<0.05$) more profit and improved economic efficiency of growth than those fed vitamin C, amaranthus, tiger nut and the mixture of the test ingredients.

4.0 CONCLUSION

In conclusion, the study revealed that during a hot-dry season, especially in the tropics, baobab pulp was an excellent replacement for industrial vitamin C. This was because birds fed baobab pulp performed better than their counterparts fed amaranthus and tiger nut in terms of final body weight, profit and economic efficiency of growth. Furthermore, their performance was similar to those fed control and vitamin C diets. However, broiler chickens fed control diet had higher mortality than those fed vitamin C and baobab.

REFERENCES

1. Egbewande OO, Olorede BR. Performance and cost evaluation of broiler chickens fed African mistletoe (*Tapinanthus bangwensis*) leaf meal. International Journal of Agriculture. 2011;3(1):109-113.
2. Ayo JO, Oladele SB. Road transport stress in food animals in Nigeria. Nigerian Veterinary Journal. 1996;1:49-57.
3. Oruseibio SM, Alu EU. Responses of broiler birds to supplementation of ascorbic acid – vitamin C. Nigerian Poultry Science Journal. 2006;4:12-15.

- 176 4. Ogbamgba KO, Omiete AJ, Wekhe SN. The comparative effects of the administration
177 of vitamins A, E and C on the growth of broilers in the tropics. *Tropical Journal of*
178 *Animal Science*. 2007;10(1-2):93-97.
- 179 5. Manfredini S, Vertuani S, Braccioli E, Buzzoni V. Antioxidant capacity of *Adansonia*
180 *digitata* fruit pulp and leaves. *Acta Phytotherapeutica*. 2002;2:2-7
- 181 6. Anonymous, Baobab Tek-Your natural ingredients supplier. Scientific Facts on
182 Baobab (*Adansonia digitata*). The fruit that spell future. 2006;
183 <http://www.baobabtek.com> Accessed on 30/05/2008
- 184 7. BOSTID. Board on Science and Technology for International Development.
185 Amaranth Modern Prospects for an Ancient Crop. 1984;74
186 <http://www.appropedia.org/amaranth> Accessed on 6/30/2008
- 187 8. Akubugwo IE, Obasi NA, Chinyere GC, Ugbogu AE. Nutritional and chemical value
188 of *Amaranthus hybridus* L. leaves from Afikpo, Nigeria. *Africa Journal of*
189 *Biotechnology*. 2007;6(24):2833-2839.
- 190 9. Ekop AS, Eddy NO. Comparative studies of the level of toxicants in the seed of
191 Indian Almond (*Terminalia catappa*) and African Walnut (*Coula edolis*). *Chem.*
192 *Class Journal*. 2005;2:74-76
- 193 10. Dianne M. Tiger nuts. *Journal of National Vegetable Society*. 2006; www.tiger
194 [nut\Tiger Nuts 1.htm](http://www.tiger-nut.com/TigerNuts1.htm) on 5/14/2008
- 195 11. Rosengarten F. 1984; *The book of Edible Nuts*. Walker & Co. ISBN 0802707699.
- 196 12. Mamman, A.B., Oyebanji, J.O. and Petters, S.W. (2000). *Nigeria: A People United, A*
197 *Future Assured (Survey of States)*. Vol. 2, Gabumo Publishing Co. Ltd. Calabar,
198 Nigeria.
- 199 13. Oluyemi JA, Robert FA. 2000; *Poultry Production in Warm Wet Climates*. Macmillan
200 Publication Ltd. London. 2nd Edition.
- 201 14. SAS. 1999; Statistical Analysis System. User's Guide. SAS Inst. Inc. 6th ed. North
202 Carolina, USA.
- 203 15. Singh KS, Panda B. 1996; *Poultry Nutrition*. Kalyani Publishers, New Delhi,
204 India.
- 205 16. Daudu OM, Ogundipe SO, Orunmuyi M. The role of vitamins in ameliorating the
206 adverse effects of hot weather in poultry production: A Review. *Proc. 14th Annual*
207 *Conf. Anim. Sci. Ass. Nig. (ASAN)*. Sept. 14th-17th LAUTECH, Ogbomoso, Nigeria.
208 2009;369 – 371.
- 209 17. Raja AQ, Qureshi A. Effectiveness of supplementation of vitamin C in broiler feeds
210 in hot season. *Pakistan Vet Journal*. 2000;20:100.

- 211 18. Abu-Dieyeh Z. Effect of chronic heat stress and long-term feed restriction on broiler
212 performance. International Journal of Poultry Science. 2006;5: 185 – 190.
- 213 19. Zuprizal M, Larbier AM, Chagneau Geraert PA. Influence of ambient temperature on
214 true digestibility of protein and amino acids of rape seed and soybean meals in
215 broilers. Poultry Science. 1993;72: 289 – 295.
- 216 20. Liener IE. Antinutritional factors in legumes seeds: state of the art. In: *Recent*
217 *Advances of Research in Antinutritional Factors in Legume Seeds*, [J. Huisman,
218 T.F.B. Vander Poel and I.E. Liener, editors]. Pudoc, Wageningen, the Netherlands.
219 1989;Pp. 6-13.
- 220 21. Bate-Smith EC. Haemanalysis of tannins- the concept of relative astringency.
221 Phytochemistry. 1973; 907-912.
- 222 22. Obun CO, Ayanwale BA. Performance, nutrient digestibility and cost evaluation of
223 raw and roasted Afzelia Africana seed meal fed finisher broiler chicks. Nigerian
224 Journal of Animal Production. 2008;35(1):9-17.
- 225 23. Esonu BO, Udedibie ABI, Emenalom OO, Madumere CC, Uchegbu OA. Evaluation
226 of oil palm (*Elaeis guinensis*) leaf meal as feed ingredient in broiler diets. Nigerian
227 Journal Animal Production. 2008;35(1):32-39.
- 228 24. Mench JA. Broiler Breeders: feed restriction and welfare. World Poultry Science
229 Journal. 2002;58: 23

230 **Table 1: Ingredient composition of experimental broiler starter diets**

Ingredients		Diets (%)					
		T ₁	T ₂	T ₃	T ₄	T ₅	T ₆
235	Maize	55.00	55.00	55.00	55.00	55.00	55.00
236	Ground nut cake	30.00	30.00	30.00	30.00	30.00	30.00
237	Fish meal	3.00	3.00	3.00	3.00	3.00	3.00
238	Blood meal	3.00	3.00	3.00	3.00	3.00	3.00
239	Wheat offal	5.00	4.00	4.00	4.00	4.00	4.10
240	Bone meal	0.80	0.80	0.80	0.80	0.80	0.80
241	Limestone	2.50	2.50	2.50	2.50	2.50	2.50
242	Salt	0.25	0.25	0.25	0.25	0.25	0.25
243	*Vitamin premix	0.25	0.25	0.25	0.25	0.25	0.25
244	Lysine	0.10	0.10	0.10	0.10	0.10	0.10
245	Methionine	0.10	0.10	0.10	0.10	0.10	0.10
246	Industrial Vit. C	-	1.00	-	-	-	-
247	Baobab pulp (Bao)	-	-	1.00	-	-	-
248	Amaran. leaf (Ama)	-	-	-	1.00	-	-

249	Tiger nut (Tig)	-	-	-	-	1.00	-
250	Bao+Ama+Tig	-	-	-	-	-	0.90
251	Total	100.00	100.00	100.00	100.00	100.00	100.00
252	Calculated:						
253	Crude protein (%)	23.39	23.23	23.23	23.23	23.23	23.25
254	ME (Kcal/kg)	2881.21	2868.65	2868.65	2868.65	2868.65	2869.91

255 **Flomix® chick/starter premix composition per 1.25kg: Vitamin A, 10,000,000I.U; Vitamin D3, 2,000,000I.U;*
256 *Vitamin E, 10,000I.U; Vitamin K3, 2,000mg; Vitamin B1, 500mg; Vitamin B2, 5,000mg; Vitamin B6, 300mg;*
257 *Vitamin B12, 10,000mcg; Biotin, 100,000mcg; Niacin, 25,000mg; Pantothenic Acid, 10,000mg; Folic Acid,*
258 *1,000mg; Antioxidant, 125,000mg, Manganese, 100,00mg; Zinc, 50,000mg; Cobalt, 250mg; Iron, 40,000mg;*
259 *Copper, 6,000mg; Iodine, 500mg; Selenium, 100mg; Choline chloride, 150,000mg*

260

261 **Table 2: Ingredient composition of experimental broiler finisher diets**

262							
263	Ingredients	Diets (%)					
264		T ₁	T ₂	T ₃	T ₄	T ₅	T ₆
265							
266	Maize	62.00	62.00	62.00	62.00	62.00	62.00
267	Ground nut cake	30.00	30.00	30.00	30.00	30.00	30.00
268	Fish meal	1.00	1.00	1.00	1.00	1.00	1.00
269	Blood meal	1.00	1.00	1.00	1.00	1.00	1.00
270	Wheat offal	3.00	2.00	2.00	2.00	2.00	2.10
271	Bone meal	0.80	0.80	0.80	0.80	0.80	0.80
272	Limestone	1.50	1.50	1.50	1.50	1.50	1.50
273	Salt	0.25	0.25	0.25	0.25	0.25	0.25
274	*Vitamin premix	0.25	0.25	0.25	0.25	0.25	0.25
275	Lysine	0.10	0.10	0.10	0.10	0.10	0.10
276	Methionine	0.10	0.10	0.10	0.10	0.10	0.10
277	Industrial Vit. C	-	1.00	-	-	-	-
278	Baobab pulp (Bao)	-	-	1.00	-	-	-
279	Amaran. leaf (Ama)	-	-	-	1.00	-	-
280	Tiger nut (Tig)	-	-	-	-	1.00	-
281	Bao+Ama+Tig	-	-	-	-	-	0.90
282	Total	100.00	100.00	100.00	100.00	100.00	100.00
283	Calculated:						
284	Crude protein (%)	20.89	20.74	20.74	20.74	20.74	20.75
285	ME (Kcal/kg)	2981.79	2969.23	2969.23	2969.23	2969.23	2970.49

286 **Flomix® grower premix composition per 1.25kg: Vitamin A, 7,500,000I.U; Vitamin D3, 1,500,000I.U;*
287 *Vitamin E, 7,500I.U; Vitamin K3, 2,000mg; Vitamin B1, 500mg; Vitamin B2, 4,000mg; Vitamin B6, 200mg;*
288 *Vitamin B12, 10,000mcg; Biotin, 50,000mcg; Niacin, 15,000mg; Pantothenic Acid, 6,000mg; Antioxidant,*
289 *125,000mg, Manganese, 100,00mg; Zinc, 50,000mg; Cobalt, 250mg; Iron, 40,000mg; Copper, 6,000mg; Iodine,*
290 *500mg; Selenium, 100mg; Choline chloride, 50,000mg.*

Table 3: Temperature readings during the study on weekly basis

Days	Week/Temp (⁰ C)							
	I	II	III	IV	V	VI	VII	VIII
1	32.33	32.67	34.00	32.83	32.33	29.33	34.67	33.33
2	33.00	30.33	32.50	29.00	31.33	32.00	32.83	34.00
3	32.50	33.67	35.00	30.00	31.67	32.67	33.67	32.33
4	32.00	33.33	35.00	32.00	30.50	32.83	34.33	35.33
5	33.33	33.00	35.33	28.67	31.83	33.83	34.33	34.33
6	32.67	35.00	33.00	31.33	31.67	33.00	33.67	35.33
7	33.00	35.00	32.50	33.00	31.83	34.67	32.33	34.67
Mean	32.69	33.29	33.90	30.98	31.59	32.62	33.69	34.79

Table 4: Growth performance and mortality of broilers fed vitamin C, baobab, amaranthus and tiger nut (g/bird)

Parameters	T1 Control	T2 Vit C	T3 Bao	T4 Amar	T5 Tiger	T6 Bao+Am +Tiger	SEM	P-value
Day old weight (g)	45.50	45.00	45.00	45.50	45.00	44.50	0.31	0.252
Final weight (g)								
4 th week	652.50 ^a	660.00 ^a	627.50 ^a	637.50 ^a	615.00 ^a	572.50 ^b	13.79	0.004
8 th week	1750.00 ^a	1500.00 ^a	1650.00 ^{ab}	1450.00 ^b	1550.00 ^b	1450.00 ^b	63.42	0.020
Daily feed intake (g)								
1-4 weeks	37.65 ^a	38.30 ^a	35.35 ^c	37.09 ^{ab}	36.05 ^{bc}	32.90 ^d	0.40	0.0001
5-8 weeks	66.85 ^a	66.65 ^a	64.78 ^b	65.20 ^b	63.55 ^c	62.53 ^d	0.22	0.0001
Daily weight gain (g)								
1-4 weeks	23.30 ^a	23.57 ^a	22.41 ^{ab}	22.77 ^{ab}	21.96 ^b	20.45 ^c	0.49	0.004
5-8 weeks	62.50	53.57	58.93	51.79	55.36	51.79	2.55	0.078
Feed conversion								
1-4 weeks	1.61	1.62	1.58	1.63	1.64	1.61	0.04	0.425
5-8 weeks	1.07	1.24	1.10	1.26	1.15	1.21	0.20	0.219
Mortality (%)								
	30.00 ^a	10.00 ^b	10.00 ^b	25.00 ^{ab}	10.00 ^b	15.00 ^{ab}	5.00	0.036

307 Means along the same row with uncommon superscripts are significantly different ($P<0.05$)

308

309 **Table 5: Nutrient retention and economic indices of broilers fed vitamin C, baobab,**
310 **amaranthus and tiger nut**

Parameter	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	SEM	P-value
Dry matter	78.19 ^a	78.34 ^a	77.31 ^a	75.79 ^{ab}	73.41 ^c	77.06 ^a	0.42	0.0001
Crude protein	48.91 ^b	49.65 ^a	48.74 ^b	49.58 ^a	47.36 ^c	47.3 ^c	0.11	0.0001
Crude fibre	40.00 ^e	52.32 ^d	65.21 ^b	58.00 ^c	62.50 ^b	68.43 ^a	0.95	0.0001
Ash	40.00 ^{bc}	66.67 ^a	50.04 ^{abc}	60.00 ^{ab}	50.00 ^{abc}	36.36 ^c	6.96	0.052
Ether extract	62.50 ^d	71.43 ^{ab}	68.75 ^{bc}	68.75 ^{bc}	66.67 ^c	74.50 ^a	1.08	0.0001
Total cost (₦)	870.00 ^{ab}	900.80 ^a	868.80 ^{ab}	881.60 ^a	867.20 ^{ab}	835.20 ^b	0.08	0.069
Feed cost (₦/kg)	428.80 ^c	489.60 ^d	392.00 ^c	598.40 ^b	563.20 ^c	652.80 ^a	0.08	0.0001
Total revenue (₦)	1454.40 ^a	1204.80 ^d	1355.20 ^b	1155.20 ^e	1254.40 ^c	1155.20 ^e	0.09	0.0001
Profit (₦)	584.40 ^a	304.00 ^{de}	486.20 ^b	273.60 ^e	387.20 ^c	320.00 ^d	0.09	0.0001
Economic Efficiency of growth	0.67 ^a	0.34 ^d	0.56 ^b	0.31 ^d	0.45 ^c	0.38 ^c	0.01	0.0001

311 Means along the same row with uncommon superscripts are significantly different ($P<0.05$)