

# VEGETATIVE GROWTH PHASE IN LOCAL RICE LANDRACES AS AFFECTED BY MOISTURE STRESS

## Abstract

Rice has been part of the most essential food crop for almost half of the world's rapid growing population. At present, it is a valuable commodity of relevant importance which drives the change in the pattern of crop preference selection both in the urban and as well the rural areas. Water scarcity has been termed the most deepened single physiological and ecological factor controlling the effect of growth and development of plants than any known other factor. Experiments have been undertaken to determine the resultant effects of moisture stress and evaluate plant biomass during the vegetative growth phase in local grown rice landraces in North Eastern Nigeria. Seeds of two different local varieties namely BG doguwa, Mai-zazzabi and NERICA as control were obtained from local farmers as they are the widely grown rice cultivars in the region. There were three treatments of irrigating once in a day (control), irrigating after every 2 days (mild) and after six days (severe), respectively. Data collection was done on the following parameters; plant height (cm), number of leaves, root length (cm), stem diameter (cm), and shoot biomass (g). The present study revealed significant reduction in plant growth and biomass accumulation because of severe water deficit. Mai-zazzabi was observed to be more tolerant to the moisture stress in terms of plant growth compared to the other two tested varieties and was noticed to accumulate higher biomass under the severe moisture stress condition.

Key: Rice; Landraces; Water/Moisture deficit

## 1.0 INTRODUCTION

One of the most manageable and prolific agricultural cropping system recognized in the world is rice farming. This is because, rice farming has adapted to varying environmental ecosystems from lowlands to highlands and from waterlogged swamps to uplands. Rice crop from ancient domestication is known to thrive in deep waterlogged soil been a semi-aquatic plant it requires a huge supply of water for its production. Rice production can be done in varying ecological conditions like irrigated lowlands, rainfed lowlands and highlands and deep-water conditions based on water availability (Hafeez *et al.*, 2007). In the World perception generally, irrigated rice is seen to have accounted for about 55 % of the whole productive farming system in its harvested area whilst contributing to about 75 % total production. Irrigated rice annual productivity has exceeded that of rain fed rice by more than 5 % (Fairhurst & Dobermann, 2002). For the meantime, acquiring resources for rice irrigation farming has come into a decline over the past years due to the rapid increase in development and industrialization which aggravates the problem of water scarcity (Gleick *et al.*, 2002). An estimation of water required to attain 1 kg of rice grain is 1900 liters as reported presently by FAOSTAT (2013).

A distinguishing physiological and as well ecological factor which affects plant growth and development is water deficit, the effect is seen to be more predominant than any other environmental factor affecting plant growth (Kramer and Boyer, 1995). Water supply and availability is a critical requirement for plant but its deficiency results in the plant becoming stressed. At the initial stage of the vegetative phase of growth in plants, water deficiency has been established to be a very important limiting factor as it affects elongation and expansion in growth (Anjum *et al.*, 2003). Slow cellular enlargements, plant tillering capacity and reduction in stem lengths because of inhibited internodal elongation are all caused by water stress.

At different levels in the stages of development such as tillering phase, panicle initiation and heading, the rice plant responds differently to moisture stress (Botwright *et al.*, 2008; Kamoshita *et al.*, 2004), yet, some associated factors of the stress like timing, intensity and duration have harmful effect on

50 plant growth. At the stage of reproduction, Liu et al., (2006) reported that flowering is more exposed to  
51 the intensity of the stress which at the end causes sterility in the spikelet.

52 Ample water is required to grow rice; rice grown in upland areas going rainless for a week and rice  
53 grown in shallow lowlands areas going rainless for 2 weeks will significantly cause a reduction in  
54 yield. Drought periods leading to low production and food scarcity were observed because of water  
55 scarcity on average yield production in rain fed conditions. Climate change onset has contributed to  
56 the intensity of frequency of droughts. Rain fed rice production in more than 23 million hectares of  
57 area in South and South West Asia has been affected by water scarcity (IPCC, 2007). Droughts  
58 recurring in Africa, have affected more than 80 % of the potential 20 million hectares of land set aside  
59 for rain fed lowland rice production (IFPRI, 2010). Increasing crop tolerance to water scarcity would  
60 be the most economic approach for maximizing productivity and to minimize agricultural use of fresh  
61 water resource. Recent studies have shown plants evolving numerous morphological, physiological,  
62 biochemical and molecular strategies to adapt to the adverse climatic effect. To fulfill this objective, a  
63 concise knowledge of the possible mechanisms lying behind water stress environment is a must.

64 Though rice is affected by some environmental factors such as moisture, pH, temperature, soil type  
65 etc. rice plant was observed to develop some mechanism to overcome or escape those unfavorable  
66 conditions. And that's the reason why it is important to study and find out moisture effect on the  
67 growth of such local landraces of rice.

## 68 2.0 METHODOLOGY

### 69 2.1 Experimental site

70 The study was carried out at Federal College of Horticulture, Dadin Kowa nursery site. The College is  
71 situated at Dadin Kowa along Gombe to Biu/Maiduguri road in Yamaltu Deba Local Government Area  
72 of Gombe State. Dadin Kowa is about 35 kilometers away from Gombe and is located in the Sudan  
73 Savannah ecological zone of Nigeria, on latitude 11° 7' 22N, longitude 11° 11' 26E, and on an altitude  
74 of 231 meters above sea level.

75 The plant was grown on a well-established bed in the nursery where required field environmental  
76 factors such as light, CO<sub>2</sub> concentration and temperature conditions were mimicked. Seeds of two  
77 different local varieties namely BG doguwa, Mai-zazzabi and NERICA as control were obtained from  
78 local farmers as they are the widely grown varieties in the region. The local rice varieties have not  
79 been subjected to any data base or germplasm depository as they now been tested for preliminary  
80 researches based on farmer usage and consumers consumption. The soil was dug from the College  
81 nursery site, three beds (for the control and drought treatments) of two by two meters were  
82 established with two replicates each and watered to field capacity for three days, and the seeds were  
83 soaked for a day prior to planting to facilitate germination. Five hills of each variety with 4 seeds per  
84 hill were sown. The experiment was a 3 x 3 factorial. The experimental set-up was a randomized  
85 complete block design (RCBD). The treatments were; irrigating once in a day (control), after every two  
86 days (mild), and after six days (severe), respectively. For the first three weeks the plants were  
87 subjected to daily irrigation with the same amount of water per bed. The beds were kept weed free by  
88 handpicking the weeds and also weeding with hoe.

### 89 2.2 Data Measurements

90 All experimental plants were subjected to before (data taken from 1 week after germination till  
91 harvest) and after harvesting agronomic measurement of data as listed below.

92 **Plant height (cm):** Measurement commenced twenty-one (21) days after planting and subsequent  
93 measurements were taken after every 7 days.

94 **Number of leaves:** This was done by counting the number of leaves on each individual tiller of the  
95 plant.

96 **Stem diameter (cm):** The stem diameter was measured using a thread which is tied to the stem of  
97 the plant and then placed on a meter rule to take the measurement.

98 **Wet shoot weight (g):** This was determined immediately after harvesting using a mini kitchen scale  
 99 weighing balance.

100 **Dry shoot biomass weight (g):** The shoots were harvested and then neatly folded and placed in  
 101 brown paper bags and placed to dry at 80°C in an oven, the shoot biomass was then weighed using  
 102 an electronic weighing scale in the laboratory.

103 **Root length (cm):** The plants were gently uprooted and soaked in water to wash off soil particles.  
 104 The length of the root was determined by using a meter rule. Measurements were taken from the  
 105 stem base to the longest root tip of the tap root.

106 **Dry root biomass weight (g):** The roots were harvested and then neatly folded and placed in brown  
 107 paper bags and placed to dry at 80°C in an oven, the roots were then weighed. The root biomass was  
 108 then weighed using an electronic weighing scale in the laboratory.


### 109 3.0 DATA ANALYSIS

110 Data collected were subjected to analysis of variance (ANOVA) using statistical computer package  
 111 Minitab(c) V. 17 (State College PA) to determine treatments effects if significant. The treatment and  
 112 variety mean were separated using the test known as Tukey Pairwise Comparisons. The level of  
 113 significance was set at 95 % confidence interval ( $P \leq 0.05$ ).  
 114 Results presented are data collected and analysed from the fifth week after germination as no  
 115 significant differences ( $P > 0.05$ ) or effects were observed before then on all parameters measured.

## 116 3.1 RESULTS

### 117 3.1.1 Plant heights (cm)

118 A general decline in plant height was observed in relation to increasing water deficit (figure 1). There  
 119 was a significant difference ( $P < 0.05$ ) in plant height among the watering regimes in W6 and W7 but  
 120 not in W8 and the first and third regimes has the tallest plants (Table 1). These may be since the plant  
 121 has developed tolerant mechanism and the reason why the growth in height continued. The reduction  
 122 in plant height in relation to increased water deficit was more pronounced in BG doguwa and NERICA  
 123 (figure 1).



124  
 125 **Figure1:** Effect of different watering regimes on the plant height of BG doguwa,  
 126 Mai zazzabi and NERICA rice land races at week 1 to week 8. Values are means of two Replications  
 127  $\pm$  Std error.

128  
 129  
 130

131 **Table 1: General linear model (P-Values) for the interaction between all the parameters versus**  
 132 **Treatment, Varieties**

Parameter	Treatment	Varieties	Treatment *Varieties
Plantheight(cm) Week 5	0.001	0.045	0.107
Plantheight(cm) Week 6	0.007	0.002	0.015
Plantheight(cm) Week 7	0.001	0.002	0.012
Plantheight(cm) Week 8	0.001	0.014	0.060
NL Week 5	0.001	0.001	0.191
NL Week 6	0.001	0.001	0.091
NL Week 7	0.001	0.001	0.054
NL Week 8	0.001	0.001	0.047
Stem diameter(cm)	0.002	0.014	0.181
Root length(cm)	0.027	0.232	0.349
Biomass weight(g)	0.001	0.001	0.043


133 \*Note. NL=Number of leaves, coloured fonts indicate significant P values < 0.05

134 **Table 2: Mean Values ± Standard error of interaction between Parameters versus Treatment**

Parameter	Treatments			P-value
	Control	Mild stress	Severe stress	
Stem diameter(cm)	1.60 <sup>a</sup> ±0.19	1.46 <sup>b</sup> ±0.19	1.39 <sup>b</sup> ±0.13	0.006
Root length(cm)	14.33 <sup>a</sup> ±1.85	13.08 <sup>ab</sup> ±1.99	12.34 <sup>b</sup> ±2.11	0.029
Biomass(g)	2.60 <sup>a</sup> ±1.30	1.43 <sup>b</sup> ±0.46	1.47 <sup>b</sup> ±0.61	0.001


135 Key: Means that do not share a letter are significantly different, coloured fonts indicate significant P values < 0.05

136 Number of leaves decreased under water deficit (figure 2). Control treatment recorded higher number  
 137 of leaves than plants in treatment 2 (mildly stressed) and treatment 3, (severe stress) respectively, the  
 138 most pronounced reduction occurred with BG doguwa land race.  
 139




**Figure 2:** Effect of different watering regimes on the number of leaves(NL) of BG doguwa, Mai zazzabi and Nerica rice land races at week 1 to week 8. Values are means of two Replications  $\pm$  Std error.

Root length noticeably reduced due to increased water deficit (figure 3). Plants of the Control treatment recorded higher root lengths than plants of treatment 2 (mildly stressed) and treatment 3, (severe stress) respectively. The more pronounced reduction occurred with BG doguwa. Stem diameter (figure 3) and as well biomass (g) (figure 4) was reduced due to the severity in water deficit. Plants watered daily (Control) had higher stem diameters(cm) (figure 3) and biomass(g) (figure 4) accumulation than plants of the other watering regimes.



**Figure 3:** Effect of different watering regimes on the root length(cm) and stem diameter(cm) of BG doguwa, Mai zazzabi and Nerica rice land races. Values are means of two Replications  $\pm$  Std error.



**Figure 4:** Effect of different watering regimes on biomass weight(g) of BG doguwa, Mai zazzabi and Nerica rice land races. Values are means of two Replications  $\pm$  Std error.

There was a significant difference ( $P < 0.05$ ) among the varieties in stem diameter (cm) and whole plant dry weight(g) as shown in (Table 3). BG doguwa had the highest stem diameter (cm) and biomass (g) in the first watering regime (control) followed by Mai-zazzabi, but Mai-zazzabi had the highest in the second (mild stress) and third (severe stress) watering regimes followed by BG doguwa.

**Table 3: Mean Values  $\pm$  Standard error of interaction between Parameters versus Varieties**

Parameter	Varieties			P-value
	BG doguwa	Mai-zazzabi	Nerica	
Stem diameter(cm)	1.55 <sup>a</sup> $\pm$ 0.21	1.52 <sup>ab</sup> $\pm$ 0.11	1.39 <sup>b</sup> $\pm$ 0.20	0.037
Root length(cm)	13.49 <sup>a</sup> $\pm$ 2.81	13.72 <sup>a</sup> $\pm$ 1.56	12.55 <sup>a</sup> $\pm$ 1.70	0.279
Biomass (g)	2.17 <sup>a</sup> $\pm$ 1.14	2.10 <sup>a</sup> $\pm$ 1.07	1.23 <sup>b</sup> $\pm$ 0.42	0.015

Key: Means that do not share a letter are significantly different, coloured fonts indicate significant P values  $< 0.05$

#### 4.0 DISCUSSION

The trend in the reduction in plant height with increase in water deficit (Figure-1) in rice agrees with results of Siddique *et al.* (2000) in wheat. Growth involves both cellular growth and development which is a process consisting of cellular division, cellular enlargement and differentiation and these processes are very sensitive to water deficit because they all depend on cellular turgidity (Jones and Lazenby, 1988). The inhibition of cell expansion is usually followed closely by a reduction in cell wall synthesis (Salisbury and Ross, 1992). This may have affected plant height of the rice. This study revealed Mai zazzabi to be generally taller than BG doguwa and Nerica at severe moisture stress conditions. This implies that Mai zazzabi can withstand higher levels of dehydration. In terms of plant height, Mai zazzabi is the most tolerant variety among the three varieties. The number of leaves decrease with increase in water deficit (figure 2). Water deficit might, inhibit photosynthesis and produce less assimilates which resulted in lower number of leaves this result agrees with the work of Hossain (2001). The plant shoot dry weights decreased in increase in water deficit. Similar results were obtained by Willumsen (1993). The reduction in shoot dry weight could be associated with reduced rate of leaf production hence low number of leaves. Reduction in leaf growth may also have

been contributed by lower rates of cell division and cell extension in the leaves. Reduction in leaf growth leads to less photosynthesis hence retarded overall plant growth as the resources required for growth processes become limited in supply (Mwai, 2002). Plants show increased root: shoot ratio during soil moisture deficit (Boyer, 1985). Similar results have also been obtained in mango rootstock seedlings (Luvaha, 2005). The differential sensitivity of roots and shoots (with root growth being less sensitive to water deficits) leads to large increases in the root to shoot ratio in drought conditions (Sharp and Davies, 1985). This may be an adaptation of Mai zazzabi rice varieties for survival under water scarcity conditions since increased root surface area allows more water to be absorbed from the soil. Shoot growth decline coupled with continued root growth would result in an improved plant water status under extreme water deficit conditions. In maize seedlings, root growth is not affected by low water potentials which are in another way completely inhibitory to shoot growth (Boyer, 1985). The three varieties may possess mechanisms of biomass accumulation under water stress conditions. In this study Mai zazzabi exhibits superior adaptation to water deficit in terms of biomass accumulation. Whole plant dry weight significantly declined with moisture deficit. This finding agrees with the results reported by Emmam et al. (2010). Water deficit may have played a role in influencing the increment in height and leaf area per plant which eventually influenced the increase in the shoot dry matter of plants. A reduction of photosynthetic surface by water deficit decreases the ability of plant to produce dry matter.

## 5.0 CONCLUSION

The present study has revealed the significance of water in rice physiological growth where water deficit has led to a reduction in plant growth and biomass accumulation. Relating to plant growth Mai zazzabi is the most tolerant among the three varieties and can accumulate higher biomass under water stress conditions.

Mai zazzabi can be recommended to farmers as the variety that is tolerant to moisture stresses especially those in the northern part of the country where there is low annual rainfall. And recommend that more research be conducted on rice to come up with improve variety that will be more tolerant and adapted to other environmental conditions not necessarily moisture to increase productivity and yield to meet up with the rapid increase in population growth and demand worldwide.

## REFERENCE

- Anjum, F., Yaseen, M., Rasul, E., Wahid, A., and Anjum, S. Water stress in barley (*Hordeum vulgare*): Effects on morphological characters. *Pakistani Journal of Agricultural Science*, 2003; 40- 43.
- Botwright Acuna, T.L., Lafitte, H.R., and Wade, L.J. Genotype and environment interactions for grain yield of upland rice backcross lines in diverse hydrological environments. *Field Crops Research*, 2008; 108, 2, 117-125. doi:10.1016/j.fcr.04.003
- Boyer J. S. Water transports. *Ann. Rev. Plant Physiology*. 1985; 36: 473-516.
- Emmam, Y., Shekoofa, A., Salehi, F., and Jalali, A.H., Water stress effects on two common bean cultivars with contrasting growth habitats. *American- Eurasian Journal of Agriculture and Environmental Science*, 2010; 9(5): 495-499.
- Fairhurt, T.H., and Dobermann, A. Rice in the global food supply. In: better crops for international Special supplementary, 2002; 37. [http://www.ipni.net/ppiweb/bcropint.nsf/\\$webindex/0E477FFC43BD62DA85256BDC00722F62/\\$file/BCI-RICEp03.pdf](http://www.ipni.net/ppiweb/bcropint.nsf/$webindex/0E477FFC43BD62DA85256BDC00722F62/$file/BCI-RICEp03.pdf)
- FAOSTAT. (December, 2013) Food security: Global food. 2, URL <http://Food-Security-Statistics@FAO.org>.
- Gleick, P.H., Wolff, E.L & Chalecki, R.R. The new economy of water: The risks and benefits of globalization and privatization of fresh water: Pacific Institute for studies in Development, environment and Security. Oakland CA., 2002; 48pp. ISBN No. 1-893790-07-X.

230 Hafeez, M.M., Bouman, B.A.M., Van de Giesen, N. and Vlek, P. Scale effects on water use and water  
 231 productivity in a rice-based irrigation system (UPRIIS) in the Philippines. *Agricultural Water*  
 232 *Management*. 2007; 92, 81–89.

233 Hossain, M.A. Growth and yield performance of some boro rice cultivars under different soil moisture  
 234 regimes. M.S. Thesis, Department Crop Botany. Bangladesh Agricultural University,  
 235 Mymensingh, 2001; Chapter 2-5.

236 IFPRI. What is the irrigation potential for Africa? A combined biophysical and socioeconomic  
 237 approach. IFPRI, Discussion Paper 00993. International Food Policy Research Institute, Washington,  
 238 DC, 2010.

239 IPCC. Climate Change 2007: Impacts, adaptation and vulnerability. Contribution of Working Group II  
 240 to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (Parry, M.L.,  
 241 Canziani, O.F., Palutikof, J.P., van der Linden, P.J. and Hanson, C.E. eds). Cambridge University  
 242 Press, Cambridge.

243 Jones B.M. and Lazenby A. The grass crop, the physiological basis of production. Chapman and Hall,  
 244 London. 1988; pp. 226-240.

245 Kamoshita, A., Rodriguez, R., Yamauchi, A., and Wade, L. Genotypic variation in response of  
 246 rainfed lowland to prolonged drought and re-watering. *Plant Production Science*, 2004; 7, 4, 406-420.

247 Khalil, H.I., Bari, A. Khan, S. and Zada, I. Genetic variation for yield components in rice. *Agricultural*  
 248 *and Biological Sciences*, 2009; 4(6): 60-64.

249 Kramer, P.J. and Boyer, J.S. Water relations of plants. Academic Press New York, USA, 1995; Pp  
 250 495-496.

251 Liu, J.X., Liao, D.Q., Oane, R., Estenor, L., Yang, X.E., Li, Z.C., and Bennett, J. Genetic variation in  
 252 the sensitivity of anther dehiscence to drought stress in rice. *Field Crops Research*, 2006; 97, 87-100.  
 253 doi:10.1016/j.fcr.2005.08.019

254 Luvaha E. The effects of water deficit on the growth of mango (*Mangifera indica*) rootstock seedlings.  
 255 M.Sc Thesis. Maseno University, Kenya, 2005.

256 Mwai.G.N. Growth response of spider plant (*Cleome gynandra* L.) to salinity. M.Sc. Thesis. Maseno  
 257 University, Kenya, 2001.

258 Salisbury B. and Ross W. Plant physiology. 4th edition, Wadsworth, Belmont, California. 1992; pp.  
 259 580-585.

260 Sharp R.E. and Davies W.J. Root growth and water uptake by maize plants in drying soil.  
 261 *Experimental Botany*. 1985; 36: 1441- 1456.

262 Siddique, M.R., Hamid, A., and Islam, M. Drought stress effects on water relations of wheat. *Plant*  
 263 *Physiology*, 2000; 41: 35-39.

264 Willumsen J. Assessment of fluctuations in water and air content of pot substrates during plant  
 265 growth. *Acta Horticulture*. 1993; 295: 249-259.

**Comment [A1]:** Is it the name of journal or the name of author?

**Comment [A2]:** It is the name of journal.