

Effects of Lead on Different Seedling Growth Attributes of Cow Pea (*Vigna unguiculata*)

ABSTRACT

Aims: This study was conducted to determine the toxicity and tolerance to lead by seedling growth of an important country legume crop cowpea (*Vigna unguiculata*).

Study Design: The seedlings of cowpea were grown in sand culture at 0, 20, 40, 60, 80 and 100 ppm of metal ions of lead salt as lead acetate.

Place and Duration of Study: The experimental site is located in the Department of Botany at the Karachi University Campus, Pakistan in 2011.

Methodology: The healthy seeds of *Vigna unguiculata* were surface sterilized with 0.2% solution of sodium hypochlorite (NaOCL) for one minute to avoid any fungal contamination. The sand was collected from the construction site of the Karachi University washed 2-3 times with tap water, distilled water and also with 5% HCl to remove any types of impurities from the soil. Seedlings were grown in sand culture at 0, 20, 40, 60, 80 and 100 ppm of metal ions of lead salt as lead acetate. The Hoagland solution was used for the supply of nutrient elements. The experiment was conducted for six weeks. Five ml of lead concentration were poured weekly and before given concentration of lead the materials of the tray were drained out to avoid any algal contaminations. The experiment was completely randomized and consists of six treatments replicated six times. After six weeks the seedlings were harvested and morphological parameters shoot, root, seedling length (cm), number of leaves and leaf area (sq. cm) was noted. The biomass production such as shoot, root, leaf and total seedling dry weight (g) was also observed along with root / shoot, leaf weight, leaf area ratio and specific leaf area. The seedlings of cow pea were dried in an oven at 80° C for 24 hours until the seedlings were completely oven dried. Leaf area, Root / shoot ratio, leaf weight ratio, specific leaf area, leaf area ratio was determined and a tolerance index was determined. The data obtained was statistically analyzed.

Results: The effects of different concentrations (0, 20, 40, 60, 80, 100 ppm) of lead on seedling growth performance of cow pea (*Vigna unguiculata*) as compared to control were observed. Lead treatment in the form of lead acetate at 100 highly affected seedling growth and biomass production of *V. unguiculata* as compared to control. Lead treatment at 40 ppm produce significant ($P = .05$). reduction in seed germination of *V. unguiculata* as compared to control. Lead treatment at 20 ppm concentration produced significant reduction in shoot length as compared to control. Root growth is an important growth variable and found negatively affected by different concentration of lead treatment. The results also showed that lead treatment in the substrate at same concentration (20 ppm) produced significant effect on root and seedling growth of *V. unguiculata*. The treatment of lead at 20 ppm produced significant ($p < 0.05$) on seedling dry weight of *V. unguiculata* as compared to control. However, increase in concentration of lead treatment at 80 ppm was found sufficient to cause significant reductions in seedling dry weight of *V. unguiculata* as compared with control. The seedlings of *V. unguiculata* were also tested for percentage of tolerance to lead. The results showed that *V. unguiculata* has high tolerance to lead at 20 ppm and lowest at 80 ppm of lead. *V. unguiculata* seedlings showed highest percentage of tolerance (92.50 %) to lead at 20 ppm and the lowest (64.50 %) at 80 of lead treatment. *V. unguiculata* seedlings showed better percentage of tolerance 73.25 % to lead at 60 ppm.

Conclusion: In conclusion, lead was toxic causing significant reduction ($P = .05$). to the seedling and biomass production of cowpea. Increased in lead concentration decreased the tolerance indices of cow pea seedlings growth.

Key words: Crop, germination, growth, lead, tolerance, toxicity.

1. INTRODUCTION

Plants are an integral part of life in many indigenous communities and modified the plant cover over wide areas [1-2]. Lead is considered highly toxic to the growth of plants. Lead (Pb) is an environmental pollutant extremely toxic to plants and other living organisms including humans [3]. A high concentrations of Pb eventually may cause cell death [4]. Soil pollution with lead is a problem of concern [5].

The concentrations of heavy metals in the environment have increased at alarmingly level. Metals are toxic to plants and fungi [6]. There are few detail reports on the impacts of metals on seed metabolism, germination, particularly roots and shoots recorded [7]. Lead is a global environmental pollutant that is present in soil, water, air and biota. The increase in concentration of heavy metal decreased plant growth and responsible to death [8]. Lead is naturally occur in substantial quantities

in the earth's surface food, water urban soil and air and lead stress causes multiple direct and indirect effects on plant growth, metabolism and also alters some physiological processes [9-10]. The metals absorbed by plants and prove toxic to plants that can be observed as growth retardation, alteration in biochemical processes [11-13].

In Pakistan substantial quantities of agricultural chemical are used annually to enhance yield [14]. The ever increase of lead concentration over the wide areas of Karachi and rural areas raising a serious questions as to its effects plant growth. Although the data on the effects of lead on cowpea seems scanty. The response of plant growth to toxic effects of heavy metals has become the subject of great interest in the field of ecology. Attention has been given in developed countries, about the effects of metal toxicities on crop growth. Therefore, this study was carried out with the aim to determine the effect of lead on seedling growth of an important country legume crop cowpea (*Vigna unguiculata*).

2. MATERIALS AND METHODS

The healthy seeds of *Vigna unguiculata* were obtained from the market and were surface sterilized with 0.2% solution of sodium hypochlorite (NaOCl) for one minute to avoid any fungal contamination. The experimental site is located in the Department of Botany at the Karachi University Campus. The sand was collected from the construction site of the Karachi University. The sand was sieved through 2.0 mm sieve and after that it was washed 2-3 times with tap water and later with distilled water. The sand was also washed with 5% HCl to remove any types of impurities. The pots with 7.3 cm in diameter and 9.6 cm in depth were filled with sand upto 2/3. All the pots were then placed in trays. Seedlings were grown in sand culture at 20, 40, 60, 80 and 100 ppm of metal ions of lead salt as lead acetate. In control, no treatment was given except distilled water. As nutrients elements were absent in sand therefore, Hoagland solution was used for the supply of nutrients. The Hoagland solution was applied for 3-4 days. Three uniform size seedlings of cowpea were transplanted in each pot. Initially, 5 ml solution of lead acetate $Pb(CH_3COO)_2$ were applied. Every week the appearance of seedlings growth was recorded. The irrigation was carried out with the tap water on daily basis. The experiment was conducted for six weeks. The nutrient solution (The Hoagland solution) was given after three days time interval for nutrition and proper growth of plant. Five ml solution of lead concentration were poured weekly and before given concentration of lead the materials of the tray were drained out to avoid any algal contaminations. The experiment was completely randomized and replicated six times.

After six weeks, the seedlings were harvested. The shoot, root, seedling length (cm), number of leaves and leaf area, specific leaf was recorded. The biomass production such as shoot, root, leaf and total seedling dry weight (g) was also determined alongwith root / shoot, leaf weight, leaf area ratio and specific leaf area. The seedlings of cowpea were dried in an oven at 80° C for 24 hours until the seedlings were completely oven dried.

Root / shoot ratio, leaf weight ratio, specific leaf area, leaf area ratio were determined by the following formulae, respectively.

Root/ shoot ratio = root dry weight / shoot dry weight

Leaf weight ratio = leaf dry weight / total plant dry weight

Specific leaf area = Leaf area / leaf dry weight

Leaf area ratio = Leaf area / Total plant dry weight

A tolerance index was determined by the following formulae as described by Iqbal and Rahmati [15]:
Mean root length in metal solution/Mean root length in distilled water X 100

Analysis of variance and Duncan's Multiple Range Test using personal computer software packages SPSS version 14.0 statistically analyzed the data obtained.

3. RESULTS

Lead treatment was found toxic to the all seedling growth parameter of *V. unguiculata*. The seedling growth performance of *V. unguiculata* were tested in different concentrations (0, 20, 40, 60, 80 and 100 ppm) of lead as compared to control (Table 1-3, Fig.1). Lead treatment was found highly toxic to all seedling growth parameter of *V. unguiculata* at 100 ppm concentration. Lead treatment at 40 ppm produced significant ($p<0.05$) effects on shoot length of *V. unguiculata* as compared to control (Table 1). Lead treatment at 20 ppm affected root growth of *V. unguiculata*. Lead treatment at 20

ppm did not produced significant effects on seedling length, while increase in lead concentration at 40 ppm significantly decreased seedling length of *V. unguiculata* as compared to control. Lead treatment at 40 ppm concentration produced toxic effects on number of leaves of *V. unguiculata* and highly affected at 100 ppm lead. The results also showed that lead treatment at similar concentration 40 ppm produced significant effect on leaf growth of *V. unguiculata*.

Table 1. Effects of lead on seedling growth of cow pea (*V. unguiculata*)

Treatments (ppm)	Shoot length (cm)	Root length (cm)	Seedling length (cm)	No of leaves	Leaf area (sq. cm)
0	48.03±0.28d	22.39±0.28e	70.33 ±0.52d	12.50 ±0.76c	28.43±0.281d
20	47.05±0.26d	21.15±0.13d	68.20 ±0.37d	11.00 ±0.77c	27.40±0.35bd
40	42.73±0.74c	19.50±0.25c	62.23 ±0.99c	9.33 ±0.95ab	25.10±0.58c
60	39.38±1.44b	17.81±0.14b	57.20±1.82b	8.66 ±0.80ab	20.31 ±1.35b
80	32.96±1.38a	16.78±0.53ab	49.75±1.89a	7.83 ±0.87a	17.63 ±0.40a
100	30.71±1.39a	16.60±0.45a	47.31±1.75a	7.66 ±0.42a	16.66 ±0.18a

Lead treatment at all concentration decreased high percentage of seedling dry weight of *V. unguiculata* (Table 2). The treatment of lead at 40 ppm produced significant ($p<0.05$) on shoot dry weight of *V. unguiculata* as compared to control. However, increase in concentration of lead treatment at 80 -100 ppm further decreased shoot dry weight of *V. unguiculata* as compared with control. Lead treatment at 20 ppm concentration significantly affected root dry weight cowpea. Lead treatment at 20 and 40 ppm concentration was less toxic for decrease in leaf dry weight of cowpea. Lead treatment at 40 ppm concentration showed more decrease in leaf dry weight of cowpea. Lead concentration of 20 ppm showed high percentage of decrease in total plant dry weight of *V. unguiculata*.

Table 2. Effects of lead on seedling dry weight of cow pea (*V. unguiculata*)

Treatments (ppm)	Shoot dry weight (g)	Root dry weight (g)	Leaf dry weight (g)	Total plant dry weight (g)
0	1.74±0.04b	0.86±0.02d	0.95±0.01c	3.55±0.06e
20	1.56±0.19b	0.84±0.03c	0.83±0.041c	3.24±0.21d
40	1.23±0.18a	0.74±0.02ab	0.77±0.04c	2.70±0.18c
60	1.03±0.07a	0.69±0.04ab	0.67±0.03ab	2.43±0.12bc
80	0.92±0.05a	0.64±0.03a	0.63±0.02a	2.20±0.08ab
100	0.86±0.03a	0.50±0.03a	0.48±0.04a	1.84±0.09a

Lead concentration of 20 ppm showed high percentage of decrease in root / shoot ratio, leaf weight, specific leaf area and leaf area ratio of *V. unguiculata* (Table 3).

Treatments (ppm)	Root / shoot ratio	Leaf weight ratio	Specific leaf area	Leaf area ratio
0	0.49±0.01b	0.26±0.01b	27.16± 0.14a	7.27±0.09a
20	0.58±0.09a	0.26±0.02a	29.18±1.08b	7.64±0.64b
40	0.66±0.08a	0.29±0.03a	31.19±1.48bc	8.87±0.62d
60	0.67±0.01a	0.29±0.01a	31.14±1.38b	9.02±0.43c
80	0.70±0.02a	0.28±0.01a	30.62±1.43b	8.78±0.36bc
100	0.58±0.02a	0.26±0.01a	36.61±2.79c	9.30±0.37bc

The seedlings of *V. unguiculata* were also tested for percentage of tolerance to lead. The results showed that *V. unguiculata* has greater tolerance to lead at 20 ppm and lowest at 100 ppm of lead (Fig. 1). *V. unguiculata* showed higher tolerance (94.46 %) to lead at 20 ppm and lowest (74.10 %) at 100 ppm of lead. *V. unguiculata* seedlings showed better percentage of tolerance (87.09 %) at 60 ppm and (74.99 %) at 80 ppm lead treatment (Fig. 1).




Fig. 1. Percentage tolerance index in seedlings of cow pea (*Vigna unguiculata*) to different concentration of lead.

4. DISCUSSION

The plants under stress condition are likely may be affected by high concentrations of pollutants. The growth in plants or any other living organisms affected directly or indirectly in the presence of heavy metals [16]. Lead is a toxic heavy metal. In present study, the effect different concentrations of lead on root, shoot, seedling length and seedling dry weight of an important bean crop cowpea (*Vigna unguiculata*) were recorded. Kasim *et al.*, [17] also showed that lead treatment at 150 mM produced significant physiological, photosynthetic and ultra structural changes in seedlings of *Vigna unguiculata*. The seedling growth parameters of cowpea responded differently to lead treatment at higher concentration as compared to control. High percentage of decrease in shoot growth of cowpea at 40-60 ppm lead treatment provided evidence that the treatment of lead in excess was inhibitory to plant growth and development. In present study, a significant decrease in number of leaves and leaf area of cowpea at 100 ppm lead treatment was also observed. Excessive amount of toxic element usually caused reduction in plant growth [18]. The permeability of metals can decreased the growth of plants. The reduction in the seedling growth of *V. unguiculata* with the increase in concentration of lead in the immediate environment provides further evidence that the lead in excess may be inhibitory to plant growth and development. Toxic metal ions enter cells by means of the same uptake processes as essential micronutrient metal ions. The amounts of metal absorbed by a plant depend on the concentrations and speciation of the metal in the soil solution. Excessive concentrations of metals result in phytotoxicity [19]. Similarly, drastic decrease in seedling growth of *Vigna radiata* (L.) Wilczek cv. Pusa Baisakhi in the presence of 1.0 mM lead acetate [20]. The root elongation tests have been used to evaluate the damage caused by toxic compounds [21] and many

species including cabbage, lettuce, carrot, cucumber, tomato and oats have been recommended for the phytotoxicity test [22]. The roots are normally has the ability to supply water and nutrients to the plants. They are also required to produce hormones, which may regulate the growth and performance of both root and shoot [23]. In present study, the root growth of *V. unguiculata* was found highly decreased at 80 ppm lead concentration. These findings are also agree with the work of [24] and reported that a low concentration of 1 μ M lead (Pb) reduced root, shoot growth of cowpea (*Vigna unguiculata*). The primary site of Pb²⁺ toxicity was the root, caused severe reductions in root growth, loss of apical dominance, the formation of localized swellings behind the root tips.

The significant decrease in seedling dry weight of *V. unguiculata* due to metal toxicity of lead was also recorded. The treatment of lead in *V. unguiculata* provided evidence that the trace element in nutrient medium if present in excess may be inhibitory to plant growth and development especially at more than 60 ppm. Toxicants accumulate in the plant when soluble forms are present in high quantities. The exact amount of accumulation depends upon the solubility of the pollutants in the soil [25]. The biomass production of *V. unguiculata* was initially non significant and decreased and with increasing the lead concentrations upto 100 ppm significantly.

5. CONCLUSION

According to tolerance test, it could be seen that tolerance to lead was higher at low concentration of lead in the seedlings of *V. unguiculata*. These results showed that the reason of tolerance against heavy metals might be a physiological association of the tolerance mechanism to these metals. The seedling growth of *V. unguiculata* showed high percentage of tolerance to lead at 20 ppm concentration. The treatment of 100 ppm concentration of lead produced lowest percentage of tolerance in seedling of cowpea.

It is concluded, that concentration of heavy metals like lead and other metals in the environment has been increased due to automobiles, industries, agro chemicals and anthropogenic activities and responsible for limiting the crop yield. Lead is found highly toxic pollutant for plants. The results showed that treatment of different concentrations of lead to the seedlings of *V. unguiculata* responded differently with the increasing contamination exposure as 20, 40, 60, 80 and 100 ppm. The response of *V. unguiculata* seedlings in the form of tolerance indices to lead treatment was found suitable pollutant indicator to study the deleterious effects of the lead. The present findings proved the deleterious effects of lead at higher concentration to the seedlings of *V. unguiculata*. The information from the present studies would be helpful in understanding the level of lead tolerance in seedlings of *V. unguiculata* while growing in lead polluted areas. Heavy reliance on metals containing agrochemicals such as fungicides, nematicide, and pesticides (Lead arsenate) can be discouraged. The continuous release of lead into the immediate environment may endanger the growth performance of other crops. Current research shows that lead treatment at different concentration has produced an important effect on seed germination and seedling growth of *V. unguiculata*.

REFERENCES

1. Bhatia H, Sharma YP, Manhas RK, Kumar K.. Ethnomedicinal plants used by the villagers of district Udhampur, J&K, India. *Journal of Ethnopharmacology*, 2014, 151 (2): 1005-1018.
2. Eyre SR. The outlook for wild nature. In "Vegetation and Soils". Edward Arnold, The Pitman Press, 2 Ed. U.K. p. 273-277. 1984.
3. Lamhamdi M, Bakrim A, Aarab A, Lafont R, Sayah F. Lead phytotoxicity on *V. unguiculata* (*Triticum aestivum* L.) seed germination and seedlings growth. *Comptes Rendus Biologies*, 2011, 334(2): 118-126.
4. WHO. Effects of heavy metals in plants at the cellular levels. In: Schuurmann G (ed). *Ecotoxicology, ecological Fundamentals, Chemical Exposures and Biological effects*. Pp 587-620. Heidelberg, Wiley. 2001.
5. Pirzda H, Ahmad SS, Rashid A, Shah T. multivariate analysis of selected roadside plants (*Dilbergia sisso* and *Cannabis sativa*) for lead pollution monitoring. *Pak. J. Bot.* 2009, 41(4): 1729-1736.]
6. Crane, S., Barkay, T. and Dighton, J. 2012. The effect of mercury on the establishment of *Pinus rigida* seedlings and the development of their ectomycorrhizal communities. *Fungal Ecology*, 5(2): 245-251.
7. Kranner, I. and Colville, L. 2011. Metals and seeds: Biochemical and molecular implications and their significance for seed germination. *Experimental and Experimental Botany*, 72(1): 93-105.
8. Kumar, P., Sudha, S., Ranjitha, T. and Kumari, B.D. 2011. Effect of chelators and mercury on growth and development of *Catharanthus roseus* (L). G Don. *Journal of Agricultural Technology*, 7(2): 281-288.

9. Diaz-Aguilar, I., Larque-Saavedra, M.U., Alcantar_Gonzalez, G. and Carrillo-Gonzalez, R. 2001. Alternation of some physiological processes in *V. unguiculata* by lead additions. *Revista International-de-Contamination Ambiental*, 17(2): 79-80.
10. Haq, R., Khan, F.F.U. and Haq, E. 2013. Adverse effect of lead acetate on light with protein of *Bactrocera cucurbitae*. *Journal of Basic and Applied Sciences*, 9: 29-296.
11. Arun KS, Cervantes C., Loza-Tavera H, Avudainayagam S. chromium toxicity in plants. *Environ. Int.* 2005, 31:739-753.
12. Shafiq M, Iqbal MZ, Athar M. Effect of lead and cadmium on germination and seedling growth of *Leucaena leucocephala*. *Journal of Applied Science and Environmental Management*, 12 (3), 2008, 61-66.
13. Shafiq M, Iqbal MZ. "Impact of Automobile Pollutants on Plants". ISBN 978-3-8443-8504-5. LAP LAMBERT Academic Publishing GmbH & Co. KG Heinrich-Böcking-Str. 6-8, 66121, Saarbrücken, Germany. 2012, 132 pp.
14. Nuzhat, A., Jameela, A., Munawar, R. and Attaurrehman. 2005. Hydrolysis of a fungicides, Buprimate by indigenous *Achromobacter* sp. *International Journal of Biotechnology*, 2(2): 357-363.
15. Iqbal, M.Z. and Rahmati, K. 1992: Tolerance of *Albizia lebbeck* to Cu and Fe application. *Ekologia* (CSFR) 11 (4): 427-430.
16. Szollosi, R. 2014. Chapter 3 – Superoxide Dismutase (SOD) and Abiotic Stress Tolerance in Plants: An Overview. *Oxidative damage to plants. Antioxidant Networks and signaling*, pp- 89-129. Academic Press.
17. Kasim, W.A., Abokassem, E.M., Rajab, G.A. and Sewelam, R.N. 2014. Allevation of lead stress toxicity in *Vigna unguiculata* by salicylic acid. *Egyptian Journal Exp. Biology (Bot.)* 10(1): 37-49.
18. Kubota, J. and Allaway, W.H. 1972. In "Micro nutrients in Agriculture". Mortvedt. J.J., Giordano, P.M., Lindsay, W.L. (ed.). Soil Science Society of America, Madison. Wis. p. 525-554.
19. Patra, M., Bhowmik, N., Bandopadhyay, B. and Sharma, A. 2004. Comparison of mercury, lead and arsenic with respect to genotoxic effects on plant systems and the development of genetic tolerance. *Experimental and Experimental Botany*, 52(3): 199-223.
20. Singh, R.P., Tripathi, R.D., Dabas, S., Rizvi, S.M.H., Ali, M.B., Sinha, S.K, Gupta, D.K., Mishra, S. and Rai, U.N. 2003. Effect of lead on growth and nitrate assimilation of *Vigna radiata* (L.) Wilczek seedlings in a salt affected environment. *Chemosphere*, 52(7): 1245-1250.
21. U.S.E.P.A. (U.S. Environmental Protection Agency), 1982. Seed Germination/Root Elongation Toxicity Test EG-12.
22. F.D.A. (Food and Drug Administration), 1982. Environmental Assessment of Technical Assistance Document 4.06. The Center of Veterinary Medicine, U.S. Department of Health and Human Services, Washington, D.C.
23. Blackman, P.G. and Davies, W.J. 1985. Root to shoot connection in maize plants and the effects of soil drying. *J. Exp. Bot.* 36: 39-48.
24. Kopittke, P.M. Asher, C.J., Kopittke, R.A. and Menzies, N.W. 2007. Toxic effects of Pb^{2+} on growth of cowpea (*Vigna unguiculata*). *Environmental Pollution*, 150(2):280-287. .
25. Treshow, M. 2010. *Terrestrial Plants and Plant Communities*. Chapter 10. *Department of "Biology*, University of Utah, Salt Lake City, Utah 84112, U.S.A. pp 225-236.