

# 1 **Phytochemical screening and evaluation of the cytotoxicity** 2 **of fruits of *Solanum torvum* Swartz (Solanaceae) on HFF** 3 **cells (Human Foreskin Fibroblasts).**

## 4 **Abstract:**

5 **Objective:** The aim of this work is to evaluate the cytotoxic activity of the 70% ethanolic  
6 extract of the fruits of *Solanum torvum* Swartz (Solanaceae) on HFF (Human Foreskin  
7 Fibroblast) cells in *in vitro* culture and to determine its phytochemical composition.

8 **Study plan:** An ethnobotanical survey was conducted in August 2015 in the Haut-Sassandra  
9 Region (Ivory Coast), on medicinal plants with multiple uses, and at the end of this survey  
10 the fruits of *Solanum torvum* have been selected then harvested in the Sub-prefecture of  
11 Bédiala (Ivory Coast). Then after drying, the ethanolic extract of the fruits was prepared and  
12 sent to France in February 2016 at the Laboratory Adaptation and Pathogenesis of  
13 Microorganisms (LAPM) of Grenoble for cytotoxic tests. Phytochemical screening was  
14 carried out at the Faculty of Biological Sciences of Félix Houphouët Boigny University (Côte  
15 d'Ivoire).

16 **Methods:** From an ethnobotanical survey, the fruits of *Solanum torvum* were harvested. The  
17 70% ethanolic extract prepared from the fruits of this plant was tested *in vitro* on divisional  
18 HFF cells after phytochemical screening.

19 **Results:** The result revealed that this extract has cytotoxic activity on the tested HFF cells. At  
20 800 µg/mL, the survival rate of HFF cells increased from 100% to 4% of living cells.  
21 Phytochemical screening revealed the presence of compounds such as alkaloids, tannins,  
22 polyphenols, saponins and flavonoids.

23 **Conclusion:** This extract is cytotoxic on HFF cells. It is therefore necessary to continue  
24 studies on toxicity and to be cautious in the use of *solanum torvum* fruits in traditional  
25 medicine.

26 **Key words:** HFF cell, Cytotoxic, Extracts, ethnobotanical survey, *Solanum*  
27 *torvum*, Phytochemicals

## 28 **1. Introduction**

29 *Solanum torvum* belongs to the family Solanaceae. *Solanum torvum* is native to Central and  
30 South America, from Mexico to Brazil and Peru. It has spread widely in the Caribbean [1]. In  
31 West and Central Africa, it is grown locally in gardens for cooking [2]. Fruits are widely used

32 in the treatment of shingles in Cameroon[3]. They are also used as a vegetable and considered  
33 an essential ingredient in the diet of the South Indian population[1]. A fruit decoction is used  
34 in Ghana for the treatment of cough, liver disease and spleen [4]. *Solanum torvum* are rich in  
35 antioxidant, Its extracts are used in the preparation of tonic and hemopoietic agents and also  
36 for the treatment of pain throughout the body [5,6]. Previous work has revealed that the fruits  
37 of *Solanum torvum* are used in the treatment of various conditions: microbial infections [7],  
38 hypertention [8], kidney disease [9] diabetes [10]. Pérez-Amador *et al.* [11] reported the  
39 presence of glycoalkaloids in the fruit of *Solanum torvum* and another study showed that very  
40 low doses of glycoalkaloids in some Solanaceae are toxic [12,13]. In view of the interest of  
41 *Solanum torvum* fruits in the traditional medicine, we have undertaken to evaluate  
42 scientifically the cytotoxicity and phytochemical screening of the fruits. An ethnobotanical  
43 survey was conducted in August, 2015 in the Haut-Sassandra Region (Ivory Coast). This  
44 survey showed that all parts of *Solanum torvum* are intensively used, especially leaves and  
45 fruits in the treatment of dermatosis and anemia. The aim of this work is to perform a  
46 phytochemical screening and a cytotoxicity study of *Solanum torvum* fruits on Human  
47 Foreskin Fibroblasts (HFF) cells.

## 48 2. Material and methods

### 49 2.1 Plant material

50 The fruits of *Solanum torvum* (Fig 1) were harvested in Biadiala in the Department of Daloa  
51 (Ivory Coast).



60 Fig 1: Leafy and fruiting twig of *Solanum torvum* (Solanaceae)

### 61 **2.3 Cellular material**

62 The cellular support consists of human HFF (Human Foreskin Fibroblasts) cells. These are  
63 human cells that testify to the toxic activity of an extract. When these cells are in culture for  
64 only 24 hours, they are in a state of mitosis (or dividing cells).

### 65 **2.4 Preparation of plant extracts**

66 After harvest, the fruits were freed of impurities, dried in the shade for a week and then  
67 pulverized with an electric grinder. The fine powders obtained were stored in glass jars to  
68 prevent mold.

### 69 **2.5 Preparation of total aqueous extract (TAE)**

70 The preparation of this extract was performed using the method described by Zirihi *et*  
71 *al*[14] which involves macerating 100 g of plant powder of species in 1L of sterile distilled  
72 water using a Blender type 7 SEVEN STAR. The homogenate was filtered over hydrophilic  
73 cotton and then on Whatman filterpaper (n°3). The aqueous filtrate thus obtained is evaporated  
74 in an oven of Med Center Venticell type at 50°C to obtain powders that constitute the total  
75 aqueous extract (TAE).

### 76 **2.6 Preparation of 70 % ethanolic extract (70 %FE)**

77 The extract was obtained by dissolving 5 g of TAE in 100 mL of ethanol 70% (v ;v) solution  
78 and then homogenized. After decantation and filtration of the alcoholic fraction on hydrophilic  
79 cotton and on Whatman filterpaper (n°3), the filtrate collected is evaporated in an oven at 50  
80 °C. The powder obtained constitutes 70 % ethanolic extract (70 %EE)[15].

### 81 **2.7 Yield calculation**

82 The yield is the quantity of extract obtained from the plant powder. It is expressed as a  
83 percentage. In practice, it has been determined by the ratio of weight of the solids content after  
84 evaporation on the weight of the dry plant material powder used for the extraction, multiplied  
85 by 100. This result is indicated by the following formula:

$$86 \quad Yd = (m \times 100) / M$$

87 Yd : Extraction yield in percentage

88 m : mass in grams of the dry extract

89 M : mass in grams of the drug powder.

## 90 2.8 Phytochemical screening

91 The identification of different chemical compounds in 70 % ethanolic was done by tubes  
92 characterization reactions. This method consists of detecting the different families of chemical  
93 compounds that may exist in plant extracts on the basis of characteristic colorations or  
94 precipitation reactions [16].

### 95 ○ Alkaloids characterization

96 The characterization of alkaloids was made using Bouchard (iodo-iodide) and Dragendorff  
97 (tetraiodopotassium bismuthate) reagent. 6 mL of 70 % ethanolic extract solution was  
98 evaporated to dryness. The residue was taken up in 6 mL of alcohol at 60 °C. The filtrate thus  
99 obtained was divided into two test tubes. In the first tube, two drops of Dragendorff reagent  
100 were added. The presence of alkaloids was characterized by observing orange-coloured  
101 precipitates. In the second tube, two drops of Bouchard reagent was added. The appearance of  
102 a reddish-brown color indicates the presence of alkaloids. A control test was made  
103 with quinine.

### 104 ○ Characterization of polyphenols

105 The polyphenols colorimetry forms colored precipitates with a solution of ferric chloride  
106 ( $\text{FeCl}_3$ ). Thus, one drop of alcoholic solution of 2% ferric chloride and 2 mL of solution of 70 %  
107 ethanolic extract was added. The formation of blue-black or green colouring more or less dark  
108 confirms to the presence of polyphenols. A control test was performed with a solution of  
109 phenol.

### 110 ○ Characterization of flavonoids

111 Flavonoids have been characterized by the reaction to cyanidin. Thus, 2 mL of 70 %  
112 ethanolic extract were evaporated to the dry sand bath. The residue thus obtained was  
113 mixed with 5 mL dilute hydrochloric acid 2 times. The mixture was collected in a test tube, in  
114 which pink-orange or violet colouration will appear. The addition of 3 drops of isoamyl  
115 alcohol intensifies this coloring and confirms presence of flavonoids. An alcoholic solution  
116 of quercetin was used as a control.

### 117 ○ Tannins characterization

118 The Stiasny reagent (Formalin 30%, concentrated HCl 1/0.5) helped to distinguish the  
119 catechin tannins (by precipitation) of gallic tannins (by saturation). Tannins cathéchi-ques: to

120 10 mg of 70 % ethanolic extract, were added 10 mL of Stiasny reagent. The mixture was heated  
121 in a water bath at 80 °C for 30 minutes. After cooling in a stream of water, observation  
122 of precipitate in the form of clear-brown flakes characterizes catechin tannins. An alcoholic  
123 solution of catechin was used as a control. Gallic tannins: For this test, the filtrate obtained  
124 from the reaction of catechol tannins characterization was saturated with sodium acetate. To  
125 this mixture was added a few drops of a dilute aqueous solution of  $\text{FeCl}_3$  at 1% (approximately  
126 1 mL). The appearance of an intense blue-black coloration indicates the presence of gallic  
127 tannins not precipitated by Stiasny reagent. An alcoholic solution of gallic acid was used as  
128 a control.

129 ○ **Terpenes characterization**

130 Sterols and terpenes characterization was made by the Liebermann-Burchard reaction. To 0.2 g  
131 of 70 % ethanolic extract, were added 5 mL of ethyl ether, then the mixture was macerated for  
132 30 minutes. The solution obtained after the maceration was filtered and then evaporated to  
133 dryness. The residue was then dissolved in 0.5 mL of acetic anhydride. Using a pipette, 2 mL  
134 of concentrated sulfuric acid were laid down at the bottom of the test tube without stirring. The  
135 appearance of brownish red or purple ring reflects the two liquid contact zone. The upper  
136 liquid turns green or purple to green or purple indicating the presence of sterols and terpenes.  
137 A control test was performed with progesterone.

138 ○ **Coumarins characterization**

139 For the detection of coumarins, 2 mg of 70 % ethanolic extract was added to 2 mL of warm  
140 water and then homogenized. The homogenate thus obtained was divided into two test tubes.  
141 Thereafter, 0.5 mL of diluted ammonia at 25% was added to the contents of one of the tubes.  
142 After observation under UV 365 nm, the presence of fluorescence in the tube where  
143 ammonia was added indicates the presence of coumarins.

144 ○ **Saponins characterization**

145 For the detection of saponins, 10 mL of 70 % ethanolic extract was introduced in the test  
146 tubes. Each tube was strongly stirred in a vertical position for 15 seconds, and then left  
147 15 minutes. The height of persistent foam is higher than 1 cm, testifying the presence  
148 of saponins.

149 **2.9 Cytotoxicity test**

150 To measure the toxicity of the ethanolic extract, the Human Foreskin Fibroblasts (HFF) cells  
 151 were inoculated in 96-well plates (CellStar) at the rate of 3000 to 5000 cells per well in 100  
 152 µl of D10 medium. These cells are kept in culture for 24 hours (dividing cells). Subsequently  
 153 they were exposed for 24 hours at different concentrations (125-800 µg /mL) in plant extract  
 154 solubilized in PBS buffer. This was done in triplicate, also for control control without plant  
 155 extract. Viability was determined using 3- (4,5-dimethylthiazol-2-yl) -2,5-diphenyl  
 156 tetrazolium bromide (MTT). The tetrazolium ring it contains is reduced in formazan by the  
 157 mitochondrial succinate dehydrogenase of metabolically active cells, which precipitates and  
 158 gives a purple color. The amount of precipitate formed is proportional to the number of living  
 159 cells. In each well, MTT is added at a concentration of 500 µg / mL and incubated for 3 hours  
 160 at 37 ° C. The formazan crystals are solubilized in 10 mM dimethylsulfoxide (DMSO). The  
 161 measurement of the optical density at 544 nm was made using a Safir spectrophotometer  
 162 (Tecan); this measurement of absorbance will make it possible to determine the relative  
 163 quantity of living and metabolically active cells [17]. The results were expressed as a  
 164 percentage of viability compared to control without plant extract. Viability rate = (Abs544  
 165 nm extract / Abs544 nm control) × 100

### 166 3.RESULTS

#### 167 3.1Yield of different extracts of fruits of solanum torvum

168 We obtained from 200 g of powder, 20 g of total aqueous extract a yield of 10 % and from 5 g  
 169 of total aqueous extract, we got 2 g of 70 % ethanolic extract or a yield of 40 %.

#### 170 3.2Phytochemical sorting

171 The phytochemical sorting performed with the extracts of fruits of *Solanum torvum* allowed to  
 172 detect the presence of various chemical groups (Table 1). They are the polyphenols, tannins,  
 173 flavonoids, saponins, and alkaloids in 70% ethanol extract.

174 Table 1 : Chemical compounds in the fruits of *Solanum torvum*

Species	Extract	Chemical compounds							
		Sap	Flav	Terp/ster	Tanins		Coum	Alc	Poly
					Gall	Cathé			
<i>Solanum torvum</i>	EE 70 %	+	+++	-	++	++	-	+	+

175 - :negative reaction ; + : positive reaction

176 EE 70 % :70 % ethanolic extract


177 **Sap:** saponins; **Flav:** flavonoids; **Terp / Ster:** terpenes / sterols; **Gall:** gallic;

178 **Cathé:** cathechic; **Coum:** coumarines; **Alc:** alkaloids; **Poly:** polyphenol

### 179 3.3Cytotoxicity test

180 Figure 2 gives the percentage of viability of the HFF cells cultured in the presence of  
181 concentrations of 100 to 800 µg / mL for the 70% ethanolic extract of the fruits of  
182 *Solanum torvum* compared to the control without plant extract. The number of cells decreases  
183 considerably as the concentration of the 70% ethanol extract of the fruits of *Solanum torvum*  
184 increases. At 800 µg / mL the number of dividing cells is 4%.The averages with the same  
185 superscript letters are not different at 5% according to the turkey test.

186



187  
188

189 Figure 2: Cytotoxicity test of 70%ethanolic extract of fruits of *Solanum torvum* on HFF  
190 dividing cells

191

192

193

194

195

196

#### 197 **4.Discussion**

198 Medicinal plants play a central role in traditional medicine. Ethnobotanical surveys  
199 conducted among traditional health practitioners have made it possible to harvest the fruits of  
200 *Solanum torvum*, which are used in the treatment of anemia, bacterial infections and several  
201 other diseases[18]. The recipes obtained from the fruits of this plant are monospecific, which  
202 is an advantage for the patients, because the associations of wrongly mixed plants, are  
203 sometimes dangerous for the health [19]. Phytochemical screening revealed the presence of  
204 chemical compounds such as alkaloids, tannins, polyphenols, saponins and flavonoids. The  
205 presence of these chemical compounds could justify the multiple activities of the fruits of this  
206 plant [20]. The cytotoxic assay performed on HFF cells showed a gradual decrease in purple  
207 staining in each well. Since the dye penetrates only in living cells, the coloring is weaker as  
208 the plant extract is cytotoxic by inhibition of HFF cells[21]. The sharp decrease in the relative  
209 amount of the dividing HFF cells could be explained by the fact that the HFF cells would be  
210 killed by the 70% ethanol extract of *Solanum torvum*. Indeed, extracts resulting in a cell death  
211 greater than 30% could be considered as cytotoxic[22]. This extract could therefore contain a  
212 chemical compound that inactivates succinate dehydrogenase, an enzyme important for  
213 mitochondrial respiration, the blockage of which would lead to cell death. This result  
214 demonstrates the cytotoxic effect of 70% ethanolic extract of *Solanum torvum*, a Solanaceae  
215 from the Ivorian pharmacopoeia on the cell line tested. Which means that the external use of  
216 the fruits of this plant would probably be dangerous for human health. This toxicity of fruit  
217 could also be explained by the presence of certain groups of chemical compounds such as  
218 glycoalkaloids which are toxic in some Solanaceae[13]. Our results on *in vitro* toxicity  
219 corroborate those [23] who worked on the same family of plants. Indeed according to the  
220 work of Busser and Baies[23] the fruits of *Solanum nigrum* L. (Solanaceae) another  
221 Solanaceae rich in glycoalkaloids and saponins are toxic in internal and external uses on an  
222 organism.

#### 223 **5.Conclusion**



224 *Solanum torvum* is an important medicinal plant of the family of Solanaceae. From the  
225 evaluation of the biological activity, it appears that 70% ethanolic extract of the fruits of  
226 *Solanum torvum* is cytotoxic on the HFF cells. We therefore plan to continue the acute and  
227 subacute toxicity studies to compare the results with those obtained *in vitro* to confirm or  
228 refute the toxicity of the fruits in this study.

229

### 230 **Acknowledgement**

231 The authors thank the Laboratory Adaptation and Pathogenesis of Microorganisms (LAPM)  
232 of Grenoble in France, where cytotoxicity studies were carried out and the traditional health  
233 practitioners of the Haut-Sassandra Region.

### 234 **References**

- 235 1. Adjanohoun J, Aboubakar N, Dramane K, Ebot E, Ekpere A, Enoworock G et al.  
236 Traditional medicine and pharmacopeia-contribution to ethnobotanical and floristic studies  
237 in Cameroon. In: CNPMS. Porto-Novo, Benin, 1996, pp. 50–52.
- 238 2. Etienne J. Solanacées médicinales et philatélie. Bull. Soc. Pharm. Bordeaux. 2005 ; 144: 311-  
239 332.
- 240 3. Emmanuel N, Catherine K, Philomène C N B, Siegfried D D, Emmanuel M M. Inventaire  
241 et caractérisation des plantes médicinales utilisées en thérapeutique dans le département de la  
242 Sanaga Maritime: Ndom, Ngambe et Pouma. J. Appl. Biosci. 2016 ; 106:10333 –10352.
- 243 3. Siemonsma J, Piluek K. Plant Resources of South-East Asia (PROSEA) Bogor. Indonesia,  
244 1994, pp. 412.
- 245 4. Kala C P. Ethnomedicinal botany of the Apatani in the Eastern Himalayan region of India. J  
246 Ethno and Ethnomed. 2005 ; 1: 1-8.
- 247 5. Sivapriya M, Srinivas L. Isolation and purification of a novel antioxidant protein from the  
248 water extract of Sundakai (*Solanum torvum*) seeds. Food Chemistry. 2007 ; 104: 510 - 517.
- 249 6. Nasir, J.Y. Solanaceae. In: Flora of Pakistan. (Eds.): S.I. Ali and E. Nasir. Fascicle 85.  
250 Pakistan Agri Res Council. 1985 ; pp. 1-61.
- 251 7. Wiat C., Mogana S., Khalifah S., Mahan M., Ismail S., Buckle M., Narayana A.K and  
252 Sulaiman M. Antimicrobial screening of plants used for traditional medicine in the state of  
253 Perak, Peninsular Malaysia. Fitoterapia. 2004; 75 (1): 68-73.
- 254 8. Fui LH, Knowledge and use of forest product as traditional medicine: the case of the  
255 forest-dwelling communities, In: Proceedings of the Conference on Medicinal Products from

- 256 Tropical Rain Forest. K Shaari, AA Kadir, ARM Ali (Eds.), Forest Research Institute of  
257 Malaysia. Kuala Lumpur. 1992 ; pp. 385-400.
- 258 9. Mohan M, Kamble S, Kasture S. Protective effect of *Solanum torvum* on doxorubicin-  
259 induced nephrotoxicity in rats, Food and Chem Toxicol. 2010 ; 48: 436-440.
- 260 10. Gandhi GR, Ignacimuthu S, Paulraj M G, Sasikumar P. Antihyperglycemic activity  
261 and antidiabetic effect of methyl caffeate isolated from *Solanum torvum* Swartz. fruit  
262 in streptozotocin induced diabetic rats, Eur J Pharmacol. 2011 ; 30: 623-31.
- 263 11. Pérez-Amador M. C, Muñoz Ocotero V, García J M, Castañeda A R, González E.  
264 Alkaloids in *Solanum torvum* Sw (Solanaceae). Inter experim Botany. 2007; 76: 39-45.
- 265 12. Tajner-Czopek A. Food Chem. 2008 ; 106 (2) :706-11.
- 266 13. Ayad K. Effect of Solanine on Arthritis Symptoms in Postmenopausal Female Albino  
267 Rats. Arab Journ of Nucl Sc and Appl. 2013 ;46(3) :279-285.
- 268 14. Zirihi GN, Kra A, Dadié ET. Etude botanique et évaluation des activités antifongiques de  
269 *Mitracarpus villosus* (MV) (Rubiaceae) et *Spermacoce verticillata* (SV) (Rubiaceae) sur la  
270 croissance *in vitro* de *A. fumigatus*. Rev de Méd et de Pharma Afr. 2007 ; 20 : 9-17.
- 271 15. Zirihi GN, Kra AKM. Evaluation de l'activité antifongique de *Microglossa pyrifolia*  
272 (Lamarck) O. Kuntze (Asteraceae) « PYMI » sur la croissance *in vitro* de *Candida albicans*.  
273 Rev Med Pharmacol Afr. 2003; 17 : 11-8.
- 274 16. Harborne J B. A guide to modern techniques of plant analysis. Springer, 3rd Edn, India  
275 (New Delhi). 1998, pp 5-32.
- 276 17. Mosman, T. Rapid colorimetric assay for cellular growth and survival : application to  
277 proliferation and cytotoxicity assay. Journal of Immunological Methods. 1983 ; 65, 55-63.
- 278 19. N'Guessan K, Kouadio K, Kouamé N'Guessan F, Traoré D, Aké-Assi L. Etude botanique  
279 des plantes emménagogues utilisées en médecine traditionnelle par les Abbey et Krobou  
280 d'Agboville (Côte-d'Ivoire). Rev Med Pharm Afr. 2008 ; 21: 43-60.
- 281 20. Zubaida Y, Ying W, Elias B. Phytochemistry and Pharmacological Studies on *Solanum*  
282 *torvum* Swartz Jour of Applied Pharma Sci. 2013 ; 3 (4), pp. 152-160.
- 283 21. Irie-N'guessan A. G, Kablan B. J., Kouakou-Siransy N. G., Leblais V., Champy P.  
284 Evaluation de la toxicité de cinq plantes antiasthmatiques de la médecine traditionnelle  
285 ivoirienne. Int. J. Biol. Chem. Sci. 2011; 5(3): 1316-1319.

286 22.Coulerie P. Etude phytochimique et pharmacologique de plantes de Nouvelle-Calédonie à  
287 potentialités anti-dengue. Thèse de Doctorat de chimie des substances naturelles. Université  
288 de la Nouvelle-Calédonie : Ecole Doctorale du Pacifique. 2012 ; 296 p.

289 23.Busser C. Baies, fruits et pseudo-fruits toxiques utilisés en médecine populaire ou en  
290 phytothérapiePhytot. 2007 ; 1: 31–36

291

292

293

294 ,

295

296

297

298

299

300