

**Using 7.1 chlorhexidine gel for umbilical cord care: implication for WHO's
recommendation for a standard cord care practice**

ABSTRACT

Neonatal mortality has been a bane especially in the developing world. Neonatal infection is the second cause of neonatal death after prematurity. The umbilical cord is one such vulnerable route via which newborns are infected, the responsible organisms most likely originate from the maternal genital tract and are acquired during labor and delivery. Such high-infection-related deaths make it necessary to identify simple and cost-effective interventions that can be implemented in all settings across the world. WHO has recommended the use of 7.1% chlorhexidine gel in areas of high neonatal mortality to replace the use of harmful substances on the umbilical stump. However, in Nigeria, studies are needed to establish the umbilical cord separation time, mothers' compliance and satisfaction with use of chlorhexidine.

Keywords: chlorhexidine gel, umbilical cord, omphalitis, neonatal mortality

Introduction

Each year, about 3 million newborns die globally (Liu, Johnson et al, 2012) accounting for 40 % of global under-five mortality (Nangia, Dhingra et al, 2016); this is of great concern especially in developing countries. Annually, serious infections account for nearly 30 % of the 3 million neonatal deaths (Sankar, Chandrasekaran et al 2016); in settings with high-mortality rates, this proportion is closer to 50 % (Oestergaard, Inoue et al, 2011). Such high-infection-related deaths make it pertinent therefore to identify simple and cost-effective interventions that can be implemented in all settings across the world. Use of topical antiseptics is one such intervention that is been envisaged to reduce the incidence of infections by preventing or limiting bacterial colonization of the skin or umbilical cord in newborns. Five countries namely India, Nigeria, Congo, Pakistan and China alone account for half (2.440 million) of the global deaths from infections and 53.3 % (1.636 million) of neonatal deaths (Liu et al, 2012). Community based



studies from developing countries found that infection accounts for about 8 to 80 % of all neonatal mortality, and as many as 42 % deaths in first week of life (Thaver & Zaidi, 2009).

In Nigeria, an estimated 276,000 neonatal deaths occurs annually; 60,000 of these deaths stem from infection, while about 20,000 of these deaths are umbilical cord related (Chorhexidine Working Group (CWG), 2016). Most of these deaths occur in rural areas and northern Nigeria (Akinyemi, Bamgboye et al, 2015). Nigeria has the second highest number of neonatal deaths in the world after India (CWG, 2016). The Nigeria demographic and health survey (NDHS) 2013, estimated its neonatal mortality rate as 37 per 1000 live births, which constituted about 54% of infant mortality. Nigeria contributed 6% to the global neonatal deaths in 2005, while the country moved from the third to the second position in terms of the highest number of neonatal deaths in the world between 2000 and 2010 (Lawn, Kinney et al, 2012).

Omphalitis is an important cause of neonatal mortality and preventing this condition and its associated neonatal mortality is of high public health importance. While the use of clean birth kits, hand washing and careful attention to hygiene in the days after birth may all be important components of a program to reduce neonatal sepsis; however, none of these interventions have to date provided unequivocal evidence that they can reduce infection-related neonatal mortality. The three randomized controlled trial systematically reviewed by Imdad, Mullany et al (2013) provided substantial evidence that umbilical cord stump treatment with 4% chlorhexidine (CHG) reduces both omphalitis and neonatal mortality, with the impact likely greater in preterm and low birth weight infants.

The umbilical stump: an important locus of infection in newborns

Newborns are particularly prone to infections for a number of reasons which includes a relatively immature immunological system, exposure to maternal vaginal organisms during the birth process and exposure to various pathogens in the first few days of life, often transmitted by physical contact with the caregivers, delivery surfaces and instruments. One vulnerable area for infection for all infants is the umbilical cord stump usually cut, tied/clamped shortly after birth, and allowed to dry and detach spontaneously after some days, the cord stump provides necrotic tissue for organisms to colonize and a point of easy entrance of microbes to the neonate (Goldenberg, McClure & Saleem, 2014).

The umbilical stump acts as a bacterial reservoir especially in first few days of life, supporting the growth of some beneficial microorganisms (commensals) whereas others are harmful (for example, clostridium tetani). Umbilical Cord infection can be limited to the umbilical cord area (omphalitis) or, after entry into the blood stream via the umbilical vein becomes systemic (neonatal sepsis) (Nangia et al, 2016). The responsible organisms most likely originate in maternal genital tract and are acquired during labor and delivery. In most developing countries, many neonatal infections are environmentally acquired because of higher number of home deliveries, unsafe traditional practices, unskilled birth attendants and unclean living conditions, all of which pose an increased risk of umbilical cord infection (Nangia et al, 2016). Local signs of umbilical cord infection may include redness, tenderness, swelling, pus, warmth and foul odour. Infectious organisms may get directly transmitted from patent umbilical vein to the systemic circulation without evident sign of local cord infection.

Current World Health Organization recommendation for umbilical cord care

In January 2014, the WHO issued a new recommendation for umbilical cord care. *Daily application of 7.1% chlorhexidine digluconate aqueous solution or gel, delivering 4% chlorhexidine to the umbilical cord stump during the first week of life for newborns delivered at home in settings with high neonatal mortality (30 or more neonatal deaths per 1000 live births). While clean, dry cord care is recommended for newborns delivered in health facilities and at home in low neonatal mortality settings. The use of chlorhexidine in these situations may be considered only to replace application of a harmful traditional substance, such as cow dung, to the umbilical cord stump* (WHO, 2013).

WHO in 1998 initially recommended dry umbilical cord care (that is, not applying anything to the cord); but study by Chlorhexidine working group (2016) has shown that 90% of mothers apply one substance or the other to the umbilical stump. This practice **of mandatorily** using substances on the umbilical cord appears to have a psychological and/or cultural inclination that requires thorough health education to curb. It was on this premise that WHO recommended use of chlorhexidine gel if at all something must be applied to the cord stump.

Example of some traditional cord care practices that called for WHO's recommendation of a standard cord care agent

Various health institutions and cultures adopt different methods of cord care. While most orthodox health institutions in Nigeria currently advocate the use of methylated spirit, WHO (2013) recently introduced 7.1% chlorhexidine in countries with high neonatal mortality; while advising no treatment at all but keeping the cord clean and dry for developed countries. Some cord care practices in some countries include:

- **Nigeria:** non beneficial practices such as use of hot compress, herbs, native chalk, salt, sand, saliva, palm oil, menthol-containing balm, petroleum jelly and toothpaste - substances were used alone or in combination and mainly applied at home (Abhulimhen-Iyoha & Abidin, 2012). Others include fomentation with dry heat, application of white powder, charcoal and bandaging to prevent infection (Adejuyigbe, Odebiyi et al, 2008)
- **Turkey:** use of dry coffee, sugared fat, olive oil, rotten tree powder, myrtle, hellebore, black sesame, burnt cloth, tying the belly with a rope and putting a buttered cloth over the infant's infected belly (Alparslan, Demirel et al, 2013).
- **Zambia:** Different substances had different purposes. If cord is too brittle, cracking, bleeding, then a substance that increases softness is been used (Vaseline, cooking or motor oil, *mabono* (wild fruit) oil, or cream from sour milk). If cord takes too long to separate, items to dry it are used such as: baby powder, charcoal dust, dried cow dung, dried chicken droppings (roaster for male babies and hen for female babies), dust from threshold of home, ash from burnt pumpkin stem, crushed *loma* (wasps nest), or mud. Any of these would be pounded or ground into a fine powder. Medicinal substances: python snake oil, breast milk, banana, cow dung, *mukunku* (tree bark), traditional herbs, and dirt from pounding stick are also used (Herlihy, Shaikh et al, 2013)
- **Bangladesh:** Turmeric, boric powder, mustard oil, ash, coconut oil, Nebanol ointment, ginger, chewed rice, heat treatment of cord (Alam, Ali et al, 2008); talcum powder, Savlon, *chular mati* (earth from a clay oven), smashed garlic (Moran, Choudhury et al, 2009).
- **Uganda:** Herbs, onions, ash from burnt papyrus, petroleum jelly, powder, saliva, ghee, soot mixed with ghee, water, butter (Byaruhanga, Nsungwa-Sabiiti et al, 2011); salty water, vaseline, normal saline, ripe banana, sap, ash (Kayom, Kakuru et al, 2015); lizard droppings (Waiswa, Peterson et al, 2010)

- **Tanzania:** Saliva, dirty doorpowder from old door, hotknife, charcoal powder, ground sea shells, burning wood, banana steam, fish bone, talcum powder, and fire steam (Dhingra, Gittelsohn et al, 2014); traditional herbs + cooking oil or water that was used to wash the genitals of an elderly woman (Mrisho, Schellenberg et al, 2008)
- **Pakistan:** Coconut oil, mustard oil, ghee, olive oil, surma/kohl, turmeric, machine oil, Antimony, detol and wheat flour (Gul, Khalil et al, 2014)
- **Napel:** Mustard oil, antiseptic, ash, mud, breast milk, herbs/spices, saliva (Karas, Mullany et al, 2012), plain water, toothpaste (Sharma, van Teijlingen et al, 2016).
- **India:** Turmeric, burning tip with castor oil lamp, antiseptic ointment (Kesterton, Cleland et al, 2009).
- **Haiti:** dirt from house threshold, burnt nutmeg, ash burned cotton, crushed charcoal, palm oil, mixture of leaves and animal dung (Walsh, Norr et al, 2015)
- **Sierra Leone:** pounded cassava (Sharkey, Yansaneh et al, 2016).

Caregivers' reasons for cord care

Studies by (Amare, 2014) found out two main reasons behind caregivers' intention of applying a substance to the umbilical cord. The first reason is to promote healing of the so called "unique wound"; secondly, is to hasten the separation of the cord either by keeping the cord stump moist or by drying it out stimulating its occurrence within the first few days of life despite the fact that studies by Imuetinyan (2011) have shown that normal cord separation time can extend for as long as 28 days in some cases. Other reasons include preventing "wind" (evil spirit) from going into the baby: to prevent pains and bad smell (Degefe, Amare et al, 2014). In Africa, mothers are often desperate to hasten the time of cord dropping; the longer time the umbilical cord takes to drop, the more likely it is that a more harmful intervention is introduced, and the greater the risk of infection. They tend to indulge in various cultural practices to hasten even the time for a healthy normal cord to separate thereby, introducing infection. This uncalled-for interventions is common in some cultures where the mother or baby's activity of daily living is restricted (for example, not allowing the mother or baby to go out, delaying baby's bath) until the umbilical stump falls off.

Chlorhexidine digluconate

Chlorhexidine digluconate is the gluconate salt form of chlorhexidine, a biguanide compound used as an antiseptic agent with topical antibacterial activity. Chlorhexidine gluconate is positively charged and reacts with the negatively charged microbial cell surface, thereby destroying the integrity of the cell membrane. Subsequently, chlorhexidine gluconate penetrates into the cell and causes leakage of intracellular components leading to cell death (National Center for Biotechnology Information, 2016). Chlorhexidine has the following advantageous properties as a topical cord care agent: it released slowly, longer duration of action, not affected by the presence of body fluids, dissolve easily in water, it has an excellent safety profile, is rarely associated with bacterial resistance, is easy to administer and costs few naira per application (Nangia et al, 2016). Available high-quality evidence from the recently conducted Cochrane review indicates that cord cleansing with 4 % chlorhexidine reduces the risk of neonatal mortality by 12 % and omphalitis/infections by 50 % in low-resource community settings including Nepal, Bangladesh, and Pakistan (Sinha, Sazawal et al, 2015). However, the review showed that in hospital settings, chlorhexidine cord cleansing reduces the risk of omphalitis/infections by 52 % and may lead to no difference in neonatal mortality as compared to dry cord care. In spite of these benefits, studies by Imdad, Bautista et al (2013) have shown that it prolongs the time for the cord to drop when compared with other methods like dry cord care. Contrary to the findings from South Asia countries, the only two randomized controlled trials conducted in Africa (Tanzania and Zambia) on chlorhexidine use gave no significant reduction in neonatal mortality when compared to dry cord care.

In Nigeria, various state government are investing into the lives of the citizens; programs has been put up to scale up the use of chlorhexidine with Sokoto **stategovernment** taking the lead, followed by Bauchi and Ogun state government. Recently, on 14th March, 2017 the governor of Kogi state flagged off the adoption of chlorhexidine gel for umbilical cord care to replace the use of methylated spirit in the state with the vision of reducing neonatal mortality.

Now how do we reconcile these high evidence studies from South Asia countries and those conducted in Africa so as to implement best policies? If the use of chlorhexidine gel prolongs umbilical cord separation time when compared with other cord care methods, it means that desperate mothers are likely to introduce other harmful methods to hasten the cord separation thereby, increasing the likelihood of infection thus, the essence WHO introduced chlorhexidine

gel is defeated. Also if there is no significant difference in neonatal mortality between the use of chlorhexidine and dry cord care in settings of high neonatal mortality (more than 30 in 1000), then there is no point incurring economic cost from the purchase of chlorhexidine. However, a detailed pooled meta-analysis from different settings across the globe is needed to inform policies.

Recommendation

Chlorhexidine gel for umbilical cord care is still a novel intervention in Nigeria, and no study was found on its use in Nigeria; More evidence based studies will be of benefit in the following areas: its effect on cord separation time, effect on mortality/morbidity, Mother's compliance and satisfaction with use, factors that will impede its use and its effect on very low birth weight babies.

Conclusion

The desire to care for the umbilical cord has been found to be universal culturally; a myriad of substances are used by different cultural groups. Though some of these substances are not totally harmful, but their therapeutic effect has not been established empirically. WHO has recommended the use of 7.1% chlorhexidine gel especially in settings with high neonatal mortality. To guarantee that mothers will use this gel judiciously as recommended; it is pertinent therefore, to put up more studies on this novel cord care agent so as to inform better policies.

References

- Abhulimhen-Iyoha, B.I., Ibadin, M.O. 2012. Determinants of cord care practices among mothers in Benin City, Edo State, Nigeria. *Niger J Clin Pract.* 152:210–3. doi:10.4103/1119-3077.97320.
- Adejuyigbe, E.A., Odebiyi, A.I., Aina, O., Bamiwuye, S. 2008. Feeding and care of low-birth weight babies in two rural communities in south-western Nigeria. *Matern Child Nutr.* 4.1.:55–64. doi: 10.1111/j.1740-8709.2007.00101.
- Akinyemi, J.O., Bamgboye, E.A., Ayeni, O. 2015. Trends in neonatal mortality in Nigeria and effects of bio-demographic and maternal characteristics. *BMC Pediatrics* 15:36-40
- Alam, M.A., Ali, N.A., Sultana, N., Mullany, L.C., Teela, K.C., Khan, N.U, et al. 2008. Newborn umbilical cord and skin care in Sylhet District, Bangladesh: implications for the

- 202 promotion of umbilical cord cleansing with topical chlorhexidine. *J Perinatol.* 28.2:61–8.
203 doi: 10.1038/jp.2008.164.
- 204 Alparslan, Ö., Demirel, Y. 2013. Traditional neonatal care practices in Turkey. *Jpn J Nurs Sci.*
205 10.1:47–54. doi: 10.1111/j.1742-7924.2012.00209.x.
- 206 Amare, Y. 2014. Umbilical cord care in Ethiopia and implications for behavioral change: a
207 qualitative study. *BMC Int Health Hum Rights.* 14:12. doi:10.1186/1472-698X-14-12
- 208 Byaruhanga, R.N., Nsungwa-Sabiiti, J., Kiguli, J., Balyeku, A., Nsabagasani, X., Peterson, S.
209 2010. Hurdles and opportunities for newborn care in rural Uganda. *Midwifery.* 27.6:775–
210 80. doi: 10.1016/j.midw.2010.02.005.
- 211 Chlorhexidine Working Group (CWG) 2016. 7.1% Chlorhexidine Digluconate for Umbilical
212 Cord Care: Introduction and scale up. [www.coregroup.org/.../CORE WEBINAR](http://www.coregroup.org/.../CORE_WEBINAR)
213 (accessed 26.09.2016)
- 214 Degefie, T., Amare, Y., Mulligan, B. 2014. Local understandings of care during delivery and
215 postnatal period to inform home based package of newborn care interventions in rural
216 Ethiopia: a qualitative study. *BMC Int Health Hum Rights.* 14:17. doi:10.1186/1472-
217 698X-14-17.
- 218 Dhingra, U., Gittelsohn, J., Suleiman, A.M., Suleiman, S.M., Dutta, A., Ali, S.M., et al. Delivery,
219 immediate newborn and cord care practices in Pemba Tanzania: a qualitative study of
220 community, hospital staff and community level care providers for knowledge, attitudes,
221 belief systems and practices. *BMC Pregnancy Childbirth.*
- 222 Goldenberg, R.L., McClure, E.M. and Saleem, S. 2014. A review of studies with chlorhexidine
223 applied directly to the umbilical cord. *Am Journal of Perinatology.* 30.8:35-75. Retrieved
224 Sept. 26, 2016 from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3875170/>
- 225 Gul, S., Khalil, R., Yousafzai, M.T., Shoukat, F. 2014. Newborn care knowledge and practices
226 among mothers attending pediatric outpatient clinic of a hospital in Karachi, Pakistan. *Int*
227 *J Health Sci (Qassim)* 8.2:167–75. doi: 10.12816/0006083.
- 228 Herlihy, J.M., Shaikh, A., Mazimba, A., Gagne, N., Grogan, C., Mpamba, C. et al. 2013. Local
229 perceptions, cultural beliefs and practices that shape umbilical cord care: a qualitative
230 study in Southern Province, Zambia. *PLoS One.* 8.11:e79191.
231 doi:10.1371/journal.pone.0079191.
- 232 Imdad, A., Bautista, R.M., Senen, K.A., Uy, M.E., Mantaring, J.B. and Bhutta, Z.A. 2013.
233 Umbilical cord antiseptics for preventing sepsis and death among newborns. *Cochrane*
234 *Database Syst Rev.* 31.5:237-286. Retrieved Sept. 20, 2016 from
235 <https://www.ncbi.nlm.nih.gov/m/pubmed/23728678/?i=4&from=/23254380/related>
- 236 Imdad, A., Mullany, L.C. Baqui, A.H., Arifeen, S.E., Tielsch, J.M., Khatry, S.K., Shah, R.,
237 Cousens, S., Black, R.E. and Bhutta, Z.A. 2013. The effect of umbilical cord cleansing

- 238 with chlorhexidine on omphalitis and neonatal mortality in community settings in
239 developing countries: a meta-analysis. *BMC Public Health* 13.3:3-15.
- 240 Karas, D.J., Mullany, L.C., Katz, J., Khatry, S.K., LeClerq, S.C., Darmstadt, G.L. et al. 2012.
241 Homecare practices for newborns in rural southern Nepal during the first 2 weeks of life.
242 *J Trop Pediatr*. 58.3:200–7. doi: 10.1093/tropej/fmr057.
- 243 Kayom, V.O., Kakuru, A., Kiguli, S. 2015. Newborn care practices among mother-infant dyads
244 in urban Uganda. *Int J Pediatr*. 2015:815938. doi:10.1155/2015/815938.
- 245 Kesterton, A.J., Cleland, J. 2009. Neonatal care in rural Karnataka: healthy and harmful
246 practices, the potential for change. *BMC Pregnancy Childbirth*. 9:20. doi:10.1186/1471-
247 2393-9-20.
- 248 Lawn, J.E., Kinney, M.V., Black, R.E., Pitt, C., Cousens, S., Kerber, K., Corbett, E., Moran,
249 A.C., Morrissey, C.S. and Oestergaard, M.Z. 2012. Newborn survival: a multi-country
250 analysis of a decade of change. *Health Policy Plan* 27.3:6-28. Retrieved Oct. 6, 2016
251 from <https://www.ncbi.nlm.nih.gov/m/pubmed/22692417/>
- 252 Liu, L., Johnson, H.L., Cousens, S., Perin, J., Scott, S., Lawn, J.E., Rudan, I., Campbell, H.,
253 Cibulskis, R., Li, M., Mathers, C. and Black, R.E. 2012. Global, regional, and national
254 causes of child mortality: an updated systematic analysis for 2010 with time trends since
255 2000. *Lancet* 379.12:2151–2161. Retrieved Sept. 10, 2016 from
256 <https://www.ncbi.nlm.nih.gov/m/pubmed/22579125/>
- 257 Moran, A.C., Choudhury, N., Uz Zaman Khan, N., Ahsan Karar, Z., Wahed, T., Faiz Rashid, S.
258 et al. 2009. Newborn care practices among slum dwellers in Dhaka, Bangladesh: a
259 quantitative and qualitative exploratory study. *BMC Pregnancy Childbirth*. 9:54. doi:
260 10.1186/1471-2393-9-54.
- 261 Mrisho, M., Schellenberg, J.A., Mushi, A.K., Obrist, B., Mshinda, H., Tanner, M. et al. 2008.
262 Understanding home-based neonatal care practice in rural southern Tanzania. *Trans R*
263 *Soc Trop Med Hyg*. 102.7:669–78. doi: 10.1016/j.trstmh.2008.04.029.
- 264 Nangia, S., Dhingra, U., Dhingra, P., Dutta, A., Menon, V.P. and Black, R.E. 2016. Effect of 4 %
265 chlorhexidine on cord colonization among hospital and community births in India: a
266 randomized controlled study. *BMC Pediatrics* 16.121:0625-0627. Retrieved Sept. 13,
267 2016 from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4971644/>
- 268 National Center for Biotechnology Information. 2016. Chlorhexidine Digluconate. PubChem
269 Compound Database. Retrieved Sept. 13, 2016 from
270 <https://pubchem.ncbi.nlm.nih.gov/compound/9552081>
- 271 Nigeria Demographic and Health Survey. 2013. National Population Commission, Federal
272 Republic of Nigeria Abuja. Retrieved Sept. 5, 2016 from
273 <http://dhsprogram.com/pubs/.../FR293.p..>

- 274 Oestergaard, M.Z., Inoue, M., Yoshida, S., Mahanani, W.R., Gore, F.M., Cousens, S., Lawn, J.E.
275 and Mathers, C.D. 2011. Neonatal mortality levels for 193 countries in 2009 with trends
276 since 1990: a systemic analysis of progress, projections and priorities. *PLoS Med.*
277 8(8):105- 189.Retrieved Sept. 5, 2016 from
278 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3168874/>
- 279 Sankar, M.J., Chandrasekaran, A., Ravindranath, A., Agarwal, R. and Paul, V.K. 2016.
280 Umbilical cord cleansing with chlorhexidine in neonates: a systematic review. *Journal of*
281 *Perinatology* 36.1:12–20. Retrieved Sept. 5, 2016 from
282 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4848738/>
- 283 Sharkey, A., Yansaneh, A., Bangura, P.S., Kabano, A., Brady, E., Yumkella, F. et al. 2016.
284 Maternal and newborn care practices in Sierra Leone: a mixed methods study of four
285 underserved districts. Health Policy Plan. 2016
- 286 Sharma, S., van Teijlingen, E., Hundley, V., Angell, C., Simkhada, P. 2016. Dirty and 40 days in
287 the wilderness: eliciting childbirth and postnatal cultural practices and beliefs in Nepal.
288 BMC Pregnancy Childbirth. 16.1:147. doi: 10.1186/s12884-016-0938-4.
- 289 Sinha, A., Sazawal, S., Pradhan, A., Ramji, S. and Opiyo, N. 2015. Chlorhexidine skin or cord
290 care for prevention of mortality and infections in neonates. *Cochrane Database*
291 *Systematic Review* 5.3:450-501. Retrieved Sept. 5, 2016 from
292 <https://www.ncbi.nlm.nih.gov/m/pubmed/25739381/>
- 293 Thaver, D. and Zaidi, A.K. 2009. Burden of neonatal infections in developing countries: a
294 review of evidence from community-based studies. *Pediatric Infectious Disease Journal*
295 28.1:3–9. Retrieved Sept. 5, 2016 from
296 <https://www.ncbi.nlm.nih.gov/m/pubmed/19106760/>
- 297 Waiswa, P., Peterson, S., Tomson, G., Pariyo, G.W. 2010. Poor newborn care practices – a
298 population based survey in eastern Uganda. BMC Pregnancy Childbirth 10:9.doi:
299 10.1186/1471-2393-10-9.
- 300 Walsh, S., Norr, K., Sankar, G., Sipsma, H. 2015. Newborn cord care practices in Haiti. *Glob*
301 *Public Health*. 10.9:1107–17. doi: 10.1080/17441692.2015.1012094.
- 302 Winch, P.J., Alam, M.A., Akther, A., Afroz, D., Ali, N.A., Ellis, A.A., Bangladesh
303 PROJAHNMO Study Group et al. 2005. Local understandings of vulnerability and
304 protection during the neonatal period in Sylhet District, Bangladesh: a qualitative study.
305 *Lancet*. 366.9484:478–85. doi: 10.1016/S0140-6736(05)66836-5.
- 306 World Health Organization. 2013. WHO Recommendations on Postnatal Care of the
307 Mother and Newborn. Retrieved Sept. 5, 2016 from [www](http://www.healthynewbornnetwork.org/bitstream/10665/97603/1/9789241506649_eng.pdf)
308 [.healthynewbornnetwork.org/bitstream/10665/97603/1/9789241506649_eng.pdf](http://www.healthynewbornnetwork.org/bitstream/10665/97603/1/9789241506649_eng.pdf)