

## **SOCIO-ECONOMIC FACTORS ASSOCIATED WITH PATIENTS WITH VESICOVAGINAL FISTULA IN MACHAKOS COUNTY, KENYA**

### **Abstract**

*Background:* Vesico-Vaginal Fistula (VVF) is traumatic fistulous tract extending between the bladder and the vagina that allows the continuous involuntary discharge of urine into vaginal vault. The patient has urine wetting and soaking her innerwear, before dripping down her legs. The accompanying smell is awful and most communities consider such women as outcasts.

*Methods:* This study was conducted among 34 patients with Vesico-vaginal fistula purposively selected from 10 sub counties of Machakos County, Kenya. To supplement the results from these patients, their case notes, 14 of the patients' husbands, 25 women group leaders from the County, 48 Community Health Workers (CHWs), and 177 healthcare workers in the County were interviewed.

*Results:* Their ages ranged between 15 and 35 years with mean and median years of 22.3 years. A majority of the patients (62%, n=21) were below 23 years, (29%, n=10) had no formal education and (82%, n=28) came from households earning less than sh6, 000 a month. All of them developed the complication as they delivered at home with either the assistance of a CHW or a relative. By the time the study was carried out, they had sought after treatment at a Referral Hospital and were awaiting appointments for surgical repair. Out of the 14 with husbands, all had monogamous marriages with (71.4%, n=10) of the husbands reporting that they tolerated their wives' conditions but could not sleep together due to the stench. The women group leaders reported that these patients were shy to face the society and tended to lead a secluded life of their own away from the glare of society. The healthcare workers attributed the VVF condition mainly from obstructed labor due to Malpresentation of the baby at delivery, Cephalopelvic disproportion (CPD), maternal distress or Cervical dystocia.

*Conclusion:* Vesico- vaginal fistula is a problem that occurs mainly amongst the young and poor mothers with prolonged obstructed labor which is not supervised by skilled medical attendants. Its prevalence can be reduced through enlightenment of the public and ensuring that mothers deliver their babies under safe hands of skilled birth attendants.

**Keywords:** Vesico-vaginal fistula, Machakos County, Socio-Economic Factors, Obstructed labor.

### **Introduction**

Vesicovaginal fistula (VVF) is an anomalous fistulous tract extending between the bladder and the vagina that allows the continuous involuntary discharge of urine into vaginal vault <sup>(1)</sup>. It is one of several vaginal fistulae such as Ureterovaginal, Rectovaginal, Colovaginal and

Enterovaginal fistulae. The patient has uncontrolled urine leak that leads to wetting and soaking of her innerwear causing unpleasant odor, before dripping down her legs. This condition has significant effects on the patient's self-esteem and leads to social stigma and often times when the condition occurs she is neglected by the family and community. She may become depressed as a result of loss of husband's affection, divorce or childlessness especially if it results from still births. Most communities consider such women as outcasts and may blame them of witchcraft and other wrongdoing. Prolonged obstructed labor is associated with pressure necrosis, edema and tissue sloughing and is the leading cause of VVF worldwide. It is responsible for 97% of VVF cases in developing countries. In the developed world, 90% of VVF cases are estimated to be secondary to accidents during gynecological surgeries such as pelvic operations and in hysterectomies where the bladder or urethra may be injured. Obstructed labor occurs when the uterus contracts normally during childbirth but the baby does not pass through the birth canal. It could occur in conditions like a big baby, an abnormally positioned baby or a small maternal pelvis or birth canal. Cultural factors that may lead to obstruction of the reproductive system include Female Genital Mutilation (FGM), making of incisions on the vaginal wall and application of substances usually with an aim of returning it to nulliparous state. Other less frequent causes of VVF include pelvic infections and trauma from insertion of foreign objects particularly in crude attempts to procure abortion<sup>(2)</sup>. Confounders would be anemia, malnutrition, unhygienic environment, and drugs that may compromise healing process. High prevalence of early marriage and subsequent childbearing, low socioeconomic status of women, lack of skilled birth attendance and lack of access to emergency obstetric services are associated risk factors. The study was carried out in Machakos County which is one of the 47 Counties in Kenya. It aimed at establishing the factors that were associated with Vesicovaginal Fistula cases. Such factors were categorized as socio-cultural, economic, programmatic and infrastructural.

## MATERIALS AND METHODS

This descriptive cross sectional study was carried out between January 15<sup>th</sup> to July 20<sup>th</sup> of 2015 in the 10 Sub Counties Machakos County, Kenya (latitudes 0° 45' South to 1°31' South and longitudes 36° 45' East to 37° 45' East), a county that is mostly semi arid and covers about 6,300 square kilometers. Clearance to conduct the study was provided by Kenyatta University graduate school, NACOSTI and the County Medical Officer of Health (CMOH). Thirty four patients with Vesico-vaginal fistula were purposively selected from 10 sub counties of the county and every individual's information was studied. Further information was collected from 14 of the patients' husbands, 25 women group leaders from the County, 48 Community Health Workers (CHWs), and 177 healthcare workers in the County. The data collected included the patients age, marital status, parity, education background, monthly income and on their occupation. Further information was sourced on perceived type, cause, duration and adjustment practices in living with the condition. Researcher facilitated questionnaires were completed; interviews with the respondents were conducted and key informant interviews and focus group discussions were carried out. Collected data was coded and descriptive statistics, cross-tabulation, and regression analysis were analyzed using SPSS Version 20.

87 The ages of the patients ranged between 15 and 35 years with mean and median years of 22.3  
 88 years. A majority of the patients (62%, n=21) were below 23 years as presented in (Table 1).  
 89 There was a highly significant negative correlation ( $r=-0.9339$ ,  $p<0.01$ ) between the incidences  
 90 of VVF and age of the patient. Perusal of the patients' hospital reports and report from the  
 91 healthcare workers attributed the VVF condition mainly from obstructed labor due to Mal-  
 92 presentation of the baby at delivery, Cephalopelvic disproportion (CPD), maternal distress or  
 93 Cervical dystocia. The study observed that (29%, n=10) had no formal education and even  
 94 among the formally educated, less than 20%, (17.6%, n=6) had post primary education (Table 2).  
 95 The mean length of time the patients had lived with the condition was about 3.5 years with about  
 96 60% of them not exceeding 4 years (Table 3). The mean age of the patient at the time they  
 97 developed the condition was found to be 18.8 years with about 95% developing it when their age  
 98 was below 27 years of age (Table 4). The study showed that most of the patients (82%, n=28)  
 99 came from households earning less than sh6, 000 a month with a mean monthly earning of about  
 100 sh3600 which was slightly more than one dollar a day (Table 5). There was a highly significant  
 101 negative correlation ( $r=-0.904$ ,  $p<0.01$ ) between the incidences of VVF and monthly income.  
 102 The study established that the 34 cases were reported from sub counties with low number of  
 103 maternity units (Table 6). There was a significant negative correlation ( $r= -0.6415$ ,  $df (10)$ ,  
 104  $p<0.05$ ); ( $t= 2.644$ ) between number of maternity units per sub county with the number of the  
 105 VVF cases studied. Almost half of the patients (47%, n=16) perceived that the general public  
 106 sympathized with their fate (Table 7)

107  
 108 **Table 1 Age of the VVF patients in the study**

Age in years	Frequency	Percentage (%)
15 – 17	8	23.5
18 – 20	6	17.6
21 – 23	7	20.6
24 – 26	6	17.6
27 – 29	3	8.8
30 – 32	2	5.9
33 – 35	2	5.9
Total	34	100.0

109  
 110  
 111  
 112  
 113 **Table 2: Distribution of VVF cases by education levels**

Education level	Frequency	Percentage (%)
-----------------	-----------	----------------

Never been to school	10	29.4
Primary school	18	52.9
Secondary school	4	11.8
Post secondary (college)	2	5.9
Totals	34	100.00

114

115 **Table 3: Period the patients had lived with the condition**

Age in years	Frequency	Percentage (%)
<2	10	29.4
2-4	11	32.3
4-6	9	26.5
6<	4	11.8
Total	34	100.0

116

117 **Table 4: Distribution of VVF patients by age at the occurrence of the condition**

118

Age in years	Frequency	Percentage (%)
12 – 15	8	23.5
15 – 18	9	26.5
18 – 21	7	20.6
21 – 24	5	14.8
24 – 27	3	8.8
27– 30	2	5.9
Total	34	100.0

119

120

121

122 **Table 5: Distribution of income levels patients with VVF cases**

Gross income level per month (kshs)	Frequency	Percentage (%)
-------------------------------------	-----------	----------------

Below 3, 000	18	52.9
3, 000 – 6, 000	10	29.4
6, 001 – 9, 000	4	11.8
Over – 9, 000	2	5.9
Total	34	100.00

**Table 6: Comparison of VVF cases in Machakos County against maternity facilities available by**

**Sub-counties**

Sub-county	Number of health facilities with maternity unit		VVF cases	
	Frequency	%	Frequency	%
Central	8	19.0	0	0.0
Kalama	2	4.8	5	14.7
Yatta	4	9.5	5	14.7
Ndithini	2	4.8	3	8
Masinga	2	4.8	5	14.7
Yathui	2	4.8	6	17.6
Katangi	2	4.8	3	8.8
Matungulu	4	9.5	1	2.9
Mwala	4	9.5	4	11.8
Athi river	4	9.5	0	0.0
Kangundo	4	9.5	2	5.9
Kathiani	4	9.5	0	0.0
Totals	42	100	34	100

**Table 7: Perception of VVF patients' perception on societal reaction towards them**

Societal reaction	Frequency (Percentage)
Shunned	9 (26.5%)
Sympathy	16 (47.1%)
Feared	6 (17.6%)
Isolated	3 (8.8%)

Total	34 (100%)
-------	-----------

## Discussion

This study showed that all the VVF incidents were among women whose ages ranged between 15 and 35 years with mean and median years of 22.3 years. These were women at their peak of fertility<sup>(3)</sup> and fecundity hence any condition affecting their reproduction would adversely affect their life. The mean age of the patient at the time they developed the condition was found to be 18.8 years. Though this mean age was within teenage, it differs from studies conducted in Kano and Zaria (Northern Nigeria) whose results showed that the disease was commoner among women who were fifteen years or younger but very rare after the age of twenty-five years.<sup>(3,4)</sup> However the observation that the incidents decreased with age is consistent with the highly significant negative correlation between the age of the patient and number of VVF cases.

Among the factors that lead to development of VVF in developing countries are marriage and conception at young age particularly where the pelvic growth had not been achieved.<sup>(5)</sup> Very young mothers are physically immature to have babies, and often suffer from obstructed labor. The head of the fetus may press against the maternal bones of the pelvis, thereby compressing the tissues of the anterior vaginal wall and the bladder. As a result, these tissues do not get enough blood supply and slough off, resulting in a fistula. The healthcare workers attributed the VVF condition mainly from obstructed labor due to Mal-presentation of the baby at delivery, Cephalopelvic disproportion (CPD), maternal distress or Cervical dystocia<sup>(6)</sup>. This eventually leads to prolonged obstructed labor which is associated with pressure necrosis, edema and tissue sloughing which is responsible for 97% of VVF cases in developing countries.<sup>(7)</sup>

The study observed that more than a quarter of the patients had no formal education and even among the formally educated, less than 20% had post primary education. This observation is consistent with Mohammad<sup>(8)</sup> who reported a strong correlation between lack of education and VVF incidents illiteracy and incidents of VVF

The study observed that the mean length of time the patients had lived with the condition was about 3.5 years. Hancock<sup>(9)</sup>, citing results from a survey done in Uganda explains the reasons behind failure of local surgeons from carrying out fistula repair as perceived view that fistula surgery is difficult, its results are poor, there was no opportunity to learn fistula surgery, lack of special instrument and equipment, and no specialist nursing care. He stresses that though not all fistulae can be repaired, fistula repair does not require special equipment and post operative nursing, though important is not complicated. Stamatakis *et al*<sup>(10)</sup> have explained that the main complication of VVF surgery is recurrent fistula formation but a typical fistula repair would be accomplished through accurate diagnostic evaluation and timely repair using procedures that exploit basic surgical principles and application.

The study showed that the mean daily earning of a majority of the patients was slightly more than one dollar. It also showed a highly significant negative correlation between the incidents of VVF and monthly income. This observation is consistent with Daru *et al*<sup>(11)</sup> who observed that VVF was problem occurring mainly amongst the illiterate and poor farmers who had prolonged obstructed labor<sup>(12)</sup>. It is observed that teenage pregnancies occur in both developed and developing countries. However 97% of obstetric fistulae are observed in developing countries<sup>(13)</sup>. Unlike in the developing world, almost all deliveries in developed world are supervised by skilled healthcare givers. Therefore, the risk of obstetric fistula is low among the mothers who deliver under supervised hands in spite of their age, early marriage or early pregnancy. Poor women are more likely to deliver at home or under unskilled hands, and risk of VVF complication increases if she is underage.

The study established a significant negative correlation between the number of maternity units per sub county and the number of VVF cases studied. Mawajdeh *et al*<sup>(14)</sup> identified five elements for assessing

the quality of prenatal care. They included patient provider relationship, technical management, information exchange, continuity and management. The women reported to be satisfied with the quality of prenatal care when these requirements were adequately met. The adequacy of health facilities motivates the women to seek for prenatal healthcare. Whenever the perceived needs are not adequately met, the tendency is to search for suitable alternatives. This is observed in the study area where 70% of deliveries occur at home, usually attended by a relative or a Traditional Birth Attendant. This choice has such consequences of severe adverse effects to both mother and child, among the outcomes are VVF<sup>(15)</sup>

Almost half of the patients (47%, n=16) perceived that the general public sympathized with their fate but others felt isolated or shunned by their family and society. Besides physical challenges, VVF is associated with medical and psychosocial complications since a mother at young age becomes a victim who cannot bear children that often times determines her value in the society. Abandonment, isolation and stigmatization were observed by Nseno<sup>(16)</sup> as the major independent variables in determining coping strategies. The more abandoned and stigmatized the woman was, the less active she was in coping, and for some patients they never regain their societal value even after surgical repair.

## Conclusion

Vesicovaginal Fistula can be prevented through adequate antenatal care that should be extended to at risk mothers who are likely to develop obstructed labor. A cesarean section should be planned for those who need it. In case of a long-standing obstructed labor, the urinary bladder should be drained continuously for a period of 5-7 days following the delivery of the baby. The ideal time for surgery of vesicovaginal fistula due to obstructed labor is after three months following delivery. By this time, the general condition of the woman improves and the local tissues are likely to be free from infection. The repair is done through either the vagina or the abdomen, or through a combined approach depending upon the choice and expertise of surgeon. Community empowerment would reduce early marriages and consequently reduce VVF cases. Timely repair of VVF cases would considerably reduce stigmatization and other challenges associated with the condition.

No funding was obtained to prepare this manuscript. The authors state they had no competing or conflict of interest

## Consent Disclaimer:

As per international standard or university standard written patients' consent and ethical approval have been collected and preserved by the author(s).

## References

1. Kimberly, L. and Alison C ( 2011 ) Advanced Reproductive Age and Fertility SOGC Clinical practice guideline Toronto, Ontario Canada
2. Bimbola, and Kemi O (2013) Obstructed labour: the main cause of vesico-vaginal fistula–Review of literature, European Journal of Research in Medical Sciences Vol. 1 No. 1, September 2013
3. Kabir, M.; Iliyasu, Z.; Abubakar, I. S. and Umar U. I, Medico-social problems of patients with vesicovaginal fistula in Murtala Mohammed Specialist Hospital, Kano *Annals of African Medicine* Vol. 2, No. 2; 2003: 54 – 57
4. Harrison K.A. Childbearing, health and social priorities: a survey of 22,774 consecutive hospital births in Zaria, northern Nigeria. *Br J Obstet Gynaecol* (Suppl) 1985; 5: 91-97.
5. Spurlock , J (2016) Vesicovaginal Fistula [emedicine.medscape.com](http://emedicine.medscape.com)

- 234 6. Korell A.N prolonged obstructed labor causing a severe obstetric fistula: a case report, J Reprod  
235 Med  
236
- 237 7. Thevenet, A. and Tita, N (2012) When is Primary Caesarean Appropriate: A Maternal and  
238 obstetrical Indications. Seminars in Perinatology, Elsevier  
239
- 240 8. Mohammad, R. H (Vesicovaginal Fistula (VVF). A problem of underdevelopment, Foundation  
241 of Women's Health Research and Development- FORWARD – Nigeria  
242
- 243 9. Hancock, B (2005) First Steps in Vesico-Vaginal Fistula Repair The Royal Society of Medicine  
244 Press Limited  
245
- 246 10. Stamatakos. M, Sargedj .C, Stasinou. T, Kontzoglou, K (2014) Vesicovaginal Fistula: Diagnosis  
247 and Management, Indian journal of Surgery  
248
- 249 11. Daru, P.H.; Karshima J.A .; Mikah, S .; Nyango, D (2011). The Burden of Vesico-Vaginal  
250 Fistula in North Central Nigeria , Journal of West Africa College of surgeons  
251
- 252 12. 11 Prevention of maternal mortality network. Barriers to treatment of obstetric emergencies in  
253 rural communities of West Africa. Stud Fam Plann 1992; 23: 279–291.  
254
- 255 13. Audu, B.M.; Kulima, A.A., and Bako, B (2008) Epidemiology ofof Vesico Vaginal Fistula: no  
256 longer a calamity of teenagers, Journal of Obstetrics and Gynecology, May, 2008  
257
- 258 14. Mawajdeh,S.; Al-Qutob, R., and Raad, F.B(2013) the assessment of Quality of Care in  
259 PrenatalServices in Irbid, Jordan. The Female Client and Health-Care Provider, International  
260 Development Reseach Centre  
261
- 262 15. Singh, A (2013 ) Mothers Choosing At Home Birth Risk Severe Complications, Higher  
263 Likelihood of Death Pub Med  
264
- 265 16. Nsemo, A. D (2014) Influence of Abandonment, Stigmatization and Social Isolation on coping  
266 Strategies of women with Vesico Vaginal Fistula in AkwaIbom State, Nigeria. Journal of Nursing  
267 and care. ISSN: 2167-1168  
268  
269  
270