

Original Research Article

Acute and Chronic Effects of methanolic extract of *Teucrium polium* on Blood Parameters and histopathology of liver and kidney in male and female mice

Abstract:

Teucrium polium L. is commonly used as medicinal plants in Algeria against a variety of human disease. In this study, the extraction of aerial parts of *T. polium* with methanol gave a dry matter yield 8.24 %. The effects of methanolic extract were examined *per os* on female rats *Albino Wistar* for six weeks. Biochemical and haematological serum parameters, and morphology and histopathology of organs of treated rats were studied. The study of acute toxicity showed a low toxicity with $LD_{50} > 2400$ mg / Kg of body weights. These data can be used to classify these plant in slightly toxic. However, the sub-acute treatment for six weeks of rats with 75, 150 and 300 mg of methanolic extract / Kg of body weight resulted in significant increases of the most studied haematological parameters. Biochemical analysis revealed significant increase of renal parameters (urea, creatinine, uric acid, Na and K), accompanied by increase of relative weight of kidney, lipidic (cholesterol) and hepatic glutamate oxalo-acetate transaminase (GOT) values in all treated rats. Histopathological examination confirmed the biochemical tests by the observation of perilobular necrosis areas, bile duct and inflammatory infiltration of the liver and presence of marked intracytoplasmic vacuoles in kidney with the dose 300 mg of methanolic extract of *T. polium* / Kg of animal body weight. Use of *Teucrium polium* L. may cause hepatotoxicity and/or nephrotoxicity after prolonged herb administration.

Key words: *Teucrium polium* L, methanolic extract, LD_{50} , biochemical parameters, Acute and chronic toxicity, Liver, Kidney.

25 **Introduction**

26 Mountain germander, *Teucrium polium* L. (lamiaceae) is originating from the south-west of
27 Asia, Europa and North Africa (Abdollahi et al., 2003). There are various reports about
28 pharmacological properties of *Teucrium polium* L. These include antibacterial (Autore *et al.*,
29 1984; Aggelis et al., 1998), anti-inflammatory, analgesic, antispasmodic and hypolipidemic
30 (Abdollahi et al., 2003; Dehghani et al., 2005) anti-ulcer, anti-nociceptive (Kaileh et al.,
31 2007), antidiabetic, diuretic (Gharaibeh *et al.*, 1988), antifongic, inotropic and chronotropic
32 (Niazmand et al., 2008) effects. Most of these effects seem been attributed to the antioxidant
33 and free radical scavenging properties of *Teucrium polium*.

34 Phytochemical investigations have shown that *T. polium* contains various compounds such as
35 flavonoids, polyphenols (Proestos et al., 2004), iridoides, tannins, essential oils and alkaloids
36 diterpenoids, principally furano neoclerodanes (Shakhanbeh and Atrousse, 2000; Parsaee et
37 Shafiee-Nick, 2006). One of these major components is teucrine A (Bruneton, 1999;
38 Ramnathan et al., 2005).

39 However, many herbal medicinal plants including *T. polium* were found to induce fatal
40 hepatic effects and severe acute liver failure with marked haematological and biochemical
41 alterations after prolonged administration (Khleifat et al., 2000). Several cases of germander
42 hepatitis were reported in Canada (De Smet 1997) and Spain (Al Varez 2001). Several reports
43 linked the consumption of *T. polium* with hepatitis in man (Mattie 1995, Mazokopakis 2004,
44 Starakis 2006). *T. polium* is consumed by many people in Mediterranean countries such as
45 Jordanians and Algerians for the treatment of several deaseases, and there is no detailed
46 information on the liver status after the consumption of the plant tea. Cytotoxic effects of
47 *Teucrium polium* on some cell lines (A549, BT20, MCF7, and PC12) has been reported by
48 Nematollahi-Mahani (2007). The IC₅₀ values for each cell line were found: A549, 90 µg/mL;

BT20, 106 µg/mL; MCF-7, 140 µg/mL; and PC12, 120 µg/mL *T. polium* ethanolic extract. All studies has been focused on aqueous or infusion (Zal et al., 2001; Rasekh et al., 2005), ethanolic (Al-Ashban, 2005; Nematollahi-Mahani et al., 2007) and ethyl acetate extracts and no toxicological study has reported the effects of methanolic extract which is rich of flavonoids and polyphenols. Finally, it is well known that every drug has been associated with hepatotoxicity almost certainly due to the pivotal role of the liver in drug metabolism. Hepatic metabolism is, first and foremost, a mechanism that converts drugs and other compounds into products that are more easily excreted and that usually have a lower pharmacologic activity than the portent compound (Poppers, 1980; Tolman, 1998). A metabolite may have higher activity and/or greater toxicity than the original drug. Metabolites of the drugs that are excreted from kidneys may also cause cellular damage leading to kidney dysfunction (Singhal *et al.*, 1998).

In this report, the focus was on the acute and chronic toxicity tests. The primary concern was to determine how toxic the methanolic extract of *T. polium* may be after acute administration to rats. Secondary, what would be the target organ for that toxicity and whether there would be any correlation between the toxic effects and the flavonoids and polyphénols contained in the plant materials after chronic oral administration to rats.

Materials and Methods

Plant material

The aerial parts of *Teucrium polium* were collected during June 2009 from Bougaa region at the north of Setif province in the north east of Algeria, identified by Prof Laouar and a voucher specimen was deposited at the Applied Biochemistry Laboratory, University Ferhat

Abbas, Setif, Algeria. The plant materials were dried at room temperature and powdered. The dry plant samples were extracted with absolute methanol. The dry extract was obtained after removing the solvent by evaporation under reduced pressure at 45° C. The extract was stored at – 20° C until use.

Animals

Experiments were performed on adult female Wistar albino rats, weighting 201.61 ± 7.04 g. The animals obtained from ‘Institut Pasteur d’Algérie’ were housed in groups of eight to ten in plastic cages at controlled room temperature. Water and food were freely available.

Determination of total polyphenol and flavonoid contents

Total polyphenols were measured using Folin-ciocalteu according to Li et al. (2007). the absorbance is measured at 765 nm. The concentration of total polyphenols is calculated starting from the calibration line established with the gallic acid and is expressed in μg of equivalent of acid gallic per milligram of extract.

Flavonoids were quantified using AlCl_3 reagent (Bahorun *et al.*, 1996). Flavonoids were measured as quercetin equivalents. One mL of *T. polium* methanolic extract (TPME) samples was dissolved in methanol, 1 mL of AlCl_3 (2% in methanol) was added, and after incubation for 10 min, the absorbance was measured at 430 nm (Boumerfeg *et al.*, 2009).

Acute toxicity assay

In order to study any possible toxic effect or changes in normal behavior, 5 groups of 7 rats were used in this experiment. The animals were fasted 24 hours before the treatment (Ulanova, 1975; Frank, 1992). The acute toxicity of the plant was studied by preparing four different concentrations of the extract (0.3, 0.6, 1.2 and 2.4 g/kg), and administered orally to four

groups of animals. The fifth group was taken as a control and given 1.0 ml NaCl 9‰. The behavioural changes, posture and mortality were checked for 24 hours. The method of Karber (Stowtchiva, 1988) was employed for the determination of acute oral lethal dose (LD₅₀).

Chronic toxicity study

Animals were divided into four dose groups of 8 animals /dose. The first group was given 1 ml normal saline and taken as a control. The second, third and forth group were given a single doses of 75, 150 and 300 mg / Kg of TPME extract by gavage daily. Body weight food consumption and clinical observations were monitored daily. Animals were fasted 3 h prior to dosing to facilitate administration of the complete dose. All animals treated for 42 days then, they were fasted for about 3 h and sacrificed by euthanasia. Immediately after decapitation blood samples were obtained directly from the neck for haematological and serum analysis. By thoracic abdominal longitudinal incision the animal abdomen was opened and the liver and left kidney were removed and the wet weights were recorded (Oduola *et al.*, 2007).

Haematological and biochemical analysis

Under ether anaesthesia all the rats were euthanized, blood samples (2.0-4.0 ml) were withdrawn by sinus retro-orbital puncture in tubes containing EDTA and immediately processed for haematological tests using Beckman coulter-automatic haematology analyzer (USA). The haematological parameters measured were mean cell volume (MCV), red blood cells (RBC), white blood cells (WBC), hematocrit (HCT), platelets (PLT), mean platelet volume (MPV), hemoglobin (HGB), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC). For biochemical analyses, 2 to 3 ml of blood were collected in a heparinized tube and centrifuged at 4000 g/ 5 min. at 4°C. The plasma obtained was stored at -20° C until use. The biochemical parameters including glucose (Glu), urea (Urea), creatinin (Creat), uric acid (UA), Na, K, cholesterol (Chol), triglycerides (TG),

glutamate oxalo-acetate transaminases (GOT), glutamate pyruvate transaminases (GPT), alkaline phosphatase (ALP) were measured at the Central Laboratory of the University Hospital (CHU) of Setif.

Evaluation of organs

The animals were weighted and euthanized by ether inhalation, all the organs/tissues were carefully examined macroscopically and the brain, lungs, heart, spleen, liver, kidneys and ovaries were weighed. The specimens were fixed in 10% formalin for 24 h, and standard dehydration and paraffin-wax embedding procedures were used. Sections (5 μ m) were cut in a microtome, adhered to glass slides with. Hematoxylin and eosin-stained slides were prepared by using standard methods and evaluated by light microscopy.

Statistical analysis

Statistical analysis was performed using Student's *t*-test for significance and analysis of variance (ANOVA) followed by Dunnett's test were done for the multiple comparison of the effect of different extract doses. Values of $p < 0.05$ were considered statistically significant. The comparison of the averages and the variances was done using SigmaStat 3.5, version 3, and SigmaPlot 10.0.

Results

The methanolic extraction gave a yield of 8.24 % from the aerial parts of *T. polium*. The total polyphenols, flavonoids quantities were 18.63 ± 3.51 mg equivalent gallic acid / g dried material and 9.14 ± 2.39 mg equivalent quercetin / g dried material, respectively.

Acute toxicity studies

Rats were individually observed during the first 30 min and regularly during the first 24 h after TPME administration. Clinical signs observed were summarized in table 1. Mortality due to different doses administered to female rat was 10 % and 20 % for 1200 and 2400 mg/Kg, respectively. LD₅₀ is higher than 2400 mg/kg body weight.

Table 1: Signs and symptoms of *T. polium* toxicity on female rat score based on the order of severity.

Extact	Dose (mg/kg)	Signs and symptoms	Time
<i>T. polium</i>	0	Normal	-
	300	piloerection, stressed rats	15 min.
	600	Piloerection, Laboured breathing, immobilization of the rats.	15 min.
	1200	Irritability, Tremblement, Labored breathing , immobilization of the rats	6 days
	2400	Paralysis, Tremor , Labored breathing, Death	30 min.

Effects of *T. polium* on body weight

Percentages of changes in body weights during the administration period are shown in Figure 1. Values for the group treated with 75 mg of TPME /kg of body weight were slowly decreased compred to those of the controls group but differencies were not significant. Whereas, administration of 150 and 300 mg/kg result in significant decrease except the fifth week with 300 mg/ Kg. It seems that, at the concentration of 300 mg/kg, there is certain adaptation of the anomals wich is not seen the the other concentrations.




Figure 1: Changes of body weight growth (%) of treated rats and control.

The macroscopic analysis of the target organs of the treated animals (liver, lung, heart, brain and spleen) did not show significant changes in color and texture when compared with the control group. Nevertheless, a significant increase in the values of the relative mass of kidneys with the 150 and 300 mg/kg are observed. The results of organs weight are summarized in figure 2.




Figure 3: Effects of methanolic extract of *T. polium* on organ relative weights of female rats in chronic toxicity. **A**; ovaries, **B**; kidneys, **C**; brain, **D**; liver, **E**; lungs and **F**; heart. The values are mean \pm SEM (n = 7-8). * p < 0.05, ** p < 0.01.

Effets of TPME on haematological and biochemical paramaters

Haematological parameters of the blood in the four groups of rats are represented in **Table 3**. A significant increase was observed of the following parameters: RBC, MPV, HCT, PLT and HGB compared to the none treated rats. Similarly, biochemical parameters in the four groups of rats, Glu, Creat, K, Urea, Na, UA and GOT were significantly increased after chronic treatment with 75, 150 and 300 mg / kg and a significant increase in GPT level with the dose of 75 mg/kg and in Chol level with the 300 mg / kg *T. polium* extract compared to controls.

Table 3: haematological data for female rats orally treated by *T. polium* extract for 6 weeks

Haematologic tests	Non-treated	75 mg/kg	150 mg/kg	300 mg/kg
RBC ($10^6/\text{mm}^3$)	5.41 ± 0.29	$7.65 \pm 1.03^{**}$	$6.36 \pm 0.16^*$	$7.41 \pm 0.32^{**}$
HCT (%)	29 ± 1.83	$39.06 \pm 0.32^{**}$	$38.96 \pm 0.71^{**}$	$36.52 \pm 2.17^*$
PLT ($10^3/\text{mm}$)	303 ± 27.76	$548 \pm 19.78^{**}$	$532.29 \pm 20.93^{**}$	$638.60 \pm 34.42^{**}$
MPV (μm^3)	6.46 ± 0.07	$7.04 \pm 0.05^{**}$	$6.71 \pm 0.09^*$	$7.12 \pm 0.11^{**}$
HGB (gr/dl)	9.90 ± 0.59	$14.23 \pm 0.08^{**}$	$13.93 \pm 0.25^{**}$	$13.62 \pm 0.61^{**}$
WBC ($10^3/\text{mm}^3$)	7.46 ± 0.82	5.80 ± 0.53	6.76 ± 0.55	6.48 ± 0.43
MCV (μm^3)	50.84 ± 0.67	50.40 ± 0.50	49.29 ± 0.47	49.08 ± 1.11
MCH (gp)	17.81 ± 0.23	18.35 ± 0.13	17.42 ± 0.26	17.63 ± 0.37
MCHC (g/dl)	35.35 ± 0.41	36.32 ± 0.19	35.53 ± 0.23	35.93 ± 0.41

* P<0.05, **P<0.01.

Table 4: Serum biochemical data for female rats orally treated by *T. polium* extract for 6 weeks


Biochemical data	Non-treated	75 mg/kg	150 mg/kg	300 mg/kg
Na (mEq/L)	134.57 ± 2.10	169.07 ± 4.74**	170.13 ± 1.32**	165.55 ± 1.81**
UA (mg/ L)	8 ± 1.21	27.14 ± 1.97**	17.13 ± 1.29*	16.53 ± 2.90*
Urea (gr/ L)	0.51 ± 0.03	0.64 ± 0.05	0.65 ± 0.03	0.61 ± 0.06
Chol (gr/ L)	0.34 ± 0.08	0.33 ± 0.026	0.35 ± 0.03	0.57 ± 0.041**
TG (gr/ L)	0.47 ± 0.084	0.39 ± 0.02	0.49 ± 0.03	0.39 ± 0.04
ALP (UI/L)	93.80 ± 7.07	74.33 ± 2.67	97.83 ± 3.31	86.30 ± 41.98
GOT (UI/L)	43.86 ± 3.77	96.67 ± 7.84**	85.17 ± 4.39**	85.00 ± 16.16**
GPT (UI/L)	96.67 ± 8.66	35.57 ± 3.09**	59.43 ± 5.22**	56.00 ± 16.76**
Creat (mg/ L)	6.58 ± 0.37	6.15 ± 0.27	6.79 ± 0.36	6.0 ± 0.45
Glu (gr/ L)	1.47 ± 0.11	1.27 ± 0.11	1.30 ± 0.06	1.24 ± 0.05

Values are expressed as mean (n = 8) ± SED, * p ≤ 0.05, ** p ≤ 0.01.


Histopathological examination

The observation of the histological slices of the liver and kidneys of treated rats compared to controls are presented in figures 3 and 4, respectively. The examination of kidneys revealed presence of intracytoplasmic vacuoles, precisely on cortical area when the dose of 300 mg/kg of *T. polium extract* was used. Histological examination of the liver showed moderate portal inflammatory infiltrates, a vascular congestion around vessels. Mild lobular necrosis, a vascular congestion and steatosis was also seen of the cuts of the rats treated with group treated by 75 mg/kg *T. polium* extract. A perilobular necrosis and an inflammatory infiltrate around the portal vein of the cuts of the rats treated with the second group. A moderate

186 inflammatory infiltrates in portal tracts were seen, proliferation of bile ducts and a portal
187 fibrosis was noted on 300 mg/kg of *T. polium* extract.



188
189 **Figure 4:** Renal Histological cuts of control group and treated rats with 300 mg/kg *T. polium*
190 extract. (hematoxyline, eosin; \tilde{A} —600). CA: cortical area, RT: renal tube, VIC: vacuoles
191 intracytoplasmic.



193
194
195 **Figure 4:** hepatic histological cuts of control group and rats treated with 75 mg/kg, 150
196 mg/kg and 300 mg/kg *T. polium* extract. (hematoxyline / eosin). **H** : hepatic cell, **CLV** :
197 centrolobular vein, **IF** : Inflammatory infiltrate, **PLN** : perilobular necrosis, **EP** : espace porte,
198 **DP** : duct proliferation, **FEP** : Fibrosis in bridge.

Discussion:

Acute toxicity

No sign of acute toxicity for female rat treated by *T. polium* extract was seen; this suggests that LD₅₀ is higher than 2400 mg/Kg of body weight. The results is in agreement with the study of Autore *et al.* (1984) and Rasekh *et al.* (2005). Hasani *et al.* (2007) dismount that the administration of this plant is influenced by the nature of the sex. However, Khleifat et al. (2001) determined LD₅₀ of 262.0 mg/kg in the rats treated by intraperitoneal way by ethanolic extract of *T. polium*. This data is correlated with the route of adopted administration which targets directly the liver (Lüllmann *et al.*, 1998; Sharp and La Regina; 1998).

Chronic toxicity

Effect on body weight

The rats present a moderately significant reduction (3%) of their weights during the sixth week with the 75 mg/kg of methanolic extract. Indeed, the loss of rat's weight is correlated in a physiological state of the animal and can be explained by a reduction of consumption of food. This result agrees with those published by Rasekh *et al.* (2005) where female rats show a significant reduction of body weight ($p < 0.05$) with dose of 100 mg/kg of aqueous extract of *T. polium*. The other groups show a normal evolution of their body weight as shows compared to the control during all the experimentation duration as Shahraki shows it and others (2005).

Haematological biochemical and parameters of rats

According to the bibliography (Khleifat *et al.*, 2001; Rasekh *et al.*, 2005), none disturbance of hematologic values was reported in rats treated by the aqueous and/or ethanolic extract of *T. polium*. Conversely the methanolic extract induced an increase in the number of red globules cells observed with the three groups treated (75, 150, 300 mg/kg), implying a potentiality of erythropoiesis. However, these data are associated with hematocrit and haemoglobin increases with doses of 75 mg/kg and 300 mg/kg. The significant increase in hemoglobin is associated with a significant increase of MCV and MCH witch indicate a tendency to macrocytose and hypochromy (Ndoutamia and Ganda, 2005). Platelets level increase on 75 and 300 mg/kg doses can be commented on by secondary hyperplaquettose associated an attack of spleen, a martial deficiency and with the myéloprolifératifs syndromes (Bain and Gupta; 2003).

The methanolic extract of *T. polium* shows a significant reduction in glucose with the dose of 75 mg/kg, this agrees with works of Gharaibeh *et al.* (1988) ; Esmaeili and Yazdanparast, (2004) ; Ghoraishian, (2006) ; Bhonde *et al.*, (2007). Several flavonoids such as quercetin and different terpenoids discovered in *T. polium* decrease the serum glucose level only in the diabetic rats. It is, therefore, possible that these effects of the areal parts of *T. polium can* be due to the flavonoids and/or terpenoids. Zal and their collaborators (2001) accorded the hypoglycemic effects of aqueous *T. polium* extract (area of Kerman, Iran) to their composition in ions: potassium, Zinc, Cadmium and Chromium .This suggests that the hypoglycemic property of this plant depends on the type of ground and the geographical area of harvest. Richness of *T. polium* in flavonoidss and polyphenolic compounds such as cirsimaritin, apigenin-7-glucoside, vicienin and luteolin-7-glucoside gifted of antioxydant activities in particular with the dose of 50 and 100 mg/kg (Ardestani and Yazdanparast, 2007; Sharififar *et al.*, 2008), thus explain the use of *T. polium* in folk medicine in the treatment of the diabetes (Hasani *et al.*, 2005). The Chol level was appreciably increased after chronic

treatment with the dose of 300 mg/kg *T. polium* extract. This is in agreement with the observations at the human ones (Khleifat *et al.*, 2001; Mazokopakis *et al.*, 2005).

The renal parameters showed a very significant in level of urea, creat Na and K. Urea increases could be explained by an increase in degradation of protein compounds, but also by an injure of renal function (Lüllmann *et al.*, 1998). Really, kidneys were clearly damaged and its histological aspect indicated a remarkable cytoplasmic vacuolization of tubular cells of cortical area and which explains the increased in relative weight of the kidney (mainly with the doses of 150 and 300 mg/kg), while the other values of the various studied organs are normal. These results agree with works of Rasekh *et al.* (2005), Khleifat *et al.* (2001) and Iriadam *et al.* (2006) who showed a significant increase in urea and creatinin level in diabetic rabbits after treatment by aqueous extract of *T. polium* (82 mg/kg *per os*), confirmed by an apoptose in some renal sections.

The values of hepatic analysis presented a significant higher in TGO with the three groups and a reduction in TGP in the first group. This reduction does not agree with work of Mattéi *et al.* (1995) and of Rasekh *et al.* (2005). However, results of Rasekh and collaborators (2005) showed that its high level only with the dose of 300 mg/kg after 44 days of treatment.

Mazokopakis and collaborators (2005) reported that a dysfunction of the liver appears after one month of treatment by an infusion of *T. polium*. In parallel, Mattéi *et al.* (1995) defer two cases of lethal hepatitis due to the treatment containing germander. Histological examinations of liver are the same ones observed in this our study in particular in treated rats with 150 and 300 mg/kg doses. These results are also supported by preceding work of Rasekh *et al.*, (2005) an acute and serious failure liver at a man after prolonged administration of *T. polium*, *T. chamaedrys* and *T. capitatum*.

Although the mechanism of hepatotoxicity of *T. polium* is not well elucidated, teucrine A and several diterpenoids neoclerodans, present in aeral parts, were to suspect like hepatotoxic precursors of this plant (Rasekh *et al.*, 2005). Experiments in mice showed a formation of toxic metabolites starting from these diterpenoids which interact with cytochrome P450 3A and the inactivation of glutathione. In fact, the detoxified diterpenoids are effective inductors of hepatocytes apoptosis (Stickel *et al.*, 2005; Stickel *et al.*, 2006). That suggests liver necrosis it is attenuated by a dual mechanism, a direct toxicity and a series of secondary immune reactions (Berardinis *et al.*, 2000). The direct cytotoxicity is known for being the fundamental cause of damage of liver in certain cases, while in others the immunological mechanisms or even a mixture of cytotoxicity and immunogenicity can be implied (Ingwale *et al.*, 2009). A covalent bond of the epoxy hydrolase will take place on the external surface of human hepatocytes and in the presence of teucrine A, could start immune reactions and induces a formation of autoantibody leading the cells to the apoptosis (Fig.).




Figure 6: Toxic mechanism of furano-diterpenoids extracts from *Teucrium polium* (Stickel *et al.*, 2005)

In conclusion, methanolic *Teucrium polium* extract perturb biochemical serum and hematological parameters related to hepatic and renal function after prolonged administration. The liver histology can indicate image of an acute or chronic hepatitis, cases of cholestasis, cytoplasmic vacuoles characteristic of renal tubes. People should consider about the use *T. polium*, particularly if they are not officially informed of their possible unfavorable reactions.

References

- Abdollahi , A., Karimpour, H., Monsef-Esfehani H. (2003). Antinociceptive effects of *Teucrium polium* L. total extract and essential oil in mouse writhing test. *Pharmacol. Res.* 48:31-35.
- Aggelis, G., Athanassopoulos, N., Paliogianni, A., Komaitis, M. (1998). Effect of *Teucrium polium* L. extract on the growth and fatty acid composition of *Saccharomyces cerevisiae* and *Yarrowia lipolytica*. *Antonie van Leeuwenhoek* . 73: 195–198.
- Ardestani, A., Yazdanparast, R. (2007). Inhibitory effects of ethyl acetate extract of *Teucrium polium* on in vitro protein glycoxidation. *Food Chem.l Toxicol.* 45: 2402-2411.
- Autore, G., Capasso, F., De Fusco, R., Fasulo, M.P., Lembo, M. ,Mascolo N., Menghini A. (1984). Antipyretic and antibacterial actions of *Teucrium polium* (L.) *Pharmacol. Res. Commun.* 1:16.
- Bahorun, T., Gressier, B., Trotin, F., Brunete, C., Dine, T., Vasseur, J., Gazin, J.C., Pinkas, M., Luycky, M., Gazin, M. (1996). Oxygen species scavenging activity of phenolic extracts from hawthorn fresh plant organs and pharmaceutical preparations. *Arzneimittel-Forsch.* 46: 1086-1089.
- Bain Barbara, J., Gupta R. (2003). A–Z of Haematology. 1^{ere} edition. Blackwell Publishing Ltd Library of Congress Cataloging-in-Publication Data. pp. 196-226.
- Beradinis, DV., Moulis, C., Maurice, M., Beaune, P., Pessayre, D., Pompon, D., Loeper, J. (2000). Human microsomal epoxide hydrolase is the target of germander-induced autoantibodies on the surface of human hepatocytes. *Mol. Pharmacol.* 58(3):542-551.
- Bhonde, R., Shukla, R.C., Kanitkar, M., Shukla, R., Banerjee, M., Datar, D. (2007). Isolated islets in diabetes research. *Indian J Med. Res.* 125:425-440.
- Boizot, N., Charpentier, J.P. (2006). Méthode rapide d'évaluation du contenu en composés phénoliques des organes d'un arbre forestier .*Le Cahier des Techniques de l'Inra.* pp. 79-82.

- Boumerfeg, S., Baghiani, A., Messaoudi, D., Khennouf, S., Arrar, L. (2009). Antioxidant properties and xanthine oxidase inhibitory effects of *Tamus communis* L. root extracts .*Phytother. Res.* 23:283-288.
- Bruneton, J. (1999). Pharmacognosie : phytochimie, plantes médicinales .3 eme édition. Paris. pp. 648-650.
- Dehghani, F., Khozani, T.T., Panjehshahin, M.R., Karbalaedoost, S. (2005). Effect of *Teucrium polium* on histology and histochemistry in rat stomach. *Indian J Gastroenterol.* 24:126-127.
- Djeridane, A., Yousfi, M., Nadjemi, b., Boutassouna, A., Stocker, c., Vidal, N. (2005). Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chem.* 97: 654–660.
- Esmaeili, M.A., Yazdanparast, R. (2004). Hypoglycaemic effect of *Teucrium polium*: studies with rat pancreatic islets. *J Ethnopharmacol.* 95:27-30.
- Frank, C.L.U. (1992). Toxicologie, Données générales procédures d'évaluation, organes cibles, évaluation du risque. Paris. pp: 73- 202.
- Gharaibeh, M., Hhamzeh, h., Salhab, A.S. (1988). Hypoglycemic effects of *Teucrium polium* *Journal of Ethnopharmacology.*2: 93-99.
- Ghoraishian, S.M. (2006). The effect of hazel-leaf decoction on blood glucose reduction in the diabetic rats. *World Journal of Medical Sciences.* 1(2):144-146.
- Hasani, P., Yasa, N., Vosough-Ghanbari, S., Mohammadira, A., Dehghan, G., Abdollahi, M. (2007). In vivo antioxidant potential of *Teucrium polium*, as compared to α -tocopherol. *Acta Pharm.* 57:123–129.
- Ingwale, D., Kshirsagar A., Ashok, P., Vyawahare, N. (2009). Role of atioxidat in the maagnet of hepatic complications. *Pharmacologyonline* 1: 238-253.
- Iriadam, M., Davut, M., Haice, G., Baba F.S. (2005). Effects of two Turkish medicinal plants *Artemisia herba-alba* and *Teucrium polium* on blood glucose levels and other biochemical parameters in rabbits. *J. Cell. Molecul. Biol.* 5:19-24.
- Kadifkova Panovska, T., Svetlana K., Stefova, M. (2005). *In vitro* antioxidant activity of some *Teucrium* species (*Lamiaceae*). *Acta Pharm.* 55:207-214.
- Kaileh, M., Berghe, W.V., Boone, E., Essawi, T., Haegeman, G. (2007). Screening of indigenous Palestinian medicinal plants for potential anti-inflammatory and cytotoxic activity. *J. Ethnopharmacol.* 113:510-516.
- Khleifat, K., Shakhanbeh, J.; Tarawneh, K. (2001). The chronic effects of *Teucrium polium* on some blood parameters and histopathology of liver and kidney in the rat. *Turk J Biol* 26 : 65-71.

- Li, H.B., Cheng, K.W., Wong, C.C., Fan, K.W., Chen, F., Jiang, Y. (2007). Evaluation of antioxidant capacity and total phenolic content of different fractions of selected microalgae. *Food chem.* 102: 771-776.
- Ljubuncic, P., Dakwar, S., Portnaya, I., Cogan, U., Azaizeh, H., Bomzon, A. (2006). Aqueous extracts of *Teucrium polium* possess remarkable antioxidant activity in vitro *Advance Access Publication.* 3(3) :329-338.
- Lüllmann, H., Mohr, K., Ziegler, A. (1998). Atlas de poche de pharmacologie. 2eme édition. Médecine-Sciences: Flammarion .Paris. pp.32-42.
- Mattéi, A., Rucay, P., Samuel, D., Feray, C., Reynes, M., Bismuth, H. (1995). Liver transplantation for severe acute liver failure after herbal medicine (*Teucrium polium*) administration. Hepato-biliary Center, South Paris University, Paul Brousse Hospital, 14 Avenue Paul Vaillant Couturier, 94800 Villejuif, France. Correspondence.
- Mazokopakis, E., Lazaridou, S., Tzardi, M., Mixaki, J. Diamantis, I. Ganotakis, E.(2004). Acute cholestatic hepatitis caused by *Teucrium polium* L. *Phytomed.* 11: 83-84.
- Ndoutamia, G., Ganda, K. (2005). Détermination des paramètres hématologiques et biochimiques des petits ruminants du Tchad. *Revue Méd. Vét.*, 156 : 202-206.
- Niazmand, S., Erfanian Ahmadpoor, M., Moosavian, M., Derakhshan M. (2008). The positive inotropic and chronotropic effects of *Teucrium Polium* L. Extract on Guinea Pig Isolated Heart .*Pharmacologyonline.* 2: 588-594.
- Parsaee, H., Shafiee-Nick, R. (2006). Anti-Spasmodic and anti-Nociceptive effects of *Teucrium polium* aqueous extract. *Iranian biomedical journal.*10(3) :145-149.
- Ramnathan, S.P., Slavoff S.A., Grundel E., White K.D., Mazzola, E., Koblenz, D., Rader J. (2005). Isolation and characterisation of Selected Germander diterpenoids from authenticated *Teucrium chamaedrys* and *T. canadense* by HPLC, HPLC-MS and NMR . *Phytochem. Anal.* 17:243-250.
- Rasekh, H.R, Khoshnood-Mansourkhani, M.J., Kamalinejad, M. (2001). Hypolipidemic effects of *Teucrium polium* in rats. *Fitoterapia.* 72:937-939.
- Rasekh, H.R., Yazdanpanah, H., Hosseinzadeh, L., Bazmohammadi, N., Kamalinejad, M .(2005). Acute and subchronic toxicity of *Teucrium polium* total extract in rats .*Iranian Journal of Pharmaceutical Research.* 4: 245-249.
- Shahraki, M.R., Arab, M.R., Mirimokaddam, E., Palan, M.J. (2007).The effect of *Teucrium polium* (Calpoureh) on liver function, serum lipids and glucose in diabetic male rats. *Iranian Biomedical Journal.* 11(1): 65-68.
- Shakhanbeh, J., Atrouce, O. (2001). *Teucrium polium* inhibits nerve conduction and carrageenan induced inflammation in the rat skin.*Turk J Med Sci.* 3:15-21.
- Sharififar, F., Dehghn-Nudeh, G., Mirtajaldini, M. (2008). Major flavonoids with antioxidant activity from *Teucrium polium* L. *Food Chemistry.*

Sharp Parrick, E., La Regina Maria, C. (1998). The laboratory rat. CRC press LLC. USA. pp. 120-152.

Stowchiva, S. (1988). Guide Manuel de pharmacologie et toxicologie. Sofia: Bulgaria Editeur: Medicinal et phys-cultura .pp. 15.

Stickel, F, Gerlinde, E., Karl, S.H. (2000). Hepatotoxicity of botanicals. *Public Health Nutrition*. 3(2):113-124.

Stickel, F., Patsenker, E., Schuppan, D. (2005). Herbal hepatotoxicity. *J. Hepatol*. 43:901-910.

Stickel F., Schuppan D. (2006). Herbal medicine in the treatment of liver diseases. *Digestive and Liver Disease*. 39: 293–304.

Ulanova. I.P. (1975). Toxicometry and prophylactic toxicology. Institute of industrial hygiene and occupational diseases, Moscow, USSR. Papers. 44-55.

Zal, F., Vasei, M., Rasti, M., Vessal, M. (2001). Hepatotoxicity associated with hypoglycemic effects of *Teucrium Polium* In Diabetic Rats. *Archives of 188 Iranian Medicine*. 4 (4): 188-192.

Al-Ashban R. M., Barrett D. A., Shah A. H. 2005. Effects of chronic treatment with ethanolic extract of *Teucrium polium* in mice. *Journal of Herbs, Spices & Medicinal Plants*. 11(4), 27-36

G. Ilhami, U. Metin, O. Munir, B. Suktru and K. Irfan, Antioxidant and antimicrobial activities of *Teucrium polium* L., *J. Food Technol*. 1 (2003) 9–16.

Nematollahi-Mahani S.N.; M. Rezazadeh-Kermani ^c; M. Mehrabani ^d; N. Nakhaee ^e Cytotoxic Effects of *Teucrium polium* on Some Established Cell Lines *Pharmaceutical Biology*, 45, 41 2007 , 295 - 298

Zal F, Vasei M, Rasti M ms*, Vessal M 2001. Hepatotoxicity associated with Hypoglycemic effects of *teucrium polium* In diabetic rats. *Archives of Iranian Medicine* 4(4): 188-192

Rasekh HR, Yazdanpanah H, Hosseinzadeh L, Bazmohammadi N and Kamalinejad M. 2005. Acute and Subchronic Toxicity of *Teucrium polium* Total Extract in Rats. *Iranian Journal of Pharmaceutical Research* 4: 245-249