

# Ethanollic Seed Extract of *Garcinia Kola* Reduces Epididymal Sperm Count And Some Serum Reproductive Hormone Concentrations In Adult Male Albino Wistar Rats.

## ABSTRACT

**Aims:** This study aims to elucidate the effect of *Garcinia kola* on serum reproductive hormones and sperm count in adult male albino wistar rats.

**Study design:** Albino rats were randomly assigned into 4 groups of 7 rats each.

**Place and Duration of Study:** Department of Human physiology, Madonna University, Nigeria.

**Methodology:** Group 1 served as the control group, while groups 2, 3 and 4 received 100mg/kg, 200mg/kg and 300mg/kg body weight, respectively of *Garcinia kola* extract orally once daily for 6 weeks, n=7. After 6 weeks of treatment, reproductive hormonal assay was carried out using the rat serum. Epididymal spermatozoa were collected and sperm count determined using heamocytometer.

**Results:** The experimental groups had significantly lower weights of testes ( $P<.05$ ), as compared with the control group. The weights of epididymis in the experimental groups were significantly ( $P<.05$ ) higher when compared to the control group. There was a significant decrease in the serum concentration of testosterone ( $P<.05$ ) in the experimental groups when compared to the control group. Semen analysis also showed a significant decrease in the sperm density ( $P<.05$ ) in the groups treated with *Garcinia kola* extract when compared to the control group.

**Conclusion:** Ethanollic seed extract of *Garcinia kola* showed a possible anti-spermatogenic consequence on treatment in male wistar rats, and may be detrimental to male reproductive health hence need to regulate its consumption rate.

**Key words:** *Garcinia kola*, epididymis, testosterone, sperm count, testes

## 1. INTRODUCTION

*G. kola* is a medium size tree up to 12m high, grown and cultivated in the moist forests of West Africa, South Africa and South East Asia. *Garcinia kola* seeds contains biflavanoids (kolaviron) capable of having anti-inflammatory [1] and natural antioxidant properties [2, 3]. *G. kola* have shown to have anti-fertility consequences [1,4,5], found to reduce testosterone secretion but increase LH and FSH secretion [1,6] and reduced sperm volume [4,6]. Alterations of serum concentrations of reproductive hormones are implicative of disordered spermatogenesis, which are undoubtedly major determinants of male fertility [7]. Thus, following the increase usage of *G. kola* in African traditional medicine and its consumption especially amongst the male population in Nigeria, this study aimed to

discover the possible effect of *G. kola* on serum reproductive hormones and sperm count, which are major determinants of male fertility.

## 2. MATERIAL AND METHODS

### 2.1 Preparation of Plant Extract

Fresh bitter kola seeds were purchased from a local market in Anambra state, Nigeria and authenticated in the botanical unit of the Department of Biological Sciences, Madonna University Nigeria. The seeds were peeled, sliced and dried in Astell Hearson Oven at 45 °C. The dried seeds were then grounded into fine powder using an electric blender.

50g of the powdered bitter kola was macerated in 250mls of ethanol for 48hrs and then filtered using Whatman filter paper, into a 500ml Beaker and the filtrate obtained was homogenized and concentrated to dryness in a water bath at 45 °C. The filtrate was left to evaporate until the extract was made into a solid form. Stock solution was prepared using appropriate methods.

### 2.2 Experimental Animals and Feeding Protocol

Twenty-Eight male Albino wistar rats were obtained from the animal house of the Department of Veterinary Medicine, University of Nigeria, Enugu Campus, Nigeria and acclimatized for two weeks before the onset of the experiment. The rats were fed with normal rat pellets and drinking water *ad libitum*. The rats were randomly assigned into four groups of seven rats each, housed in wire mesh cages (14hrs light and 10hrs dark cycle). Group 1 served as the control group, while groups 2, 3 and 4 received 100mg/kg, 200mg/kg and 300mg/kg body weight, respectively of extract orally once daily for six weeks. Rats in all four groups received normal rat chow and drinking water *ad libitum* during the feeding regimens. At the end of the feeding regimen the animals were sacrificed, blood samples and some organs collected for analysis.

### 2.3 Sample Collection

At the end of the 6 weeks experiment, the animals were anesthetized in a chloroform chamber, and blood was obtained via cardiac puncture using a 5ml syringe attached to a needle (21 SWG). Blood samples from each animal were put in a labeled non-heparinized sample tubes, allowed to stand for three hours in iced water and later centrifuged at 7000g for 10mins. Serum was then collected and stored at -15°C for reproductive hormonal assay. After blood collection, the animals were cut open with the aid of a dissection set and some internal organs (Testis and Epididymis) were collected and weighed. The semen from the epididymis was collected for sperm analysis (sperm count).

### 2.4 Sample Analysis

The serum testosterone and LH concentrations were determined using the enzyme linked immunosorbent assay (ELISA). The epididymis and testes were carefully removed, rinsed in normal saline solution and weighed using an electronic weighing balance. Epididymal spermatozoa were collected and sperm count was done by method of Freud and Carol [8].

### 2.5 Statistical Analysis

Data are expressed as Mean + Standard Error of Mean (SEM). Results obtained from this study were analyzed using Statistical Package for Data Analysis (SPSS) version 17.0 for windows. Analysis of Variance (ANOVA) was used to compare means, and values were compared at  $P < 0.05$ . Post Hoc multiple comparisons for difference between groups were established by Tukeys HSD.

## 3. RESULTS AND DISCUSSION

### 3.1.1 Effect of *G. kola* on male reproductive hormones

Results in Table 1 shows significant ( $P < 0.05$ ) increase in the serum concentration of LH, between 200mg/kg *G. kola* treated group ( $4.41 \pm 0.37 \mu\text{ml}$ ) and 300mg/kg *G. kola* treated group ( $4.51 \pm 0.04 \mu\text{ml}$ ) when compared to control group ( $4.21 \pm 0.04 \mu\text{ml}$ ) when compared.

Serum concentration of testosterone, in 200mg/kg *Garcinia kola* treated group ( $12.1 \pm 0.03 \text{nmol/ml}$ ) and 300mg/kg *Garcinia kola* treated group ( $10.72 \pm 0.39 \text{nmol/ml}$ ) when compared to control group ( $13.4 \pm 0.14 \text{nmol/ml}$ ) was significantly ( $P < 0.05$ ) lower. While there was no significant

81 difference between control group ( $13.4 \pm 0.14\text{nmol/ml}$ ) and 100mg/kg *Garcinia kola* treated group  
82 ( $12.9 \pm 0.04\text{nmol/ml}$ ).

### 83 3.1.2 Effect of *G. kola* on male reproductive organs

84 Table 2 shows significant ( $P<0.05$ ) increase in the weight of the epididymis in 200mg/kg *Garcinia kola*  
85 treated group ( $1.13 \pm 0.07\text{g}$ ) and 300 mg/kg *Garcinia kola* treated group ( $1.46 \pm 0.12\text{g}$ ) when  
86 compared to the control group ( $0.64 \pm 0.02\text{g}$ ) . While there was no significant difference between  
87 control group ( $0.64 \pm 0.02\text{g}$ ) and 100 mg/kg *Garcinia kola* treated group ( $0.65 \pm 0.11\text{g}$ ).

88 The weights of the testes in Table 2 shows significant ( $P<0.05$ ) decrease in the 100mg/kg  
89 *Garcinia kola* treated group ( $1.35 \pm 0.09\text{g}$ ), 200mg/kg *Garcinia kola* treated group ( $1.31 \pm 0.05\text{g}$ ) and  
90 300mg/kg *Garcinia kola* treated group ( $0.74 \pm 0.1\text{g}$ ) when compared to the control group ( $1.53 \pm$   
91  $0.16\text{g}$ ).

### 92 3.1.3 Effect of *G. kola* on Epididymal sperm count

93 Results in Table 3 shows significant ( $P<0.05$ ) decrease on sperm count in 100mg/kg *Garcinia kola*  
94 treated group ( $33.57 \pm 0.9 \times 10^6/\text{ml}$ ), 200mg/kg *Garcinia kola* treated group ( $21.44 \pm 0.4 \times 10^6/\text{ml}$ ) and  
95 300mg/kg *Garcinia kola* treated group ( $11.14 \pm 0.24 \times 10^6/\text{ml}$ ) when compared to the control group  
96 ( $58.93 \pm 0.47 \times 10^6/\text{ml}$ ).

97 **Table 1. Effect of *G. Kola* extract on male reproductive hormones in male wistar albino rats**

98 Group	Control	Group 2	Group 3	Group 4
LH ( $\mu\text{ml}$ )	$4.21 \pm 0.04$	$4.33 \pm 0.01$	$4.41 \pm 0.37^{(*)}$	$4.51 \pm 0.04^{(*)}$
Testosterone(nmol/ml)	$13.4 \pm 0.14$	$12.9 \pm 0.04$	$12.1 \pm 0.03^{(*)}$	$10.72 \pm 0.39^{(*)}$

99 Values are expressed in mean  $\pm$  SEM, (\*) statistically significant at  $P<0.05$  compared to control groups

102 **Table 2. Effect of *G. Kola* extract on male reproductive organs in male wistar albino rats**

103 Group	Control (g)	Group 2 (g)	Group 3 (g)	Group 4
Epididymis	$0.64 \pm 0.02$	$0.65 \pm 0.11$	$1.13 \pm 0.07^{(*)}$	$1.46 \pm 0.12^{(*)}$
Testes	$1.53 \pm 0.16$	$1.35 \pm 0.09^{(*)}$	$1.31 \pm 0.05^{(*)}$	$0.74 \pm 0.1^{(*)}$

104 Values are expressed in mean  $\pm$  SEM, (\*) statistically significant at  $P<0.05$  compared to control groups

107 **Table 3. Effect of *G. Kola* extract on epididymal sperm count in male wistar albino rats**

108 Group	Control (g)	Group 2 (g)	Group 3 (g)	Group 4
Sperm count ( $\times 10^6/\text{ml}$ )	$58.93 \pm 0.47$	$33.57 \pm 0.9^{(*)}$	$21.44 \pm 0.4^{(*)}$	$11.14 \pm 0.24^{(*)}$

109 Values are expressed in mean  $\pm$  SEM, (\*) statistically significant at  $P<0.05$  compared to control groups

## 112 3.2 Discussion

113 In this study, the effect of ethanolic extract of *G. kola* on sperm count, male reproductive  
114 hormones and reproductive organs was investigated. *G. kola* has been found to have high  
115 concentrations of saponnins. Saponnins decreases plasma concentrations of cholesterol and  
116 increase bile acid production [10]. Testosterone is a steroid hormone, therefore decreased plasma  
117 cholesterol will reduce the level at which cholesterol is being synthesized [11]. Direct action of *G. kola*  
118 on the testes may have caused inhibition of gonadotropic action on the testes. This was shown by  
119 Price et al. [12] who observed an irreversible combination of saponins with membranes in animal

cells, thus rendering the membrane non semipermeable. Other possibilities include preventing the release of pituitary gonadotropins and/or elevation of blood levels of testosterone (by inhibition of hepatic metabolism) thereby inducing negative feedback effect on gonadotropin release. This may be the mechanism in which *G. kola* enhances the serum levels of LH in rats. The most plausible explanation of the observations in male rats in this study could be that *G. kola* inhibits gonadotropic action on the testes. This is in collaboration with studies done by Udoh and Patil [13] which showed that phenolic compounds (saponins) are antispermatogenic.

The significant increase in the weights of the epididymis of the rats in the treatment groups of *G. kola* is in line with studies done by Oluyemi et al. [14], who reported an increase in the weights of the epididymis in rats treated with 100 and 200 mg/kg body weights of *G. kola*. Decrease in weights of the testes are in collaboration with studies done by Akinloye et al. [15] who reported a decrease in the weights of testes of male albino wistar rats fed with *G. kola* extract. This may be due to the reduction of leydig cells population in the interstitial spaces, slight reduction in the seminiferous luminal spermatozoa concentration and derangement of cells of the spermatogenic series with increase in the interstitial spaces [15].

The sperm density of rats treated with 100mg/kg body weight of *G. kola* extract showed a marked decrease when compared to the control group. Udoh [16] found out that long term administration of *G. kola* caused marked spermatogenesis arrest. This may be primarily due to decreased production of testosterone by the testes [14]. The decreased sperm count observed in this study may also be an implication of the reduced testosterone and LH concentration, which are major regulators of spermatogenesis [17].

#### 4. CONCLUSION

It can be concluded from this study that Ethanolic seed extract of *G. kola* resulted in reduced serum reproductive hormone concentrations, a dose dependent decrease in sperm count in male wistar rats, and may be detrimental to male reproductive health hence need to properly regulate its consumption rate and usage in African traditional medicine.

#### CONSENT (WHERE EVER APPLICABLE)

Not applicable

#### ETHICAL APPROVAL (WHERE EVER APPLICABLE)

For manuscripts involving animal experiments, Authors may use the following wordings for this section "All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee"

#### REFERENCES

1. Braide VB. Anti-inflammatory effect of kolaviron, a biflavonoid extract of *Garcinia kola*. *Fitoterapia*, LXIV; 433- 436.
2. Olatunde FE, Akanni OO, Emerole GO. Antioxidant and scavenging activity of flavonoid extract (Kolaviron) of *Garcinia kola* seeds. *Pharmaceut. Biol.* 2002;40:107-16.
3. Terashima K, Takaya Y, Niwa M. Powerful antioxidative agents based on garcinoic acid from *Garcinia kola*. *Bioorg Med Chem*, 2002;10(5):1619-25.
4. Akpantah AO, Oremosu AA, Noronha CC, Ekanem TB, Okanlawon AO. Effects of *Garcinia kola* seed extract on ovulation, oestrous cycle and foetal development in cyclic female sprague-dawley rats. *Nig. J. Physiol Sci.* 2005;20(1):58-62.
5. Esomonu UG, El-Taalu AB, Anuka JA, Ndodo ND, Salim MA, Atiku MK. Effect of ingestion of ethanol extract of *Garcinia kola* seed on erythrocytes in Wistar rats. *Nig. J. of Physiol Sci.* 2005;20(1):30-2.

- 173 6. Oluyemi KA, Jimoh OR, Adesanya OA, Omotuyi IO, Josiah SJ, Oyesola TO. Effects of crude  
174 ethanolic extract of *Garcinia cambogia* on the reproductive system of male wistar rats (). Afri.  
175 J. Bio. 2007;6(10).
- 176 7. Umoh IO, Emmanuel OA, Nna VU. Aqueous seed extract of *Cola nitida rubra* reduces serum  
177 reproductive hormone concentrations and sperm count in adult male albino Wistar rats. Nig  
178 Med J. 2014;55(6):456.
- 179 8. Freud M, Carol B. Factors affecting haemocytometer counts of sperm concentration in human  
180 semen. J Reprod Fertil, 1964;8:149-55.
- 181 9. Oakenfull D, Sidhu GS. Could saponins be a useful treatment for hypercholesterolaemia?.  
182 European J. of Clin Nutr. 1990;44(1):79-88.
- 183 10. Guyton AC, Hall JE. Reproductive and Hormonal Functions of the Male. Textbook of medical  
184 physiology. 2000;11:918-29.
- 185 11. Price KR, Johnson IT, Fenwick GR, Malinow MR. The chemistry and biological significance of  
186 saponins in foods and feedingstuffs. Critical Reviews in Food Science & Nutrition.  
187 1987;26(1):27-135.
- 188 12. Udoh P, Patil DR. Effects of gossypol acetate on pituitary-adrenal axis in male albino rats.  
189 Contraception, 1992; 45(3):263-71.  
190
- 191 13. Akinloye AK, Igharna OO, Olaniyi MO, Alaka OO, Oke BO. Preliminary investigations on the  
192 effects of *Garcinia kola*,(bitter kola). on the rabbit's testis and epididymis. Trop. Vet.  
193 1999;18:49-54.
- 194 14. Udoh FV. Effect of extract of *Garcinia kola* seeds and reproductive organs of the male rats.  
195 M.Sc.Thesis, University of Calabar,1 Calabar Nigeria, 1998, 132pp.
- 196 15. Seeley R, Stephens T, Tate P. Anatomy and Physiology. 6<sup>th</sup> ed. Mc Graw: Hill Publishers. p.  
197 1017-30.