

# PRELIMINARY ASSESSMENTS AND RENOPROTECTIVE EFFECTS OF METHANOL EXTRACT OF *Parquetina nigrescens* ON DIABETIC WISTAR RATS

## ABSTRACT

**Aim:** Preliminary studies and renoprotective effects of methanol extract of *Parquetina nigrescens* (MEPN) in diabetic rats were investigated.

**Methods:** Twenty-five rats divided into five groups (n=5) were used for this study. Groups 1 and 2 served as normal control and diabetic untreated respectively and each received 0.3ml distilled water. Groups 3-5 served as diabetic groups treated with 100mg/kg, 200mg/kg MEPN and 100mg/kg metformin respectively. Glucose 6 phosphate dehydrogenase (G6PDH), lactate dehydrogenase (LDH), superoxide dismutase (SOD), glutathione peroxidase (GPx), catalase (CAT) activities and albumen (Alb) level were determined using randox kits. The kidney histology was done using haemotoxylin-eosin stain. Results were analyzed using ANOVA with statistical significance taken at  $P < .05$ .

**Results:** Phytochemical screening showed the presence of alkaloids, cardenoloides, anthraquinones, tannins and flavonoids. Gas Chromatography Mass Spectrometry showed twenty-two active compounds. G6PDH and CAT significantly increased in the diabetic treated with MEPN and metformin compared with normal control. G6PDH, CAT, GPx and Alb levels were significantly increased in diabetic treated with MEPN and metformin groups when compared with diabetic untreated. BUN and CRT significantly decreased in diabetic treated with MEPN and metformin groups when compared with diabetic untreated.

**Conclusion:** MEPN possesses active components which may be useful in ameliorating the oxidative stress and renal dysfunction in diabetes mellitus.

**Keywords:** *Parquetina nigrescens*, active compounds, oxidative stress, renal dysfunction, diabetes mellitus

## 1.0 INTRODUCTION

Plants have been used for medicinal purposes long before history and many commercially available drugs are medications that were originally from plants [1]. There are synthetic medications produced to treat various forms of illnesses but, because of the cost implications and their side effects, most people prefer alternative medications to meet with their primary health needs [2]. Many medicinal plants have been reported to be useful in treatment of diabetes and its complication although diet, insulin, and other oral hypoglycemic agents have remained the mainstays of therapy for the diabetic patients for decades [3], many local plants have also been identified and tested for their antidiabetic properties. Among the medicinal plants that have been used in treatment of diabetes mellitus are *Acacia arabica*, *Aegle marmelos* and *Agrimony eupatoria*. *Acacia arabica* have been shown to cause hypoglycaemic effect in rats by stimulating insulin release [4]. *Aegle marmelos* have been shown to possess antihyperglycemic activities in streptozotocin induced diabetic rats by improving glucose utilization [5]. Similarly, aqueous extract of *Agrimony eupatoria* evoked stimulation of insulin secretion from the BRIN-BD11 pancreatic beta cell line in vitro, an effect which was found to be glucose independent [6]. *Parquetina nigrescens* is an herbaceous, perennial twine belonging to the family Asclepiadaceae. *Parquetina nigrescens* is commonly found in secondary forest and around villages in Senegal and Nigeria. It is a perennial plant with twining stems and a woody base, shortly tapering 10 to 15 cm long, 6 to 8 cm broad, smooth and long stem. In Nigeria, this plant has been shown to be useful in the treatment of anaemia [7]. This plant has also been used in the treatment of fever, inflammatory and painful disorder [8]. It has also been used in the treatment of several other ailments



which include diarrhoea, gonorrhea, menstrual disorders, insanity, intestinal worm infections, skin lesions and erectile dysfunction [9-11]. It has also been reported that aqueous extract of *Parquetina nigrescens* caused significant anti-nociception using the hot plate and formalin tests [8] and anti-inflammatory effects by reducing leucocyte migration during the process of inflammation [12]. Although several studies have shown that hyperglycaemia, if uncontrolled may results in several complications, with chronic kidney disease (CKD) being the leading cause of morbidity and mortality [13]. In this present study, the renoprotective effect of methanol extract of *Parquetina nigrescens* was investigated in diabetic wistar rats.



**Figure 1: *Parquetina nigrescens* leaves**

## **2.0 MATERIALS AND METHODS**

### **2.1 Plant collection, identification and extract preparation**

The Fresh leaves of *Parquetina nigrescens* were collected from the metropolis of Ibadan. It was identified at the Department of Botany, University of Ibadan with voucher specimen number UIH-22475 deposited in the herbarium. The plant materials were air-dried for a period of six weeks and grinded using Thomas milling machine (2mm Sieved). About 1823g of the grinded materials was soaked in 9 Liters of methanol for 72 hours. The mixture was filtered with cheesecloth and the filtrate concentrated under reduced pressure at 40°C for 20 min using a rotatory evaporator (Gallenkamp UK). The residue yielded 53.58g of methanol extract of *Parquetina nigrescens* (MEPN) which was later stored at 2-8°C prior to Physiological investigation.

### **2.2 Toxicological study and calculation of median lethal dose (LD<sub>50</sub>)**

Thirty-five rats weighing between 100-150g were used for this study and were divided into seven groups of five rats per group. Group 1 received 0.3ml distilled water; Groups 2-7 were orally given graded doses of MEPN at 500mg/kg, 1000mg/kg, 2000mg/kg, 3000mg/kg, 4000mg/kg and 5000mg/kg respectively. Rats were placed under continuous observation for 6hours. After 24 hours, rats were sacrificed under mild anesthesia (sodium thiopental 30mg/kg *i.p*) to observe changes in internal structure according to OECD method [14]. The median lethal dose was calculated as described by Karber *et al.*, 1931 [15].

### **2.3 Experimental design**

Twenty-five Wister rats weighing (100-150g) were obtained from the Central Animal House, College of Medicine, University of Ibadan, Nigeria. They were housed in well aerated cages, maintained on standard rat chow with free access to drinking water according to the guidelines and regulations of the National Institute of Health (1985) and approved by Animal Care And Use Research Ethics committee of the University of Ibadan (Reference no: 7/551/153A). Twenty-five rats were divided into 5 groups of 5 rats per group. Group 1 served as normal control, group 2 served as diabetic untreated, groups 3, 4 and 5 were diabetic treated with 100mg/kg MEPN, 200mg/kg MEPN and 100mg/kg Metformin respectively.

### **2.4 Induction of diabetes mellitus**

Diabetes was induced after a 24hour fast in group 2, 3, 4 and 5 by single intra-peritoneal injection of alloxan monohydrate (Sigma Aldrich, U.S.A) at a dose of 120mg/kg using the method



described by Carvalho *et al.*, 2003 [17]. After 72 hours of alloxan administration, only rats with fasting blood glucose level of 250mg/kg and above were considered diabetic and selected for this study.

## 2.5 Blood sample collection, determination of oxidative stress markers and renal function test

Rats were treated orally for 28 days after which rats were mildly exposed to sodium thiopental anesthesia (30mg/kg *i.p.*). Blood samples were obtained from each rat through retro-orbital sinus. The blood samples were centrifuged at 3000 r.p.m to obtain serum which was taken into another plain bottle using pasture pipette. Glucose 6 phosphate dehydrogenase (G6PDH), Lactate dehydrogenase (LDH), Superoxide dismutase (SOD), Catalase (CAT) activities and serum albumen level were determined using commercially available randox kits and their absorbance measure using spectrophotometry procedure as described by (Avinash *et al.*, 2017) [18]. Blood Urea Nitrogen (BUN) and Creatinine levels were determined using commercially available kits and their absorbance measured using spectrophotometry procedure as described by Evans *et al.*, 1968 and Khaldun *et al.*, 2017 [19-20].

## 2.6 Statistical analysis

Results obtained were analyzed using one way analysis of variance (ANOVA) followed by Neuman's keul post-hoc test. Data were expressed as mean  $\pm$ SEM with the level of statistical significance taken at  $P < .05$ .

## 3.0 RESULTS

### 3.1 Phytochemical screening and GC-MS studies on MEPN


Table 1 and Figure 1 showed Phytochemical screening and Gas Chromatography Mass Spectrometry (GC-MS) analysis of MEPN respectively. The phytochemical screening showed positive results for alkaloids, cardenoloids, anthraquinones, tannins and flavonoids. GC-MS analysis of MEPN showed the presence of twenty-one (22) bio-active compounds which include Alpha-phellandrene(5.819), Cymene(6.117), Beta-curcumene(12.005), Alpha-begarmotene(12.126), Beta-bisabolene(12.262), Naphthalene(12.455), Diethylphthalate(12.995), 1,12-tridecadiene(15.033), 5-ethyl-2-furaldehyde(15.283), 7-hexadecyne(15.478), Pyranos(15.576), decanoic acid(15.867), Mannos(16.275), Thiophene(16.438), Hexadecanoic acid(16.813), Octadecanoic acid(17.710), Catrienoate(17.772) Phytol(17.878) Octadecatrienoate(18.274), Octadecanoic(18.388), 9-Octadecenamide(20.222), Squalene(28.028).

**Table 1: Phytochemical analysis of methanol extract of *Parquetina nigrescens***

Phytochemicals compounds	Methanol extract of <i>Parquetina nigrescens</i> MEPN
Alkaloids	+
Cardenoloids	+
Anthraquinones	+
Saponins	-
Tannins	+
Flavonoids	+

+ = Present      - = Absent





**Figure 2: Gas Chromatography Mass Spectrometry (GC-MS) analysis of MEPN**

### 3.2 Toxicological study of different doses of MEPN administered orally in rats

Table 1 showed the effect of graded doses of MEPN in normal rats. There were no cases of mortality in the experimental rats and this means that the LD<sub>50</sub> of MEPN was greater than 5000mg/kg. However, fecal materials and urine were found in group administered with 5000mg/kg MEPN.

**Table 2: Toxicological study of different doses of MEPN administered orally in rats**

S/n	Groups	Mortality x/N	Symptoms (0-6 hrs)
Group 1	0.3ml	0/5	Nil
Group 2	500mg/kg	0/5	Nil
Group 3	1000mg/kg	0/5	Nil
Group 4	2000mg/kg	0/5	Nil
Group 5	3000mg/kg	0/5	Nil
Group 6	4000mg/kg	0/5	Nil
Group 7	5000mg/kg	0/5	Frequent Defecation and urination

### 3.3 Anti-oxidative parameters in normal, MEPN and Metformin treated groups.

Table 3 showed changes in oxidative stress parameters in normal and treated rats. There was significant increase ( $P<.05$ ) in G6PDH activities in normal control, diabetic treated with 200mg/kg MEPN and 100mg/kg metformin groups when compared with diabetic untreated and diabetes treated with 100mg/kg MEPN respectively. There was also significant increase in G6PDH activities in diabetes treated with 100mg/kg metformin when compared with normal control and diabetes + 200mg/kg MEPN. Lactate dehydrogenase (LDH) was significantly lower ( $P<.05$ ) in diabetes treated with 100mg/kg MEPN when compared with diabetes treated with 100mg/kg metformin.

Catalase (CAT) activities significantly increase ( $P<.05$ ) in normal control, diabetes + 100mg/kg MEPN, diabetes + 200mg/kg MEPN, diabetes + 100mg/kg Metformin when compared with diabetic



untreated. There was significant increase in CAT activities in normal control, diabetes + 200mg/kg MEPN, diabetes + 100mg/kg Metformin when compared with diabetes treated with 100mg/kg MEPN. A significant increase was also observed in CAT activities in normal control and diabetes + 200mg/kg MEPN when compared with diabetes treated with 100mg/kg meformin

Glutathione peroxidase (GPx) significantly increased ( $P<.05$ ) in normal control, diabetes treated with 200mg/kg MEPN and 100mg/kg Metformin groups when compared with diabetic untreated and diabetes treated with 100mg/kg MEPN respectively.

The level of albumen was significantly higher ( $P<.05$ ) in normal control, diabetes treated with 100mg/kg, 200mg/kg MEPN and diabetes treated with 100mg/kg Metformin when compared with diabetic untreated.

165

**Table 3: Antioxidative parameters in normal, MEPN and Metformin treated groups.**

Experimental groups	G6PDH (U/L)	LDH (U/L)	SOD (U/L )	CAT (U/ml)	GPx (U/L)	ALB (mg/dl)
Normal control (0.3ml distilled water)	37.78 ± 3.29	27.17 ± 1.19	149.2±7.98	166.6±5.65	7.49±0.94	2.38±0.12
Diabetic untreated (0.3ml distilled water)	20.52 ± 0.52 <sup>a</sup>	23.30 ± 3.84	143.0±27.09	113.1±3.48 <sup>b</sup>	0.78±0.26 <sup>a</sup>	1.32±0.04 <sup>b</sup>
Diabetes + 100mg/kg MEPN	20.80 ± 2.18 <sup>a</sup>	17.49 ± 1.09 <sup>c</sup>	165.0±8.52	131.5±7.884 <sup>a</sup>	0.99±0.39 <sup>a</sup>	2.35±0.16
Diabetes +200mg/kg MEPN	30.46 ± 0.61	23.36 ± 3.50	140.0±20.31	168.4±2.36	4.53±0.72	2.48±0.14
Diabetes + 100mg/kg Metformin	45.14 ± 2.98 <sup>#</sup>	32.18 ± 3.24	141.0±13.45	186.2±4.85 <sup>#</sup>	5.43±1.46	2.58±0.03

167

168a were expressed as Mean ± SEM;  $P<0.05$ . <sup>a</sup> indicate values significantly different from normal control, 169 diabetes + 200mg/kg MEPN, diabetes + 100mg/kg Metformin. <sup>b</sup> indicate values significantly different from 170 normal control, diabetes + 100mg/kg MEPN, diabetes + 200mg/kg MEPN, diabetes + 100mg/kg Metformin. <sup>#</sup> 171 indicate values significantly different from normal control, diabetes + 200mg/kg MEPN. <sup>c</sup> indicates value 172 significantly different from diabetes + 100mg/kg metformin (n=5)

173

### 174 3.4 Blood Urea Nitrogen in normal, MEPN and Metformin treated groups


175 Figure 2 showed blood urea nitrogen (BUN) in normal, diabetes treated with MEPN and metformin 176 rats. BUN significantly increased ( $P<.05$ ) in normal control, diabetes treated with 100mg/kg, 200mg/kg 177 MEPN and 100mg/kg Metformin when compared with diabetic untreated. BUN significantly increased 178 ( $P<.05$ ) in diabetes treated with MEPN and metformin treated groups when compared with diabetic 179 untreated. There was no significant difference in MEPN treated groups when compared with normal 180 control but, BUN significantly increased in metformin treated group when compared with normal 181 control.

182

183



184




185

186 **Figure 3: Blood Urea Nitrogen in normal, diabetes treated with MEPN and Metformin groups.**

187 Data were expressed as Mean ± SEM;  $P < .05$ . <sup>a</sup> indicates value significantly different from the normal control, diabetes + 100mg/kg MEPN, diabetes + 200mg/kg MEPN, diabetes + 100mg/kg Metformin. <sup>b</sup> indicates value significantly different from normal control, diabetes + 100mg/kg MEPN, diabetes + 200mg/kg MEPN (n=5).

### 191 3.5 Creatinine level in normal, MEPN and Metformin treated groups.

192 Figure 3 showed creatinine level in normal, MEPN and Metformin treated groups. There was significant increase ( $P < .05$ ) in creatinine level in diabetic untreated group when compared with normal control. Similarly, creatinine level significantly increased in 100mg/kg MEPN treated group when compared with normal control. However, there was significant decrease ( $P < .05$ ) in creatinine level in diabetes treated with 200mg/kg MEPN when compared with diabetes untreated and diabetes treated with 100mg/kg MEPN.




198

199 **Figure 4: Creatinine level in normal, MEPN and Metformin treated groups.**



Data were expressed as Mean  $\pm$  SEM;  $P < .05$ . <sup>a</sup> indicate values significantly different from the normal control, diabetes+200mg/kg MEPN, diabetes+100mg/kg metformin (n=5)

### 3.6 Photomicrographs of kidney in normal, MEPN and Metformin treated rats



**Plate 1 (A – E):** Shows sections stained with H & E showing architecture of the Kidney in A (Control), B (Diabetic untreated), C (Diabetes + 100mg/kg MEPN), D (Diabetes + 200mg/kg MEPN), E (Diabetes + 100mg/kg Metformin). Plate 1A showed architecture of the kidney with normal glomerulus and capsular space (white Arrow), Plate 1B showed architecture of the kidney with degenerated glomerulus and Bowman's capsule with inflammatory cells (Black Arrows). Plate 1C& E showed architecture of the kidneys with distorted glomerulus. Plate 1D showed architecture of the kidney with glomerulus and capsular space comparable with the normal control (Plate 1A) (Blue Arrow) X 400.

### 4.0 Discussion and Conclusion

This present study investigates the renoprotective effects of methanol extract of *Parquetina nigrescens* (MEPN) in alloxan induced diabetic rats. The acute toxicity test performed in the experimental rats using MEPN showed no case of mortality in various groups treated and this suggests that MEPN is non-toxic even at higher doses administered. According to Hodge and Sterner (2005) [20] toxicity scale [21], a plant extract with an LD<sub>50</sub> < 2000mg/kg is said to be toxic and since the LD<sub>50</sub> of MEPN from our study was >5000mg/kg therefore, MEPN is said to belong to a non toxic category of medicinal plant. However, this finding is not in agreement with the report of Lyon *et al.*, [22] who had earlier reported that the latex part of the plant is toxic and even used in making poison.



Diabetes mellitus is a multi-factorial disease in which increased oxidative stress plays an important pathogenic role [23]. A low Glucose-6-phosphate dehydrogenase (G6PDH) activity has been considered to play a role in increased oxidative stress because of less NADPH produced during pentose phosphate pathway [24]. The significant decrease in G6PDH activities observed in diabetic untreated is consistent with the reports of Wan *et al.*, 2002 [25] who showed that positive correlation exist between a reduced G6PDH activities and diabetes mellitus. In diabetic untreated, less NADPH is likely produced to reduce the oxidized glutathione in the pentose phosphate pathway leading to altered glucose tolerance [26]. The increase in G6PDH observed in 200mg/kg MEPN group indicates that more reduced glutathione may have been produced to mop up free radicals that may have been generated in diabetes mellitus [27]. The increase in G6PDH observed in metformin treated group suggests that metformin may also have the potential of reducing oxidative stress as well [26].

The decrease observed in Catalase (CAT) activities in diabetic untreated may indicate that more hydrogen peroxide is produced and if this is allowed to accumulate may be potentially toxic at high concentration [28]. However, the increase observed in the treatment groups indicates that more hydrogen peroxide may have been broken down to harmless water and oxygen [28].

**Albumen** is a multifunctional protein with 585 amino acids and one reduced cysteine residue (Cyst 34) and a molecular weight of 66kDa [29]. The significant decrease in albumin in diabetic untreated may likely indicates a decrease in the quantity of the reduced Cyst 34 residues in albumin to scavenge hydroxyl radicals [30]. However, the increase observed in albumin level in the treatment groups suggests an increase in reduced cyst 34 in albumin which is converted to sulfenic acid that is important in redox modulation of reactive species [31]. It is also likely that MEPN may have reversed methionine sulfoxide that may have been produced in diabetic untreated back to methionine residue in albumin which is highly susceptible to oxidative damage [32] and these effects observed in 100mg/kg and 200mg/kg MEPN were comparable to metformin treated group.

Creatinine is a metabolic waste product produced from muscle creatine and excreted through the kidneys [33]. The significant increase in creatinine level in diabetic untreated rats is an indication of kidney damage, sclerosis, inflammation and decreased glomerular filtration rate [34] as shown by the photomicrograph (Plate 1B). This observation is consistent with previous reports which identified creatinine as a basic marker of renal dysfunction in diabetes mellitus [35]. However, the significant decrease in creatinine level in 100mg/kg and 200mg/kg MEPN was comparable to that of metformin treated group and this implies that MEPN may have the ability to improve kidney functions in diabetic condition so that more waste products can be filtered from the blood and excreted in urine as shown by the photomicrograph (Plate 1C&D) [36].

Blood urea nitrogens (BUN) are nitrogenous waste products produced from breakdown of protein into ammonia which undergoes deamination by liver enzymes to produce urea which are excreted through the kidneys. The increase in BUN in diabetic untreated rats may be due to damage to the glomerulus of the renal tubular cells from uncontrolled hyperglycemia [37] as shown by the photomicrograph (Plate 1B). However, oral administration of MEPN at 100mg/kg and 200mg/kg significantly decrease urea nitrogen and this implies that MEPN could also repair the damaged renal tubular cells and increase excretion of urea nitrogen in the urine as shown by the photomicrograph (Plate 1C& D).



In summary, administration of MEPN caused increased G6PDH activities; an effect which may be due to its ability to reduce oxidized glutathione. MEPN caused increased Catalase activities, an effect which may result from decreased formation of hydrogen peroxide. Similarly, MEPN caused increased albumen level; which may be due to availability of more cysteine 34 amino acid residues for redox modulation of free radicals. MEPN also caused decreased level of creatinine and blood urea nitrogen, an effect which may be due to its ability to repair the damaged renal tubular cells, reduce thickening of glomerulus and improve ultrafiltration. This various effects of MEPN may be due the presence of phytochemicals and active compounds present in the plant extract as shown by the results of phytochemical screening and Gas chromatography mass spectrometry analysis.

## 6.0 REFERENCES

1. Schipmann U, Leaman DJ, Cunnigham AB. Impact of cultivation and gathering of medicinal plants on biodiversity: Global trends and issues: in biodiversity and the ecosystem approach in Agriculture, forestry and fisheries. FAO. 2002; 1-21
2. Zhang J, Yulong H, Tiande H, Yunpu W. Effect of Azadirachta indica leaf extract on serum lipid profile changes in normal and Streptozotocin induced diabetic rats. African Journal of Biomedical Research. 2005; 8:101- 104.
3. Ismail N, Becker B, Strzelczyk P. Renal disease and hypertension in non-insulin-dependent diabetes mellitus. Kidney int. 1999; 55: 1–28.
4. Singh LW, Traditional medicinal plants of Manipur as antidiabetics. J Med Plant Res. 2011; 5(5): 677-687.
5. Ayodhya S, Kusum S, Anjali S. Hypoglycaemic activity of different extracts of various herbal plants Singh. Int J Ayurveda Res Pharm. 2010; 1(1): 212-224.
6. Bnouham M, Ziyat A, Mekhfi H, Tahri A, Legssyer A. Medicinal plants with potential antidiabetic activity-a review of ten years of herbal medicine research (1990-2000), Int J Diabetes Metab. 2006; 14:1-25.
7. Saba AB, Oyagbemi AA, Azeez OI, Antidiabetic and haematinic effects of *Parquetina nigrescens* on alloxan induced type-1 diabetes and normocytic normochromic anaemia in Wistar rats African Health Sciences. 2010; 10(3): 276 – 283
8. Owoyele BV. Studies on the Analgesic, Anti-Inflammatory and Antipyretic Effects of *Parquetinan nigrescens* Leaf Extract J Ethnopharmacol. 2008; 122 (1): 86-90.
9. Sofowora A, Medicinal Plants and Traditional Medicine in Africa, 2<sup>nd</sup> edition. Spectrum Books Limited (Publisher), Ibadan, Nigeria, 1993;134–156.
10. Iwu, M.M.: Handbook of African Medicinal Plants. CRC Press, Boca Raton. 1993; 351.
11. Odetola AA, Oluwole FS, Adeniyi BA, OLatiregun AM, Ikupolowo OR, Labode O, Busari, KO, Shorinola JA. Antimicrobial and gastroprotective properties of *Parquetina nigrescens* (Afzel.) Bullock. Journal of Biological Sciences. 2006; 6:701–705.
12. Olajide OA, Makinde JM, Okpako DT. Evaluation of the anti-inflammatory property of the extract of *Combretum micranthum* G. Don (Combretaceae). Inflammopharmacology. 2003; 11: 293–298.



- 308 13. Seyed Bahman Ghaderian, Fatemeh Hayati, Shokouh Shayanpour, Seyed Seifollah Beladi  
309 Mousavi . Diabetes and end-stage renal disease; a review article on new concepts J Renal Inj  
310 Prev., 2015; 4(2): 28–33.
- 311 14. Fisher K. The Impact of School Infrastructure on Student Outcomes and Behaviour”, Schools  
312 Issues Digest, Department of Education, 2000 Training and Youth Affairs, Australian Government.  
313
- 314 15. Karber G. Beitrag zur kollektiven Behandlung pharmakologischer Reihenversuche. Arch. Exptl.  
315 Pathol. Pharmacol, 1931; 162:480-483.
- 316 16. National Institute of Health. Guide for the Care and Use of Laboratory Animals. NIH publication.  
317 1985; 85 – 23.  
318
- 319 17. Carvalho E, Carvalho N, Ferreira L. Experimental model of induction of diabetes mellitus in rats.  
320 Acta Cir Bras. 2003; (18)
- 321 18. Avinash Krishnegowda,<sup>a</sup> Nagaraja Padmarajaiah, <sup>b</sup>Shivakumar Anantharaman, <sup>c</sup>KrishnaHonnur<sup>b</sup>:  
322 Spectrophotometric assay of creatinine in human serum sample Author links open overlay panel  
323 Arabian Journal of Chemistry., 2017; 10: 2  
324
- 325 19. Evans RT. Manual and automated methods for measuring urea based on a modification of its  
326 reaction with diacetyl monoxime and thiosemicarbazide. J Clin Pathol. 1968; 21(4): 527–52  
  
327
- 328 20. Khaldun MohammadAl Azzam,<sup>a</sup> WafaaEl Kassed New: Simple, sensitive and validated  
329 spectrophotometric method for the determination of salicylhydroxamic acid in capsules and raw  
330 material according to the ICH guidelines 2017. Author links open overlay panel Egyptian Journal  
331 of Basic and Applied Sciences.  
332
- 333 21. Hodge, A. Sterner, B.: Toxicity classes. In: Canadian center for occupational Health and safety  
334 (2005) (<http://www.ccohs.ca/oshanswers/chemicals/id50.htm>).
- 335
- 336 22. Lyon RC, Taylor JS, Porter DA. Stability profiles of drug products extended beyond labeled  
337 expiration dates. Journal of Pharmaceutical Sciences. 2006; 95:1549-60.  
338
- 339 23. Moussa, S.A. Oxidative stress in diabetes mellitus. Romanian J. Biophys. 2008; 18: 225–236  
340
- 341 24. Adinortey MB, Owusu, R.K. Galyuon, I.K.A. Ekloh, W. Owusu, I. Larbi, D.A.: G6PD Deficiency – a  
342 potential risk factor for development of diabetes mellitus. Journal of Medicine and Medical  
343 Science. 2011; 28: 1017 – 1021
- 344 25. Wan GH, Tsai SC, Chiu DT. Decreased blood activity of glucose-6-phosphate dehydrogenase  
345 associates with increased risk for diabetes mellitus Endocrine. 2002; 19:191-195  
346
- 347 26. Mehta A, Mason PJ, Vulliamy TJ. Glucose-6-phosphate dehydrogenase deficiency. Clin.  
348 Haematol., 2000; 13: 21-38.  
349
- 350 27. Gaskin RS, Estwick D, Pedi R. G6PD deficiency: its role in the high prevalence of diabetes  
351 mellitus. Ethnicity and Disease., 2001; 11: 749- 754
- 352 28. Góth L, Nagy T, Paragh G, Káplár M. Blood Catalase Activities, Catalase Gene Polymorphisms  
353 and Acatlasemia Mutations in Hungarian Patients with Diabetes Mellitus. Glob J Obes Diabetes  
354 Metab Syndr. 2016; 3(1): 001-005



- 355 29. Marjolaine Roche<sup>1</sup>, Philippe Rondeau<sup>1</sup>, Nihar Ranjan Singh, Evelyne Tarnus, Emmanuel  
356 Bourdon: The antioxidant properties of serum albumin *FEBS Letters*. 2008; 582: 1783–1787
- 357 30. Gutteridge, J.M. Antioxidant properties of the proteins caeruloplasmin, albumin and transferrin. A  
358 study of their activity in serum and synovial fluid from patients with rheumatoid arthritis. *Biochim.*  
359 *Biophys. Acta*. 1986; 869, 119–127.
- 360 31. Oettl K, Stauber RE. Physiological and pathological changes in the redox state of human serum  
361 albumin critically influence its binding properties. *Br. J. Pharmacol.*, 2007;151:580–590.
- 362 32. Bourdon E, Loreau N, Lagrost L, Blache, D. Differential effects of cysteine and methionine  
363 residues in the antioxidant activity of human serum albumin. *Free Radic. Res.*, 2005; 39:15–20.
- 364 33. Stryer L, Tymoczko JL, Berg J. *Biochemistry*, 5th edition. New York, 1997; 324-50.
- 365 34. Henry JB. *Clinical diagnosis and management by laboratory methods*. Philadelphia, PA, 2001;  
366 187- 196.
- 367 35. Cholongitas E, Shusang V, Marelli L, Nair D, Thomas M, Patch D. Renal function assessment in  
368 cirrhosis - difficulties and alternative measurements, 2007; *Aliment Pharmacol Ther*. 26:969–78.
- 369 36. Meyer TW, Hostetter TH, Uremia. *N Engl J Med.*, 2007; 357(13):1316–25.
- 370 37. Molitoris BA. Acute kidney injury. In Goldman L, Ausiello D, editors. *Cecil Medicine*. Philadelphia,  
371 Pa: Saunders Elsevier., 2007; 121.