

Original Research Article

Potential of column purified fractions of *Allium cepa* against intermediate host of urinary schistosomiasis (*Bulinus globosus*) in Sokoto, Nigeria

ABSTRACT

Potential of column purified fractions of *Allium cepa* bulb against intermediate hosts of urinary schistosomiasis (*Bulinus globosus*) was conducted in laboratory condition. The fresh bulbs of *A. cepa* were purchased from Ramin Kura market of Sokoto, identified and authenticated by taxonomist. The bulbs were sliced in to pieces, air dried and powdered. Extracts were obtained using methanol as polar then purified with silica gel as a stationary phase while N-hexane and ethyl acetate (1:1) as mobile phase. Thirteen fractions each fraction containing 10ml of the effluent was collected, the collected extracts were left open for evaporation for 48hours. Ten adult *B. globosus* were immersed in 3liters of water containing different concentrations of the fraction and each treatment was replicated three times with control in the same condition without treatment, observations were recorded after 24hours up to 96hours. The toxicity experiment showed that fractions (F₇, F₈, F₆ and F₉) were most toxic fractions, LC₅₀ after 96 hours was 19.371mg/l. based on findings from this research it can be concluded that, *A. Cepa* was very potent and can be used for control of *B. globosus* in order to prevent urinary schistosomiasis infection in endemic areas and drugs industries may use the extracts of these plants for production of molluscicides

Keywords: *Allium cepa*, *Bulinus globosus*, Ramin kura, Schistosomiasis, Molluscicides.

INTRODUCTION

Urinary schistosomiasis also known as Bilharziasis or snail fever, is a disease caused by infection of digenetic blood trematode (*Schistosoma haematobium*) [1]. Urinary Schistosomiasis remains among the most important public health problems in Nigeria [2], [3]. Generally, various factors that are responsible for the increment and persistent of infection with schistosomiasis in Nigeria include climatic changes, proximity to water bodies, irrigation and dam construction as well as socio-economic factors such as occupational activities and poverty as well as abundance of infected intermediate host such as *Bulinus* species [4]. Although, use of molluscicides to eradicate snail vector is considered to be the best method to eliminate schistosomiasis, because killing of intermediate host disrupts the life cycle of the parasite and stop the transmission of infection [5]; synthetic molluscicides widely used for the effective control of the intermediate host remain cost-effective and have rapid toxicity to the aquatic environment [6]. Moreover, most of the synthetic molluscicides such as Niclosamide, endothall, chloramines, ozone and hydrogen peroxide are toxic to non-target animals and may have long-term detrimental effects on the aquatic life [7], [8], [9].

38 Although, several groups of compounds present in various plants were determined to have
39 toxic effect to target organisms at acceptable doses [10], [11], [12]; there is still an urgent
40 need for highly toxic plants in order to avoid transmission of the parasitic disease like urinary
41 schistosomiasis [13]. Furthermore, in the environment like Sokoto where a large fraction of
42 population is still dependent on lakes, rivers and dams water for their everyday supplement;
43 it is highly injurious to use synthetic chemicals to control the disease.

44 In addition, *Allium cepa* is available, accepted, cheap and safe plants, which have been used
45 for many reasons (such as food, flavour, medication etc.) among Sokoto people. Therefore,
46 potential of column purified fractions of *Allium cepa* bulb plants on intermediate host of
47 urinary schistosomiasis was determined to serve as an increment for cheaper, non-toxic and
48 efficient way of controlling urinary schistosomiasis infections in Sokoto.

49 **MATERIALS AND METHODS**

50 **Collection and Identification of Plant materials**

51 Red globe onionbulbsused for this study were purchased from Ramin Kura market Sokoto,
52 each of the plant bulbwas identified and authenticated by taxonomist, in the herbarium,
53 Botany Unit, Department of Biological Sciences, Usmanu Danfodiyo University Sokoto.
54 Voucher number collected for the identified plants was UDUH/ANS/0088

55 **Preparation of the Plants Powder**

56 Fresh bulbs of *Allium cepa*was washed with borehole water, sliced in to smaller pieces and
57 air dried under shade for 14 days, the dried bulbs and cloves were grinded into fine powder
58 using a grinder and stored in air-tight container [14].

59 **Preparation of the Plants Methanol Extract**

60 The Methanolic extractsof *Allium cepa*was prepared using cold maceration method
61 described by [15]. 100g was weighed and transferred into clean sterile bottle, soaked in to
62 300ml of Methanol, tightly covered and allowed for 72hours at room temperature, the
63 suspension was stirred occasionally after each 24hrs and filtered in to a sterile bottles using
64 wet man filter paper No. 1. The filtrate was left in a vacuum for 48hors then finally taken for
65 column purification.

66

67

68 **Column Purification**

69 The plant extract was purified according to method described by [16]. The column with
70 95×45 size was placed in the vertical position; 140mg of cotton wool was inserted and
71 pushed down to the bottom of the column which reached 1.5cm from down of the column to
72 avoid escape of silica gel; one hundred and twenty gram (120g) of a dried stationary powder
73 of silica gel (60-120 mesh) was added in to the column; 100ml of mobile phase (hexane)
74 was added to flushed through the column and made it wetted; while flushing through, the

column was slapped several times and ensured the air bobbles were removed; dropping funnel was attached to the top of the column and extract to be purified was poured gently in to the column and sank in to the silica gel; 80ml of ethyl acetate and hexane in ratio of 1:1 (i.e. 40ml:40ml) was added continuously and simultaneously through the funnel from the top of the column with carefully open stop cock until the extract eluted; from each plant extract; 13 fractions were collected and each fraction was 10ml; the fractions were left for two days to evaporation

Snail Collection

Adults *Bulinus globosus* snails with shells length between 9 to 11mm long were collected from Kwalkwalawa River of Wamakko Local Government, Sokoto State, between 10:00am to 12:00noon whenever needed using a scoop as described by [17]. The scoop comprised wooden frame, supporting mesh and mounted handle. To collect the snails, the scoop net was immersed and pushed to 16 to 20m; the scoop was lifted upward vertically to ensure proper collection; the collected snails were transferred in to plastic bucket containing borehole water which was left open for 24 hours after fetched, the snails were brought to the parasitology laboratory, Department of Biological Sciences, Usmanu Danfodiyo University Sokoto.

Identification of *Bulinus globosus* snail

The snails were taken to Department of Zoology, faculty of life sciences, Ahmadu Bello University Zaria for confirmation of the animals. Snails were identified at Museum of Natural History of the Department of Zoology by malacologist as *Bulinus globosus* and 7B was assigned as cabinet number for the animal. Finally, the snails were taken back to the parasitology laboratory Usmanu Danfodiyo University Sokoto for toxicity study.

Maintenance of the Snails

In the Parasitology Laboratory of Department of Biological Sciences, Usmanu Danfodiyo University, Sokoto; identified snails were kept in groups of 40 in plastic buckets containing borehole water left for 24 hours after fetched. Water in the plastic buckets was changed twice in a week to eliminate the contaminants so as to prevent fouling. The snails were kept for a period of three weeks in the laboratory before experiment [17].

Toxicity Experiment for the Intermediate Host

Each purified fraction of *Allium cepa* was tested against *Bulinus glubosus* snails according to toxicity experiment described by [18]. For experimental test of each fraction, ten adult snails were kept in plastic buckets containing concentration of the fraction (the treatment were 17.15mg/l; 18.20 mg/l; 20.25 mg/l; 26.20 mg/l; 31.75 mg/l; 38.90 mg/l; 40.20 mg/l; 39.75 mg/l; 37.25 mg/l; 34.45 mg/l; 26.20 mg/l; 21.75 mg/l and 14.35 mg/l) for 96 hours; each of the fraction was replicated three times; control animals were kept in similar conditions, but without treatment for each three replicates. Mortality was recorded after each 24 hours (i.e.

after 24hours; 48hours; 72hours and 96hours respectively) and dead snails were removed to avoid any contamination in aquarium water. Absence of response to a needle probe was considered as evidence of the snail dead. After every 24hour of the experimental set up, mortality of each treated group was calculated using Abbott's formula by subtracting number of survival in the treated group from number of survival in the untreated (control) group multiply by one hundred and recorded.

Data Analysis

The data obtained from mean mortality were analysed using one way analysis of variance (ANOVA) to determine statistically significant differences between means at $P < 0.05$; lethal concentration (LC50) at a confidence interval of 95%, using probate analysis method was estimated in the Minitab statistical software package. .

RESULTS AND DISCUSSION

Potentials of column purified fractions of *AlliumcepaBulinus globosus* was time and concentration dependence, it was observed that; immediately after introducing the experimental snails in to various concentration of the fractions, the snails attempt to escape from toxicity of the plants by moving around (distressing) and becoming firmly attached to the wall of experimental container; shells of the snails also were filled up. Nevertheless, swelled pores were observed on the foot of some snails, shell's colour of most snails changed from brown to dark brown; mortality ranged between 100% (10.0) and 20% (2.0) with LC50 19.371mg/l after 96 hours in to various fraction. F₇ had higher mortality of 100% (10.0) followed by F₈ with 93.3% (9.33); F₆ showed 76.7% (7.67) mortality and mortality of 63.3% (6.33) was observed in F₉; in the same way, F₁₀ and F₅ showed mortality of 53.3% (5.33) and 46.7% (4.67) respectively. Although, F₄ and F₃ showed mortality of 43.3% (4.33) and 40% (4.00) respectively; also F₂, F₁, F₁₁ and F₁₂ showed mortality of 36.7% (3.67), 33.3% (3.33), 33.3% (3.33) and 30% (3.0) respectively. Lastly, F₁₃ had lowest mortality of 20% (2.00) (table 1). P value was <0.05, hence, there was significant difference for the mortality of the snails after 96 hours in to different concentrations of *Allium cepa*

Table 1: Potential of column purified fractions of *Allium cepa* (L) on *Bulinus globosus* after 96 hours

Fraction Number	Concentration (mg/l)	No. of Snails in the Concentration	Mean Mortality of the Snails	Mortality of the snails (%)
F ₁	14.90	10	3.33±0.33 ^f	33
F ₂	17.70	10	3.67±0.33 ^f	37
F ₃	21.45	10	4.00±0.00 ^e	40
F ₄	25.10	10	4.33±0.33 ^e	43
F ₅	30.90	10	4.67±0.33 ^d	47
F ₆	38.15	10	7.67±0.33 ^b	77
F ₇	42.15	10	10.00±0.00 ^a	100
F ₈	38.75	10	9.33±0.33 ^a	93
F ₉	35.05	10	6.33±0.33 ^c	63

F ₁₀	34.30	10	5.33±0.33 ^d	53
F ₁₁	24.85	10	3.33±0.33 ^f	33
F ₁₂	22.60	10	3.00±0.00 ^f	30
F ₁₃	11.90	10	2.00±0.00 ^g	20

141 Values are expressed as Mean±SEM of three replicates. Values in row having different
142 superscript differs significantly at $P<0.05$ level (One Way ANOVA).

143 Discussion

144 An attempt to escape from toxicity, filling up of the shells of the experimental snails, swelled
145 pores on the foot of some snails and changed of shells colour was due to toxicity of the
146 plants extracts on *Bulinus globosus*; another researchers also observed movement of snails
147 to the side of the container in an attempt to escape from the test media
148 containing *Althernanthera sessilis* treated water [19].

149 Potential of *Allium cepa*, on *Bulinus globosus* were concentration and time dependent,
150 because the higher the concentration in the fraction of the plants the higher mortality was
151 observed, similar finding was reported by [21] on the toxicity of *Agave filifera* plant extracts
152 on *Biomphalaria alexandrina*. [20], also reported similar observation while studying
153 molluscicidal activity of *Bauhinia variegata* and *Mimusops elengi* plant extracts against
154 the *fasciola* vector *Lymnaea acuminata*.

155 Snails treated in F₇, F₈, F₆ and F₉, showed higher mortality and mean mortality of these
156 fractions were significantly different from that of other fractions may be due to potent
157 phytochemicals such as alkaloid, flavonoid, saponins, glycosides and tannins were eluted in
158 higher quantities in these fractions; this was in line with the report of several researchers
159 such as [21]. who reported that, potency of plants on the snails was due to presence of
160 saponins content [23], also reported that, higher toxicity occurred due to higher amount of
161 tannins compounds and higher amount of flavonoid; similarly, glycoside and alkaloid
162 components were reported with higher toxicity to the snails by [24]. Although, LC₅₀ value in
163 column purified fraction of *Allium cepa* was 19.4mg/L, the value was in the same range with
164 thatof Niclosamide (11.8 mg/L) which is one of the standard molluscicidal drugs as reported
165 by [24].

166 CONCLUSION AND RECOMMENDATIONS

167 Column purified fractions of *Allium cepa* showed that, F₇, F₈, F₆ and F₉were most toxic
168 fractions; the toxicity maybe due to composition of active phytochemicals such as alkaloid,
169 saponins, glycosides, tannins and flavonoid in higher amount.

170 The promising results obtained from column purified fractions of *Allium cepa* (bulbs)
171 encourages further investigations for the phytochemicals present and there quantity in each
172 fraction.

Governmental and nongovernmental organizations should enlighten public on the effectiveness of the plants for control of intermediate host of urinary schistosomiasis (*Bulinus globosus*) since the plant is inexpensive, readily available and environmentally safe.

REFERENCES

1. Downs JA, van Dam GJ, Chungalucha JM. Association of schistosomiasis and HIV infection in Tanzania. *Am J Trop Med Hyg.* 2012; 87:868-73.
2. Bala AY, Ladan MU, Mainasara M. Prevalence and Intensity of urinary Schistosomiasis in Abarma Village, Gusau, Nigeria. *Science World Journal.* 2012; 7(2):197-6343.
3. Abiola F, Adenowo BO, Bolajoko IO, Abidemi PK. Impact of human schistosomiasis in sub-Saharan Africa; *Brazilian J. of Infect. Diseases.* 2015; 19:196-205
4. Mostafa MH, Sheweita S, AConnor PJ. Relationship between Schistosomiasis and Bladder Cancer, *J. Clin Microbiol Rev.* 2012; 12:1
5. Wynn TA, Thompson RW, Cheever AW, Mentink-Kane MM. Immunopathogenesis of schistosomiasis. *Immunol Rev.* 2014; 201:156-67.
6. Bakry FA, Mohamed RT, El-Hommossany K. Biological and biochemical responses of *Biomphalaria alexandrinato* some extracts of the plants *Solanum siniacum* and *Artemisia judaica* L. *Pesticide Bio-chemistry Physiology.* 2011; 99, 174-180.
7. Osman, EA, Hilali, AH, Koko WS, Muhmoud AA. Molluscicidal activity of some Sudanese wild plants. *Journal of Science and Technology.* 2012; 8: 111-118.
8. Ngassapa O, Beecher CWW, Pezzuto JM, Farnsworth NR, Henderson TO, Boye GL. Isolation of Echinocystic Acid-3-O-sulfate. A new triterpene, From *Tetrapleura tetraptera*, and Evaluation of the mutagenic potential of molluscicidal extracts and isolates. *J. Nat. Prod.* 2013; 56(11):1872-1877.
9. Khalifa AM. Study of the chemical constituents of *Sesbania aegyptiaca* (sesban) as a molluscicidal and livestock food. Monoufia University, Menoufia. *World J Urol.* 2010; 30:31-38.
10. Hamad MM. Phytochemical and biological studies of some plants of family's solanaceae and pittospora as molluscicides. *Cairo University, Cairo.* 2012; 14, 3-11.

- 11.Lema A. A preliminary report on the molluscicidal property of Endod
(*Phytolacca dodecandra*) *Ethiopia Medical Journal*. 2013; 123:187–190.
- 12.Maillard M, Adewunmi CO, Hostettmann K. New triterpenoid N-
Acetylglycosides with molluscicidal activity from *Tetrapleura tetraptera*.
Helvetica Chimica Acta. 2009; 72:668–674.
- 13.Sampath KP, Debjit B, Biswajit C, Pankaj T. *Allium cepa*: A traditional
medicinal herb and its health benefits *J. Chem. Pharm. Res*. 2010; 2(1): 283-
291
- 14.Handa SS, Khanuja SPS, Longo G,Rakesh DD. Extraction Technologies for
Medicinal and Aromatic Plants, Italy: *United Nations Industrial Development
Organization and the International Centre for Science and High Technology*.
2008; (1st Edn), No. 66.
- 15.Nesti FS, Wilmar M, Alice V. Toxic activities of hexane extract and column
chromatography fractions of rodent tuber plant (*Typhonium flagelliforme*
Lodd.) on *Artemia salina* Indones. *J. Agric. Sci*. 2013; 14: 1: 1-6
- 16.Kanchan L, Singh DK, Singh VKS. Characterization of the molluscicidal
activity of *Bauhinia variegata* and *Mimusops elengi* plant extracts against
the *fasciola* vector *lymnaea acuminata*Rev. *Inst. Med. Trop. S. Paulo*.
2012; 54:.3
- 17.Stirewalt MA. Effect of snail maintenance temperatures on development
of *Schistosoma mansoni*, *Experimental Parasitology*. 2004: 3(6):504-516
- 18.World Health Organization. Molluscicide screening and evaluation. Informal
meeting of investigators on molluscicide screening and evaluation held during
1965; 17-21 November 1964. Geneva.
- 19.Azare BA, Okwute SK, Kela SL. Molluscicidal activity of crude water leaf
extracts of *Alternanthera sessilis* on *Bulinus* (Phy) *globosus* *African Journal
of Biotechnology*. 2007; 6(4):441-444.
- 20.Rawi SM, El-Gindy H, Haggag AM, Elhassan AA, Kader AA. New possible
molluscicides from *Calendula micrantha officinalis*and *Ammi majus*plants.
Physiological effect on *Biomphalaria alexandrina*and *Bulinus truncatus*.
Journal of Egyptian Geriatrics Society Zoology. 2011; 16, 69-75.
- 21.Singh VK, Singh DK. Characterization of *allicin*, as a molluscicidal agent in
Allium sativum (garlic). *Journal Biol. Agricul. Horticul*. 1995; 12:119.

- 239 22. Clisiane CS, Silvan SA, André LMS, Elis CVA, Antônio SD, Nicole P. D. et al.
240 Evaluation of the toxicity and molluscicidal and larvicidal activities
241 of *Schinopsis brasiliensis* stem bark extract and its fractions. *Revista*
242 *Brasileirade Farmacognosia*. 2014; 24,298-303.
- 243 23. Babeet ST, Rekha V. Phytochemical evaluation and molluscicidal activity of
244 *Andrographis paniculata* *Acta Trop*. 2010; 56: 4
- 245 24. Singh DK, Agarwal RA. Correlation of the anticholinesterase and molluscicidal
246 activity of the latex of *Euphorbia royleana* Bioss on *Lymnaea acuminata*. *J Nat*
247 *Prod*. 1984; 47:702-5.