

**Comparative Toxicity effect of Methanolic and petroleum ether extracts of Some Ethnobotanicals Against *Anopheline* (Diptera: Culicidae) in the North Eastern Nigeria**

**ABSTRACT**

Owing to the adverse effect of synthetic pesticides, there has been increasing need to search for natural and environment friendly pesticides of plant origin as agents for control of vector of mosquito-borne diseases. This study was aimed at investigating the larvicidal activities of North eastern ethnobotanicals against vector of mosquitoes-borne diseases. Five mosquito repellent plants used by the indigenous people of north eastern, Nigeria were analysed for their effectiveness against *Anopheline* mosquitoes. The methanol and petroleum ether were analysed for their phytochemical properties. The methanol and petroleum ether extracts were investigated for phytochemical compounds with *larvicidal* activities against *anopheline* 3rd instar larvae through using deferent extracts of methanol and petroleum ether of different concentration that ranged from 50-200ppm. The *larvicidal* were observed after every six hours for total period of 36hours. The extracts of *Hyptis suaveolens* and *Azadirachta indica* of both methanol and petroleum ether extracts showed high toxicity effects against *anopheline*. The test plants contained phenol, alkaloids, flavonoids, tannins, *azadrachtin*, glycosides saponins, terpenoids and steroids.

In conclusion, the present plant extracts have potentials for development of new and safe control products for mosquitoes. As natural occurring insecticides, these plants derived materials that could be useful as an alternative for synthetic insecticides.

**Key words: Larvicidal, Insecticidal, Ethnobotanicals, Anopheline and Phytochemical Analysis**

**INTRODUCTION**

Mosquito (*Diptera: Culicidae*) is a family of small, midge-like flies which are considered by World Health Organisation (WHO), as the most dangerous insect pest to man (1). There are many types of mosquitoes living in the tropical and sub-tropical regions of the world, such as *Aedes*, *Anopheles* and *Culex* (2). Female mosquitoes are generally considered as blood-eating pests. According Michigan Mosquito Control Organization (MMCO), only few mosquito species are harmless or even useful to humanity, but most are considered as a nuisance, or even deleterious human. The female feeds on blood and in the process, they transmit extremely harmful human and livestock mosquito-borne illnesses (3). Mosquito-borne illness are diseases caused by bacterial, viruses or parasites transmitted by mosquitoes. World Health Organization (WHO) reported that the diseases transmitted by mosquitoes include: malaria,

dengue, *filariasis*, *zika virus chikungunya*, yellow fever, encephalitis (4). Mosquito-borne diseases contribute significantly to morbidity and mortality in most tropical countries. Despite the advances in techniques and products used for their control, mosquito continue to pose serious public health problems and some of the methods are compounding more problems due to environmental pollution that has a connection to human health hazard. Various products used for mosquito control have varying degree of effectiveness especially chemicals which are hazardous to environmental ecosystem and human being. The chemical products are costly and sometimes not available in some interior villages. Result of mosquito repellent based on chemicals have remarkable profile, but some are toxic against the skin, nervous system, eye irritation, nasal irritation and some result to worse problem such as brain swelling in children, anaphylactic shock, low blood pressure (5). Insects repellent of plant origin has minimal environmental effects and as well as provide a means of personal protection that could go a long way in reducing burden due to mosquito and environmental pollution. The use of plant derivatives in mosquito protection instead of chemicals could reduce the costs and environmental health effects (6). The obvious benefits would provide public health protection and an environmentally safer alternative. The significant effects could be worldwide because of the existence of these types of plants in countries where production could be possible done simply and economically to provide a viable form of protection against disease vectors (7). The inference derived from this study will go a long way in reducing burden due the pesticide and burden due to the mosquito borne disease.

#### **EXTRACTION OF ORGANIC MOLECULES**

Soxhlet apparatus were used for the plant extraction. The various plants extract for both methanol and petroleum ether extracts were concentrated using water bath which removes the methanol and hexane component leaving behind only the components of the various extracts, which were used for toxicity bioassay (8).

## 65 **Phytochemical Screening**

66 This was carried out using the procedure described by Evans (9) and Sofowora (10). Test  
67 were carried out on *Azadirachta indica* (seed, Leaf and stem), *Eucalyptus globulus* (leaf),  
68 *Occimum Kilimanscharicum* (whole plant), *Citrus senensis* (orange peels), samples to detect  
69 the presence or absence groups of phytochemical compounds such as alkaloids, glycosides,  
70 flavonoids terpenoids, saponins, phenols, Azadrachtin, steroids, and tannins

## 71 **TOXICITY BIOASSAY**

72 Third instar larvae of anopheline were collected from the rice fields and some natural water  
73 bodies in Yola, Adamawa State, Nigeria. Larvicidal bioassay were also carried out in  
74 insectary prepared for the course of this study in Yola, Adamawa state.

75 The larvicidal activities of these extracts were evaluated in static Bioassays. The method  
76 adopted from that described by World health organisation (WHO), (11). One millilitre of  
77 various plant extracts was measured and emulsified with 3 drops of Tween <sup>+</sup>80 from a needle  
78 tip. The emulsified was made up to 1 litre with distilled water to form 1000ppm stock  
79 solutions. For all the stock solutions, serial concentration was prepared. The ranges start from  
80 50ppm-200ppm. From each concentration, 250ml of all extracts was measured and introduce  
81 into separate labelled 500ml of specimen bottles. Twenty 4<sup>th</sup> instars larvae of Anopheline  
82 mosquitoes were introduced to each beaker. Each treatment has five replicates. Mortality  
83 served as the end point of the test and result were used to determine the lethal concentration  
84 (LC50), of the various plant extracts. The LC50 is defined as the lethal concentration of the  
85 bioactive extracts that kills 50% of the test species. Larvae was considered death if there is no  
86 moving or no respond to gentle probing with a fine glass rod) three times, 10 second each.  
87 Mortalities were recorded at after 36 hours for the various plant extracts and the control (only  
88 distilled water).

89

## 90 Data Analysis

91 Data obtained from this investigation were subjected to analysis using Duncan Multiple  
92 Range Test for analysis of variance. the lethal concentrations (LC50 and LC90) of the  
93 extracts were determined by Probit analysis (12) using SPSS Version 17

## 94 RESULT

95 The result of the **Table 1**, showed that out of the three hundred (300) people interviewed  
96 about ethnobotanicals that were being used as agent of mosquito repellent within their  
97 localities, 145(48.3%) of people interviewed showed that *Hyptis suaveolens* was being used  
98 within their localities, as mosquito repellent. Followed by *Eucalyptus globulus* 60(20%) and  
99 *Azadirachta indica* 45(15%). *Citrus senensis* 20(6.67%) shows to be the most unpopular  
100 plant for mosquito repelling in the region. The result also shows that *Hyptis suaveolens* has  
101 the highest percentage (58.3%,  
102

103 **Table 1: Percentage of plants in use per state**

State	<i>H. suaveolens</i>	<i>C.senensis</i>	<i>O. kilimanscharicum</i>	<i>E. globulus</i>	<i>A. inndica</i>
Location	N0. (%)	No. (0)	No. (%)	No. (%)	No. (%)
Adamawa	35(58.3)	0(0)	10(16.6)	10(16.6)	5(8.3)
Borno	10(16.6)	0(0)	0(0)	10(16.6)	40(66.6)
Bauchi	40(66.7)	0(0)	20(33.4)	0(0)	0(0)
Gombe	30(50)	15(25)	0(0)	15(25)	0(0)
Taraba	30(50)	5(8.3)	0(0)	25(42.7)	0(0)
Total	145(48.3)	20(6.67)	30(10)	60(20)	45(15)

104  
105 66.67%, 50% and 50%) of plant used against mosquitoes in four states of North eastern  
106 Nigeria that includes Adamawa, Bauchi, Gombe and Taraba respectively, followed by

*Eucalyptus globulus* (41.67%, 25% 16.7%, and 16.7%) of Taraba, Gombe, Adamawa, and Borno respectively. The locals in Borno State used *Azadirachta indica* (66.67%) more than any other plant products in the area as mosquito repellent.

#### **Qualitative Phytochemical Analysis**

In this study, primary metabolites of different solvent extracts of some ethnobotanical used against mosquito in North eastern Nigeria, were analysed qualitatively (**Table 2**). The extracts of different plants showed diverse phyto-profile with reference to solvents. Methanol and petroleum ether were used as solvents for the extraction. Methanol extracts demonstrated maximum occurrence of phyto- constituents (54/63) such as *flavonoid*, *Saponins*, *glycosides*, *tannins*, *terpenoids*, *steroids*, *alkaloids*, *Azadirachtin* and *phenol*. The Extracts derived from neem fruits showed presence of *glycosides*, *alkaloids*, *saponins* and *flavonoids* while *steroid*, *tannin*, *phenol* and *terpenoid* were absent. In the case of petroleum ether extracts, the result showed low occurrence of phyto-constituents (39/63) and absence of *terpenoids* were observed in all the neem plant extracts. Methanol extracts, of orange peels (8), *Hyptis suaveolens*, (8), *O. killimanscharicum* (8), *E. gloubulus* (8) and neem leaves (8) showed all the metabolites under observation are present.

**Table 2: Qualitative Phytochemical Analysis**


Sample(M/P)	Phen	Terp	Aza	Alk	Sap	Flav	Glyc	Ster	Tan	Total(M)	Total(P)
Neem seed	-	-	+	+	+	+	+	-	-	5	
	-	-	+	+	+	-	+	-	-		4
Neem stem	+	+	+	+	+	+	-	-	+	8	
	+	-	+	+	+	+	-	-	+		6
Neem leaf	+	+	+	+	+	+	+	+	+	9	
	+	-	+	+	+	+	+	+	+		8
O.kilimanscharicun	+	+	-	+	+	+	+	+	+	8	
	+	+	-	+	-	+	+	+	+		7
Orange peels	+	+	-	+	+	+	+	+	+	8	
	-	+	-	+	+	+	+	-	-		5
<i>Hyptis suaveolens</i>	+	+	-	+	+	+	+	+	+	8	
	-	+	-	+	-	+	+	+	-		5
<i>E. globulus</i>	+	+	-	+	+	+	+	+	+	8	
	+	-	-	+	-	+	+	-	+		5
Grand total										54/63	39/63

Keys: P. petroleum ether extract, M. methanol extracts, Phen= phenol, Glyc= glycosidae, Tan= tanins, Ster= steroids Terp= terpenoids, Flav =flavonoids, Alk= Alkaloids and sap =saponins.  
 Note: Total observation made was 63 for the 7 sample plants used for methanol extract and the same observation were made for petroleum ether extracts

## Effects of solvent used in extraction of extracts on *Anopheline* 3<sup>rd</sup> instar larvae.

Figure I, shows that all the treatment agents of methanol and petroleum ether extracts showed high significant differences to control against *anopheline*. The control showed 0 mortality at all stages during the experiment. In general, LC90 of methanol extracts (181.94, 239.29, 200.58, 145.40 and 160.70), of neem seed, *O. kilimanscharicum*, orange peels, *H. suaveolens* and *E. gloubulus* respectively, are more effective than LC90 of petroleum ether extracts (737.41, 435.747, 384.42, 251.56 and 426.03) of neem seed, *O. kilimanscharicum*, orange peels, *H. suaveolens* and *E. gloubulus* respectively. Neem stem (755.09ppm,) and neem leaf (2484.34ppm) of methanol extracts showed low toxicity effects against 3rd instar larvae of *anopheline* when compared to petroleum extracts LC90 of neem stem (478.95ppm) and neem leaf (516.01) respectively.

The LC50 methanol extracts (63.91, 77.76, 66.88, 53.07 and 56.67ppm) of neem seed, *O. kilimanscharicum*, orange peels, *H. suaveolens* and *E. gloubulus* respectively, are the concentration of methanol extracts that killed 50% of the 3rd instar larvae of *anopheline* larvae that were exposed to these concentrations. These results showed higher effectiveness against 4<sup>th</sup> instar of *anopheline* than the LC50% petroleum ether extracts (193.20, 126.86, 111.18, 79.67 and 123.60) of neem seed, *O. kilimanscharicum*, orange peels, *H. suaveolens* and *E. gloubulus* respectively.



**Figure 1: Effects of solvent used in extraction of extracts on *Anopheline* 3rd instar larvae**

## DISCUSION

Ethnobotany are well known as the study of region's plants and their practical uses through the traditional knowledge of local culture and people. The health benefits of ethno-botanical are mainly accounted to the presence of many active phytochemical constituents in various parts of these plants. Present study was conducted with an objective to identify the best extraction solvent, which can be used to obtain the maximum amount of the phytochemicals from the shade dried ethnobotanicals used against mosquitoes in the North eastern Nigeria.

The present study, started with ethnobotanicals survey in north eastern Nigeria, and investigations revealed that *Hyptis suaveolens*, *Eucalyptus globulus*, *Citrus senensis* and *Ocimum kilimanscharicum*, were popularly used, by some local populace. The present study shows that *Hyptis suaveolens* [145 (48.3%)] proved to the most popular plant in the region that has been used as mosquito repellent. The findings of this Present study agree with the works of. (13) and. (14). The result of the survey shows that *Hyptis suaveolens* has the



highest percentage (58.3, 66.67, 50.0, 50.0%) of plants that are used in Adamawa, Bauchi, Gombe and Taraba respectively, followed by *Eucalyptus globulus* (41.67%, 25% 16.7%, and 16.7%) of Taraba, Gombe, Adamawa, and Borno respectively. The locals in Borno prefers using *Azadirachta indica* (66.67%) more than any other plants product in the area as mosquito repellent. Their preference is based on how it worked for them.

Results have shown that all the effect of the treatment agents of methanol and petroleum ether extracts used against anopheline larvae mosquitoes are significantly different from each other. The LC50 methanol extracts (63.91, 77.76, 66.88, 53.07 and 56.67ppm) of neem seed, *O. kilimanscharicum*, orange peels, *H. suaveolens* and *E. gloubulus* respectively, showed higher larvicidal effect than petroleum ether with LC50 (193.20, 126.86, 111.18, 79.67 and 123.60). LC90 values of methanol extracts (181.94, 239.29, 200.58, 145.40 and 160.70), of neem seed, *O. kilimanscharicum*, orange peels, *H. suaveolens* and *E. gloubulus* respectively, are also more effective than LC90 of petroleum ether extracts (737.41, 435.747, 384.42, 251.56 and 426.03) of neem seed, *O. kilimanscharicum*, orange peels, *H. suaveolens* and *E. gloubulus* respectively. This agrees with the reports of (15) and (16) that showed methanolic extracts as the best treatment agent against many mosquito species than some of the extracts from other solvents. This may be attributed to high polarity effect of methanol than the petroleum ether as reported by (17)

In the case of Neem stem (755.09ppm,) and neem leaf (2484.34ppm) of methanol extracts, opposite is the case, they showed low toxicity effects against 3<sup>rd</sup> instar larvae of *anopheline* when compared to petroleum extracts with LC90 of neem stem (478.95ppm) and neem leaf (516.01) respectively and this is contrary to the reports of (16) and (15), that reported higher larvicidal effect of methanolic extracts than the petroleum ether extracts but is in agreement with the report of Komalamisra et al. (18) where petroleum ether extracts of some Thai plants that showed LC50 values between 11.2 and 18.84mg/L which are far better than the methanol

extracts that showed LC50 values between 13.2 and 45.2mg/L. The difference in the amount phytochemical constituents extracts of different plants by different solvent may depend on the physiology and the polarity effect of the individual solvents used.

## CONCLUSION

The extracts exhibited *larvicidal* effects on the 3<sup>rd</sup> instar larvae exposed to them at different concentration. The *larvicidal* effects are concentration dependent. Generally, *Hyptis suaveolens* extracts proved to be most effective treatment agent used, followed by neem seed extract and *Ocimum kilimanscharikum*. The most ineffective treatment agent observed was neem leaves and orange peels extracts. This research may serve as scientific basis lend credence to the claim by the local populace that this plants material has some metabolites that mosquitoes are comfortable with it which causes their repellence. It justifies the claim that the selected plants are efficacious in the management of mosquito populations. It also concludes that potency of these plants is depended on the solvent used for the extraction and the dose administered.

## REFERENCES

1. World Health Organisation (2015). Malaria facts sheets N° 94
2. Donald, J. (2004). Mosquitoes. Associate research professor in entomology [on line] Available: <http://www.rei-rutgers.edu/~insects/moslife.htm>.
3. Michigan Mosquito Control Organization, (2013). Michigan and Mosquitoes- their Biology and Control.
4. Caraballo, H. (2014). Emergency department management of mosquito-borne illnesses. Malaria dengue and West Nile virus. *Emergency Medicine Practice* 16 (5): 1-5
5. Patel E. K., Gupta A., Oswal R.J. (2012). A review on: Mosquito repellent methods. *International of pharmaceutical, Chemical and Biological Sciences*. **2**: 310-317

- 219 6. Rahuman, A. A., Bagavan, A., Kamaraj, C., Saravanan, E., Zahir A. A., Elango G.  
220 (2009). Efficacy of larvicidal botanical extracts against *Culex quinquefasciatus* Say  
221 (Diptera: Culicidae) Parasitology research 104 (6):1365-1372.
- 222 7. Marta F. M. and Sarah J.M. (2011). Plant based insect's repellents: a review of their  
223 efficacy, development and testing. malarial journal.
- 224 8. Amusan A.A.S, Idowu A.B., Arowolo F.S. (2005): Comparative Toxicity effect of  
225 bush tea leaves (*Hyptis suaveolens*) and orange peel (*Citrus sinensis*) oil extract on  
226 larvae of the yellow fever mosquito *Aedes aegypti*.
- 227 9. Evans W. C. (2002). Treese and evans Pharmacognosy. 15<sup>th</sup> edition. W. C. saunders  
228 company Limited 135-150
- 229 10. Sofowara, A. (2006). Medicinal plants and Traditional Medicine in Africa. Spectrum  
230 Books Limited Ibadan, 150- 156
- 231 11. World Health Organisation (2005). Guidelines for Laboratory and field testing of  
232 mosquito larvicides. World Health Organisation Communicable Disease Control,  
233 Prevention and Eradication. *Who pesticides Evaluation Scheme*  
234 WHO/WHOPES/GDP,1-  
235
- 236 12. Finney, D. J. (1971). Probit analysis. Cambridge University Press. United Kingdom 66-  
237 72.  
238  
239
- 240 13. Egunyomi A., Gbadamosi I.T., and Osiname K.O. (2010). Comparative effectiveness  
241 of ethnobotanical mosquito repellents used in Ibadan, Nigeria. *Journal of*  
242 *Applied Biological Sciences*.36: pp.2383 -2388.  
243
- 244 14. Kweka, D. Elimingaya, Franklin.Mosha, Asanterabi Lowossa, Aneth M.M., Jovin K.,  
245 Emmanuel F., Esther E.L., Johns M., Michael, J. M., Charlse P. M., Filemoni, T.,  
246 Michael A. M., Rajabu, M., Grace C., Michael A. M., and Emmanuel A. T. (2008).  
247 Ethnobotanicals study of some mosquito repellent plants in the north eastern  
248 Tanzania. *Malaria Journal* 7: 152
- 249 15. Dixon D., Jeena G, (2017). Comparison of different solvents of phytochemical  
250 extraction of Potentials from *Datura metel* plant leaves. *International Journal of*  
251 *Biological Chemistry* 11:17-22.
- 252  
253 16. Kadri, A., Felhi., Daoud, A., Hajloui, H., Mnatgui, K. Gharsalah, N. (2017). Solvents  
254 extraction effect on phytochemical constituent's profiles, antioxidant and anti-  
255 microbial activities and functional group analysis of *Ecballium elaterium* seeds and  
256 peels fruits. *Food Science and Technology* 37(3): 483-492
- 257 17. Felhi, S., Baccouch, N., Ben Salah, Smaui, S., Alluche, N., Gharsallah N., Kadri A.  
258 (2016a). Nutritional constituents' phytochemical profiles, invitro antioxidant and anti-  
259 microbial properties and gas chromatography-mass spectrometry analysis of various  
260 solvents and extracts from grape seeds (*vitis vinifera* L). *Food science and*  
261 *bacteriology* 25(6): 1537-1544

- 262 18. Komalamisra, N., Trongtokit, Y., Rongsriyam, Y. and Apiwathnasorn, C.  
263 (2006). Screening for larvicidal activity in some thai plants against four  
264 mosquito vector species southeast Asia *Journal of tropical medicine of public*  
265 *health*. 36(6):1412-1422.

UNDER PEER REVIEW