

POTENTIAL OF FIVE LOCAL PLANT AS PROTECTANT AGAINSTS HIDE BEETLE,
DERMESTES MACULATUS DEGEER INFESTING DRY CAT FISH ; *CLRIAS*
GARIEPINUS BURCHELL.

ABSTRACT

Efficacy of varying amounts of five selected local plant namely ; *Balanite aegyptiaca*, *Eugenia aromatica*, *Piper guineense*, *Ocimum gratissimum*, and *Ziziphus mauratania* against the mortality, oviposition, progeny emergence and weight loss of adults hide beetle, *Dermestes maculatus* was studied under ambient conditions. The plant were pulverized into powders and applied as 2.0, 4.0 and 6.0 g per 50 g of dried fish. Three pairs of newly emerged adults of Hide beetle were inoculated into each of the treated fish in beaker, arranged in a Complete Randomized Design (CRD) in the laboratory. The result showed all the plant powders tested evoked significant mortalities of adults at highest concentration (6) after 3 and 7 days of post treatment when compared with untreated control. The rate of oviposition was drastically suppressed by all plant powders especially at higher concentration. The number of adults emerged were significantly lower than control at ($p>0.005$). Similarly all the plants powders significantly prevented weight loss at highest concentration after 12 weeks of post treatments periods. These results have shown that tested plants demonstrated potential ability against adults hide beetle infesting dried fish.

Keywords: *Dermestes maculatus* , mortality, oviposition , Smoked fish.

INTRODUCTION

Dried fish damage in storage, transportation as well as marketing by insect pest are particularly important in Nigeria because they significantly reduce the availability of the produce as good sources of protein (Adedire and Lajide, 2000). The quantitative and qualitative loses of dried fish during storage, transportation and marketing by insect are due to infestation by beetles species of the genus *Dermestes* and *Necrobia* (Haines,1991). These cause losses ranging from 22-50% in Nigeria (Rolling and Hayward, 1963). In addition, the fish are often render unfit for human consumption and aesthetically unacceptable to the buyers (Baba *et al.*, 2014).

The control of these pest in Northern Nigeria is primarily dependent upon repeated application of synthetic chemicals such as chloripyripos, methyl, permethrin,

cypermethrin, BHC, and “Otapiapia” (Locally formulated) onto fish carton for protection against insect pest (Igene *et al.*, 1998; Agbolagba *et al.*, 2011). Although synthetic chemical are effective to protect stored dried fish, the availability, cost and right application ,constitute the major problems toward the use of synthetic pesticide by illiterate fish processors (Booke *et al.*, 2006). Hazard associated with food can result in injury and harm to human health. Millions of people world- wide suffer from sort of food poisoning each year due to uncontrolled application and abuse of synthetic chemical(Abdusalam,2015). With these demerit of synthetic chemical currently world- wide interest centered on search for alternative pesticide to stored product by the use of botanical pesticide.

Many Nigeria medicinal plants species have been cited as very important in pest control of stored grain, legumes and dried fish (Onu and Baba, 2003; Adebote *et al.*, 2006; Akinwumi *et al.*, 2006, Olayinka, 2004; Jose and Adesina, 2014 and Baba *et al.*, 2014). These provide a solution to the problem emanating from the use of synthetic chemical, thus, this study sought to search for natural preservation material that are cheap and consumers friendly. The present work is aimed at investigating the potential of five plants powders as protectant against *D. maculatus* adults.

Materials and Methods

Preparation of plant powders

The sample of *Balanite agyptiaca* (K) and *Ziziphus mauratania* (L) were obtained within the main campus of Usmanu Danfodiyo University, Sokoto, while the Fresh leaves of *Ocimum gratisimum* (L.) and dry fruit of *Piper guinense* and *Eugenia aromatic* were bought from Sokoto Central Market, Sokoto. The samples were washed and dried under shade and milled into fine powder using mortar and pestle.

They were sieved through 0.2 mm mesh size. Each of the plant powders was kept in a separate plastic container with tightly fitted cover prior to the use.

Preparation of fish sample

The samples of dried *Clarias gariepinus* was purchased from fish mongers at Sokoto Central Market, the fish samples were disinfected by heat treatment in the laboratory drying cabinet at 60°C for 1 hour and allow to cool at room temperature as adopted by (Onu and Baba, 2003).

Collection of Hide Beetle and Maintenance of insect culture

Different stages of Hide beetle were obtained from naturally infested fish collected from Sokoto central market fish stalls. Several adults pairs of *D. maculatus* were obtained and kept in glass jars and fed with dried fish (*Clarias* species), the jars were covered with Muslin cloth and tight with rubber band. Wet cotton wool were supplied in each jar to provide water requirements for oviposition as suggested by Hill (1990). The adults laid eggs which hatched into larvae and change to pupae. The pupae were picked up and transferred in to separate jar to obtained newly emerged adult, which were used for the experiment. The culture maintained for the regular supply of newly emerged adults.

Effect of plant powders on mortality, Progeny emergence and weight loss of fish by *D. maculatus*.

Fifty gram (50 g) of dried fish was dusted with 0, 2, 4 and 6g of each plant powder. Corresponding amount dried fish was dusted with 0.5g of cypermetrin (2.0%) powder. Six newly emerged adult (0-72 hrs) of *D. maculatus* were introduced into each of the beaker (500 ml), a cotton wool soaked in distilled water was provided to cope with the water requirement of adults, these beakers were covered with muslin cloth to allow gaseous exchange and prevent insect from

escaping. There were three (3) replicates. Mortality data were taken at one (1) three (3) and seven (7) days post treatment. After seven days both live and dead insect were discarded, eggs and larvae were counted. The set up was kept for the emergence of F1 adult from 35 days after treatment until all adult emerged. Weight loss due to *D. maculatus* infestation was determined at 4, 8 and 12 weeks of post treatment. Percentage weight loss was calculated as difference between the initial and final weight of fish divided by initial and multiplied by 100.

Data Analysis

Data obtained were subjected to analysis of variance (ANOVA) using general linear Model (Multivariate Produce of statistical package for Social Science (SPSS, 2007) and means found to be significant were separated using Duncan Multiple range test at 5% level of significant.

Results

Table 1, shows the mortality of adults *D. maculatus* recorded at 24hrs, 48hrs and 7days intervals due to effect of plant powder treatments. The result revealed the mortality increase with increase in amount of powder and length of exposure. There were relatively low mortality rates at 24hrs and 72hrs of post treatment. Higher mortality were recorded at 7days of post treatment period by *E. aromatic* (61.11%) and *B. aegyptiaca* (44.44%) and *P. guinense* (33.33%) at application of 6g/50g of dried fish. These were statistically significant ($P < 0.05$) compared to other test plant and control. Lower mortality of adult at highest concentration after seven days post treatment was observed in *O. gratisimum* (27.77), and *Z. mauratania* (27.77) .

Table 2, showed the effect of plant powders on the oviposition and adults emergence of *D. maculatus* on treated dried fishes. All the plant powders tested at all concentrations inhibited the number of eggs laid by *D. maculatus* these were significant different ($p < 0.05$) compared with oviposition on untreated dried fish.

Fewer number of eggs were deposited on fishes treated with *P. guinense* at all concentration and *E. aromatic* at 4.0 and 6.0 concentration. But higher number of eggs were observed on fish treated with *O. gratisimum*, *B. aegyptiaca* and *Z. mauratania* at all concentration.

It is also evident from Table 2, that percentage of larval and adult emergence in untreated dried fish was significantly higher ($p < 0.005$) than percentage of larval and adult emergence in all treated dried fishes. There was little emergence of adult (0.3%) at lower concentration (2.0g) and no emergence of adults on dried fish treated with *P. guinense* at 4.0 and 6.0g/50g of dried fish. But few emergence of adult *D. maculatus* were also observed on dried fish treated with *E. aromatica*, *B. aegyptica*, *Z. mauratania* and *O. gratisimum* at all application rate compared with control.

Table 3, shows the weight loss of dried fish treated with plant powders, it is evident from the table that the weight loss of the treated dried fish with *P. guinense*, *E. aromatic* were significantly lower ($P < 0.05$) than control at all rate of application at 4, 8 and 12 weeks of post treatment period, but there were no significant difference between the weight loss of dried fish treated with powder of *B. aegyptiaca*, *Z. mauratania* and *O. gratisimum*, at lower rate of application of 2.0g/50g of fish after 4, 8 and 12 weeks of post treatment periods

Discussion

The result obtained in this study showed that the powder of *E. aromatic* seem to be the most effective in killing adult *D. Maculatus* especially at high doses, followed by *B. aegyttica* and *P. guinense* treatment and *O. gratisimum* and *Z. mauratania* produce moderate result when compared with other tested plant powders after 7 days of post treatment period. All the powder treatment caused significant mortality of *D. maculatus* (adult) compared with control. The effectiveness of *E. aromatica*, *B. aegyptica* and *P. guinense* could be attributed to

high toxicity of these plant powders, this is consistent with the finding of others workers who have reported the effectiveness of various plant powders as dried fish and grain protectant against various insect of stored product. Ashamo and Akinneye (2004) reported that powders of *E. aromatic* and *Z. officianle* were effective against the yam moth *Euzopherodes vapidella* at 0.5g, 1.0g and 1.5g concentration within 72 hours of application producing 100% mortality of moths. Hoda *et al.* (2016) reported the insecticidal activity of *Balanite aegyptica* and *commiphora molmol* against blowfly *lucilia sericata* (Diptera: calliphoridae). Abdullahi and Muhammad (2004) assessed toxic potential some plant powders on the survival and development of *C. maculatus* and recorded 100% mortality after 8 days post treatment when *P. guinense* powder was used at the rate of 1g/50g cowpea seed. Similarly Adedire and Lajide (2000) also reported the susceptibility of different ages of *Callosobrunchus maculatus* and *D. maculatus* respectively to *P. guinense* at all rate of application. The observed low mortality of *O. gratisimum* and *Z. mauratania* could be due to low toxicity, differences in test insect, geographical location as well as environmental condition at which the study was carried out. Which correspond with report of Iloba and Ekrakene (2006) who reported effectiveness of botanical insecticide varies with species of insect, with percentage adult mortality of *C. maculatus* ranging from 83.3 to 93.4% for *Azadrachta indica* ; 86.7 to 90% for *O. gratisimim*, 93.4 to 96.6% for *Hyptis suavelons* , while *S. zeamaise* recorded mortality of 13.3 to 50% for *A. indica*, 26.7 to 50% for *O. gratismum* and 13.3 to 30% for *H. suaveolens* powders at rate of 1.5, 2.5 and 3.5/80g cowpea seed after 7 days of post treatment. Maud *et al.*, (2013) reported weak oral acute toxicity test of *O. gratisimum* aqueous extract on Wister mice.

It is observed from the result of the study that oviposition and percentage of adult emergence by *D. maculatus* were significantly lower in powder treated fish against untreated fish. Few number of eggs were oviposited by each of the beetle

on treated fish and consequently low larvae and adult emergence were observed ; this could be as a result of high adult mortality caused by the treatment before oviposition or plant powder had deleterious effect on mating, egg laying, repellence or development of immature stage which in turn result in lower adult emergence. This is agreement with findings of Donpedro (1985) who reported that the orange peel powder reduced progeny development of *D. maculatus*. Asawalam and Emosairue (2006) reported that 0.1g of *P. guinense* powder at 50g of maize grain result in significant reduction of adult emergence. Ofuya and Dawodu (2001) had reported when *P. guinense* powder was applied at five rate (0.1, 0.2, 0.3, 0.4, and 0.5g/20g of cowpea seeds) results in significant reduction in oviposition and adult emergence of *Callosobrunchus maculatus*. Ajayi,(2015) reported plant product obtained from clove (*S. aromatica*), West African black pepper (*P. guinense*), alligator pepper (*Aframomum melegueta*). Ethiopian pepper (*X. aethiopica*) and dried rhizome of ginger (*Z. officinale*) plant powder significantly reduce oviposition of *S. zeamais* compared with control and the powder of West African black pepper and clove at all dosage rate completely inhibited progeny emergence.

The result of the investigation reveals all the plant powder used at different amount cause considerable reduction in weight loss of dried fish infestation. There was significant ($p<0.05$) reduction in percentage weight loss of smoked fish treated with powder of *P. guinense* and *E. aromatic* compared to other plant powder tested and untreated fish. The reduction in weight loss reflect the number of surviving larvae and adult emerged from respective treated fish. The highest weight loss was recorded in untreated (control) this could be as a result of feeding activities of *D. maculatus* larvae, on the fish which subsequently cause highest weight loss. The result confirmed the earlier report of Akob and Ewete (2007), that ash and some bioactive plant species cause oviposition deference or ovicidal action resulting in reduced progeny production of stored product insect that reduce seed

damage(weight loss). Fasakin (2003) reported that percentage weight Loss of smoked fish Infested by *D. maculatus* was significantly reduced in fish treated with oil extracts from *A. melegueta*, *P. guinense* and *M. myristica* to 4.43, 3.72 and 8.46% respectively. Adesina *et al.*, (2012), reported that treatment of maize with *Secamone afzellei* Plant powder against *Sitophilus zeamais* at 2.0g/ 20g seeds significantly suppress Adult emergence and percentage weight Loss were significantly reduced at all level of application i.e 0.5/20g, 1.0/20g, 1.5g/20g 2.0g/ 20g of maize seeds.

Conclusion

A study was conducted to evaluate the effect of five plant powders on mortality of adults, oviposition, adult emergence and weight loss of the treated fish by *D. maculatus* infestation. The five plant species, namely; *Balanite agyptiaca*, *Eugenia aromatic*, *Piper guinense*, *Ocimum gratisimum* and *Ziziphus mauratania*. Result of the study revealed that all the plant powders tested were effective in control of *D. maculatus*. The powder of *Eugenia aromatica*, *Balanite agyptiaca* and *P. guinense* recorded the higher mortality of adult and the powders of *O. gratisimum* and *Z. mauratania* gave the least mortality of adult *D. maculatus*. However, the powdered of *P. guinense* and *E. aromatica* were found to result in lowest rate of oviposition, adults emergence and weight loss of the treated fish. The above mention finding revealed that powdered that powders of *P. guinense* and *E. aromatica* could be use as grain protectant against *D. maculatus*. Therefore, it is recommended the use of these powder for control of *D. maculatus* infestation during processing, storage; transportation and market of smoked dried fish .

226

227

228

229

230

231

232 Table 1: Percentage mean mortality of adult *D. maculatus* on dried fish treated with plant powders.

Plant Powders	Concentration (g)/ (50 g fish)	Post treatment period in Days		
		1 day	3 days	7 days
<i>B. aegyptiaca</i>	2	0.00 ^c ± 0.00	5.55 ^{bc} ± 5.55	11.11 ^{de} ± 5.56
	4	0.00 ^c ± 0.00	11.11 ^{bc} ± 5.55	11.11 ^{de} ± 5.55
	6	0.00 ^c ± 0.00	27.77 ^{bc} ± 5.56	44.44 ^{bc} ± 5.56
<i>E. aromatica</i>	2	11.11 ^{bc} ± 1.11	11.11 ^{bc} ± 1.11	22.22 ^{cde}
	4	16.66 ^{bc} ± 0.96	22.22 ^{bc} ± 1.11	27.77 ^{cde}
	6	22.22 ^b ± 5.5	33.33 ^b ± 0.96	61.11 ^{ab}
<i>O. gratisimum</i>	2	00.00 ^c ± 0.00	00.00 ^c ± 1.11	00.00 ^c ± 0.00
	4	00.00 ^c ± 0.00	00.00 ^c ± 1.11	11.11 ^{de} ± 5.55
	6	5.55 ^b ± 5.55	16.66 ^{bc} ± 0.96	27.77 ^{cde} ± 5.56
<i>P. guinense</i>	2	00.00 ^c ± 0.00	5.55 ^{bc} ± 1.11	11.11 ^{de} ± 0.00
	4	55.55 ^{bc} ± 5.55	16.66 ^{bc} ± 0.96	27.77 ^{cde} ± 5.56
	6	16.66 ^{bc} ± 0.96	16.66 ^{bc} ± 0.96	33.33 ^{cd} ± 9.6
<i>Z. mauratania</i>	2	5.55 ^{bc} ± 5.55	5.55 ^{bc} ± 5.55	11.11 ^{de} ± 5.55
	4	5.55 ^{bc} ± 5.55	11.11 ^{bc} ± 5.55	16.66 ^{cde} ± 0.00
	6	16.66 ^{bc} ± 9.6	16.66 ^{bc} ± 9.6	27.77 ^{cde} ± 5.55
Cypermeterin	0.5g	55.53 ^a ± 5.55	72.22 ^a ± 4.69	72.22 ^a ± 4.69
Control	0.00	0.00 ^e ± 0.00	0.00 ^e ± 0.00	0.00 ^e ± 0.00

235 Table 2: Effect of plant powders treatment on oviposition and f1 adult emergence

Plant powders	Concentration (50g fish)	Oviposition	Number of larvae emerged	F1 adults
B. aegyptiaca	2	82.33c ± 4.48	65.53 ^c ± 5.72	17.85 ^{ed} ± 0.81
	4	53.33fgh ± 2.60	43.12 ^{ef} ± 2.01	14.15 ^{de} ± 1.11
	6	51.67gh ± 4.48	41.88 ^f ± 5.72	5.53g ± 0.81
E. aromatica	2	61.00 ^{efg} ± 0.57	51.28 ^{de} ± 1.99	11.38 ^f ± 1.11
	4	16.33 ⁱ ± 0.88	12.25 ^g ± 0.49	3.08 ^{gh} ± 1.63
	6	4.66 ^{ij} ± 2.60	3.70 ^{gh} ± 1.99	1.23 ^{gh} ± 0.62
O. gratisimum	2	95.00 ^b ± 1.52	79.76 ^b ± 1.03	22.77 ^b ± 2.4
	4	79.00 ^{cd} ± 1.73	62.67 ^c ± 3.21	14.15 ^{de} ± 1.23
	6	42.67 ^h ± 2.72	35.61 ^f ± 2.48	13.54 ^{de} ± 3.2
P. guinense	2	10.33 ^{ij} ± 0.33	6.55 ^{gh} ± 0.28	0.31 ± 0.31
	4	6.66 ^{ij} ± 2.96	5.69 ^{gh} ± 2.53	0.00 ^h ± 0.00
	6	0.00 ^j ± 0.00	0.00 ^h ± 0.00	0.00 ^h ± 0.00
Z. mauratania	2	70.00 ^{de} ± 1.15	56.98 ^{cd} ± 3.47	18.77 ^{bc} ± 1.34
	4	64.00 ^{de} ± 3.00	52.71 ^d ± 2.80	16.62 ^{cde} ± 3.70
	6	62.00 ^{ef} ± 4.04	51.85 ^{de} ± 1.02	112.92 ^{ef} ± 1.59
Cypermethrin	0.5g	9.66 ^{lj} ± 7.31	4.7 ^{gh} ± 3.6	0.00 ^h ± 0.00
Control	0.00	117 ^a .00 ± 1.52	93.73 ^a ± 3.70	29.33 ^a ± 0.33

236

237

238 Table 3: Effect of plants powders on weight loss of dried fish due to *D.maculatus* infestations

Plant powders	Concentration/(50g fish)	Post treatment Period (Day)		
		30	60	90
<i>B.aegyptiaca</i>	2	28.23 ^e ± 1.51	31.27 ^{ab} ± 1.44	31.97 ^{abc} ±0.90
	4	21.46 ^{cd} ±1.68	23.32 ^{de} ± 1.1	24.49 ^{de} ± 0.53
	6	18.82 ^d ± 6.1	22.05 ^e ± 1.08	23.04 ^e ± 6.4
<i>E.aromatica</i>	2	22.27 ^{cd} ±2.29	26.97 ^{cde} ±2.06	29.85 ^{abcd} ±0.94
	4	12.65 ^e ±1.97	13.65 ^f ±2.60	14.34 ^f ±2.06
	6	10.36 ^e ±0.77	11.53 ^{fg} ±0.62	27.89 ^{bcd} ±2.52
O.gratisimum	2	22.27 ^{cd} ±2.29	26.97 ^{cde} ±2.06	29.85 ^{abcd} ±0.94
	4	12.65 ^e ±1.97	13.65 ^f ±2.60	14.34 ^f ±2.06
	6	10.36 ^e ±0.77	11.53 ^{fg} ±0.62	27.89 ^{bcd} ±2.52
<i>P. guinense</i>	2	11.44 ^e ±0.54	12.9 ^{fg} ±0.24	13.13 ^f ±0.21
	4	8.77 ^e ±1.532	10.31 ^{fg} ±1.06	12.50 ^f ±0.42
	6	7.89 ^e ±0.33	8.06 ^g ±0.26	8.11 ^F ±0.21
<i>Z. mauratania</i>	2	29.27 ^{ab} ±0.22	32.44 ^a ±1.01	32.84 ^{ab} ±0.48
	4	26.51 ^{abc} ±2.61	26.42 ^{cde} ±2.32	28.51 ^{abcde} ±2.74
	6	23.32 ^{bcd} ±0.57	25.00 ^{cdc} ±0.46	26.31 ^{cde} ±0.63
<i>Cypermtrin</i>	0.5	8.15 ^e ±0.86	11.07 ^{fg} ±0.57	11.14 ^e ±0.60
<i>Control</i>	0.00	29.22 ^{ab} ± 1.50	32.29 ^a ± 2.28	34.58 ^a ±2.03

REFERENCES

- Abdullahi, Y. M. and Muhammad, S. (2004). Effects of Some Plant Powders on Cowpea Weevil *Callosobrunchus maculatus* During Storage. *African Journal of Biotechnology*, **5** (10): 901 – 906.
- Abdulsalam, A. S. (2015). Expert Warns on dangers of pesticides. *Daily Trust*(Tuesday, August, 25, 2015).
- Abolagba, O. J., Igene, J. O. and Usifoh, C. O. (2011). Studies of Pesticide Residues in Smoked Cat Fish (*Clarias gariepinus*) in Nigeria: Some Health Implication. *Australian Journal of Basic and Applied Science*, **5** (5): 496-502.
- Adebote, D. A., Abolude, D. S., Oniye. S. J., Oladodo, S. S. and Hassan, M. M. (2006). Larvicidal and repellent action of *Detarium microcarpum* seed oil against the larvae of *Dermestes lardarius* (Coleopteran: Dermestidae) in dried *Clarias gariepinus* fish. *Journal of Entomology* **38**(3)245-253
- Adedire, C. O. and Lajide, L. (2000). Effect of pulverised plant materials on fish damage and growth performance of fish beetle. *Dermestes maculatus* (Degeer). *Journal of Entomology*. **32**:215-218.
- Adesina, J. M., Ofuya, T. I. and Afolabi, L. A. (2012). Insecticidal Activity of *Secamone afzelli* (Schult) K. Schum Powder in the Control of *Stiphilus Zeamais* (Mots) (Coleopera, Curculianidae) *J. of Agricultural Technology* **8** (1) 117-124.
- Ajayi, F. A. (2015). Insect, Plants and Humanity; the Organic Agriculture and Stored Products Protection Axis, 10th in *Augural Lecture, Nasarawa State University, Keffi*, Onaivi Printing and Pub. Co. Ltd, Keffi Nasarawa State; Nigeria.
- Akinwumi, F. O., Fasakin, E. A and Adedire, C.O. (2006) Progeny inhibiting effect of four plant product against the leather beetle and copra beetles on smoked African mud cat fish *Clarias gariepinus* Burchell . *Journal of Biological Science*. **6**(6): 1023 -1028,
- Akob, C. A. and Ewete, F. K. (2007). The Efficacy of Ashes of Four Locally Used Plant Materials Against *Sitophilus Zeamais* (Coleoptera Curculionadae) in Cameroon. *International Journal of Tropical Insect Science* **27** (1): 21 – 26.
- Asawalam, E.F. and Emosairue, S. O. (2006). Comparative efficacy of *Piper guinense* Schum and Thonn and Primiphos Methyl on *Sitophilus zeamais* (Motschulsky). *Tropical and Subtropical Agroecosystem*; **6**:143-148.

271 Ashamo, M. O. and Akinneye, J. O. (2004). Toxicity of Powder of some Tropical Plants Against
 272 the Yam Moth *Eutrophedes vapidella* Mann. (Lepidoptera: Pyralidae), *Nigerian Journal of*
 273 *Experiment Applied Biology*, 5:63 – 68.

274 Baba, G. O., Ayandande, F. A., Erhabor, T. A. Majolagbe, M. O., Sulaiman, Y. D., Fodoyin, A.
 275 S. and Musa, K. (2014). Potential of Neem Product as Protectant Against Adult Mortality;
 276 Rate of Oviposition and Emergence of F1, Generation of *Dermestes Maculatus* Degeer on
 277 Dried Fish. *International Journal of Applied Research and Technology* 3 (4): 53-62.

278

279 Booke, S. J., Vanlon, J. J. A., Vanttuis, A., Kossou, D. K. and Dicke M. (2001). The Use of Plant
 280 Material to Protect Stored Leguminous Seed Against Seed Beetle: A Review. The
 281 *Netherlands: Backhuys Publish* P. 108.

282 Don-Pedro, K. N. (1985). Toxicity of some citrus peels to *Dermestes maculatus* Degeer and
 283 *Callosobruchus Maculatus* (F). *Journal of Stored Products Research* 25 (2): 81-86.

284 Fasakin.E. A, (2003). Use of plant oil extract as protectant against storage insect pest,
 285 *Dermestes maculatus* Degger in smoked fish: In Eyo, A. A. 18th Ann. *Conference of Fish.*
 286 *Society Nigeria Proc.* P 1-6.

287 FDF, (2005). Federal Department of Fisheries: Federal Ministry of Agriculture and Rural
 288 Development Report of Presidential Committee on *Fisheries and Aquaculture*
 289 *Development.* P 1-60.62

290 Green A. A (1967). The protection of dried Sea-Fish in south Arabia from infestation by
 291 *Dermestes frischii*, Kug. (Coleoptera: Dermestidae). *Journal Stored Product Research*, 2
 292 (4);1-50

293 Haines, C. P. (1991). Insect and Arachnids of Tropical Stored Products, their Biology and
 294 Identification. *A Training Manual Natural Research Institute, Chatham, Kent, UK*, 246P.

295 Harbone, J. B. (1984). Phytochemical Methods 2nd Edition, E'LBs Pp 12-56.

296 Hill, D. S. (1990). *Pest of Stored Products and their control.* Belhaven Press London, 74-75.

297 Hoda, S. M., Falimy, M. M., Attia, M. M., Rabab, M. E., Shalaby, H. A. and Massoud, A. M.
 298 (2016). The insecticidal activity of two medicinal plant (*Commiphora molmol* and *Balanites*
 299 *aegyptiaca*) Against Blowfly *Lucilia sericata* (Diptera: Calliphoridae) *International Journal of*
 300 *advanced Research in Biological Science*, 3 (3) 144 – 158.

301 Igene, J. O., Abolagba, O. J. and Ikeme, A. I. (1998). Use of Pesticide on Cured Fish in Nigeria.
 302 *Report and Proceeding of the Sixth FAO Expert Consultation on Fish Technology in Africa,*
 303 *Kisumu, Kenya*, 126-130.

304 Iloba, B.N. and Ekrakene, T. (2006). Comparative assessment of insecticidal effect of
 305 *Azadirachta indica*, *Hyptis suaveolens* and *Ocimum gratissimum* on *Stiphilus zeamais* and
 306 *Callosobrunchus maculatus*. *Journal of Biological Sciences* 6 (3): 626-630.

307 Jose, A. R. and J. M. Adesina (2014). Larval Susceptibility of *Dermestes Maculatus* (Degeer)
 308 (Cleoptera: Derestidae) to *Secamone Afzelli* K. Schum Leaf on Smoked Dried Fish.
 309 *International Journal of Aguaculture* Vol. 4 (17): 101, 10: 5376.
 310

311 Lale, N. E. S (2002). *Stored Product entomology and acarology in Tropical Africa*, Mole
 312 Publication (Nig.) Ltd, Maiduguri, Nigeria.

313 Maud, K. M., John, P. B., Patrick, O., Alex, K., Arop L.D., Jusua, O. O. and Mutabola, T. M.
 314 (2013). Oral Acute Toxicity of Selected Botanical Pesticide Plant Used by Substance Farmers
 315 around the Lake of Victoria Besin. *African Journal Environmental Science and Technology* Vol.
 316 7 (3) pp 93 – 101.

317 Ofuya, T. I. and Dawodu, (2001). Aspect of insecticidal action of *Piper guineense* Schum and
 318 Thorn fruit powder against *Callobrosbrunchus maculatus* (F) Coleoptera: *Bruchidae*.
 319 *Nigerian Journal of Entomology*. **19** : 40-50.

320 Olayinka, O. J. (2014). Effect of Two Eco-Friendly Insecticides (Dennetta Tripetela) and Piper
 321 Guineense) Against *Dermestes maculatus* (Degeer) on Smoked Dried Clarias Gane Pinus
 322 (Pisces: Claridae); *IOSR Journal of Environmental Science. Toxicology and Food*
 323 *Technology* (10SR-J(STFF) Vol. 8 (7): 37-43.

324 Onu, I. and Baba, G. O. (2003). Evaluation of neem products (*Azadrachta indica* (A. Juss;
 325 Melaceae) for the control of Dermestid beetle. *Dermestes maculatus* (Coleoptera
 326 Dermestidae) on dried fish. *Nigerian Journal of Entomology*, **20**: 105-115

327 Rolling. M. J. and Hayward. L. A. W. (1963). Aspect of the dried fish trade in Nigeria. With
 328 particular reference to lake Chad. *Tropical Stored Product information* **5**: 162.

329 Sokoto State Government (2009). *Sokoto State Diary* Sokoto State Government Printing Press,
 330 Sokoto, Nigeria.

331 SPSS (2007). Statistical, Packages for the Social Science Ver. 16 SPSS Inco.

332