

Original Research Article**THE SPECTRUM OF PSYCHIATRIC MORBIDITY IN SURGICAL
WARDS OF A STATE GOVERNMENT HOSPITAL IN SUB-
SAHARAN AFRICA.****Abstract**

Aim: To determine the incidences and types of psychiatric disorders, mortality, bed stay and management challenges found in our surgical in-patients.

Methods: A three year retrospective study in which all cases with co-morbid psychiatric disorders admitted into the surgical wards of Central Hospital Benin city, Nigeria were studied.

Results: Sixty surgical patients with psychiatric co-morbidity made up of 40 males and 20 females in a ratio of 2 to 1 were studied. 2.3% of all surgical admissions during the study period had psychiatric co-morbidity.

The patients' ages ranged from 18 to 90 years with a mean age of 44.7 years and median age of 45 years.

Post-operative delirium cases formed the bulk of psychiatric co-morbidity (50%) while road traffic accidents involving vagrant psychotics (31.7%) made up the majority of the surgical disorders seen in this study.

The mortality rate in this study was 40% and 87.5% of deaths occurred in cases that developed delirium after surgery.

The bed stay of the patients ranged from 11.7 to 60.9 days with a mean of 33.1 days.

Conclusion: There were management challenges, long bed stay and a high mortality in surgical patients with psychiatric co- morbidity.

A greater care of road traffic accident cases and early identification of mental illness in surgical patients are important. An alternative pain drug control for sickle cell anaemia patient is imperative.

Enhanced knowledge and awareness of psychiatric illnesses in the surgical wards is needful.

KEY WORDS:- Spectrum of Psychiatric disorder, Surgical In-patients

BACKGROUND

30 The prevalence of psychiatric disorders among surgical patients is common and
31 ranges between 23% and 50%¹⁻⁶. It is interesting to note that a remarkable
32 proportion of these cases remain unrecognised by the medical staff.^{7,8}

33 These disorders may occur equally amongst both sexes¹ or more frequently in
34 the female^{2,5,6,9} depending on the location of the study. The older age groups
35 are commonly more affected^{2,5}. It has been recognised that morbidity and
36 mortality are higher in surgical patients with psychiatric co-morbidity.^{6,7,9,10,11,12}

37 Bariatric, oesophageal, plastic, orthopaedic (hip, amputation operations),
38 major abdominal, cardiac and aortic, organ transplant and cancer operations
39 are associated with a high incidence of psychiatric morbidity^{13,14,15,16},

40 Other predisposing events are severe trauma from burns, road traffic and
41 industrial accidents, and assaults^{15,17}.

42 Some of these groups of patients present with disruptive symptoms and
43 behaviours, poor decision-making capacity, substance abuse, problems with
44 coping styles and strategies, preoperative anxiety and health related
45 phobias^{5,15}. Some of the management challenges involved in the care of these
46 patients include problem of obtaining informed consent, treatment adherence,
47 and discontinuation of anti-depressants before anaesthesia⁵.

48 Others challenges encountered are in the administration of psychiatric drugs
49 when the patient is on nil per oral and intra- and post-operative hypotension
50 resistant to intravenous catecholamine administration^{5,18}.

51 The most prevalent symptoms these patients present with are those of
52 depression, anxiety, bipolar and post traumatic stress disorders, alcohol abuse
53 and dependence, and post-operative delirium^{3,5,8,16,19}.

54 Some of these patients are extremely disruptive to care, have bizarre speech,
55 expression and behaviour and thought disorders thereby causing a lot of
56 distress to the doctors, nurses and their relations. Patient contact avoidance
57 may occur as a result¹⁴. Despite all these, the provision of basic psychosocial
58 support for these patients by the medical staff is paramount¹⁶.

59 In the Surgical Unit of Central Hospital Benin City, we noticed an upsurge of
60 patients with psychiatric disorders from vagrant psychotics involved in road

61 traffic accidents, major trauma cases and from complex surgical emergency
62 procedures carried out on very ill patients.

63 The objective of this study therefore, was to determine the types of psychiatric
64 disorders seen in our surgical wards and their impact on management,
65 morbidity and mortality and bed stay.

66 **PATIENTS AND METHOD**

67 All patients admitted into the surgical wards of Central Hospital, Benin City,
68 from December 2013 to December 2016 who also had co-morbid psychiatric
69 disorders were retrospectively reviewed. The diagnosis of psychiatric morbidity
70 was made by the consultant psychiatrist through a consult. Case files of the
71 patients under review and nurses' ward records during the period were
72 studied. Data collated included patients' demographics (age and sex), surgical
73 and psychiatric diagnoses, a nursing and management challenges, and
74 associated morbidities, and mortalities. Data analyses involved the use of
75 simple ratios, ranges, means, medians and percentages.

76 **RESULTS**

77 During the three-year period under study, a total of 2,562 patients were
78 admitted into our surgical units. There were 1,711 males and 851 females in
79 the ratio of 2 to 1. A total of 60 surgical patients with psychiatric co-morbidity
80 were seen, made up of 40 males and 20 females also in the ratio of 2 to 1. In
81 this study, 2.3% of our surgical patients had psychiatric co-morbidity.

82 Table 1 displays the age distribution of 60 patients in the study. Age ranged
83 from 18 to 90 years with a mean age of 44.7 years and median age of 45 years.
84 However, 60% of the cases were over 40years.

85 Table 2 shows the distribution of psychiatric disorders. There was a
86 preponderance of post-operative delirium cases followed by psychosis (mostly
87 in the vagrant cohort).

88 Table 3 shows the spectrum of surgical morbidity seen. Road traffic accidents
89 especially involving vagrant psychotics formed the bulk of the surgical
90 disorders seen in this study.

91 Table 4 displays psychiatric disorder, surgical co-morbidity and plausible
 92 associations. Sepsis and pre-morbid psychosis appear to be most prominent
 93 associations.

94 Table 5 reveals that a majority of the mortality occurred amongst cases that
 95 developed delirium after surgery.

96 Table 6 displays the nature of difficulties encountered in the management of
 97 these cases which included disruptive behaviour and absconding from care
 98 amongst others.

99 Table 7 shows the bed stay of the patients. This ranged from 11.7 to 60.9 days
 100 with a mean of 33.1 days. The highest length of stay occurred in the drug
 101 abuse (60), psychotic vagrant (60.9) and bipolar disorder (60) groups.

102 **TABLE 1: AGE DISTRIBUTION**

Age	No of patients
<20	2
20-30	13
31-40	9
41-50	16
51-60	5
61-70	10
71-80	4
81-90	1
<hr/>	
Total	60

103

104

105

106 **TABLE 2: DISTRIBUTION OF PSYCHIATRIC DISORDERS**

	PSYCHIATRIC DISORDER	NO OF PATIENTS
107		
108	DELIRIUM	30
109	DRUG ABUSE	5
110	ATTEMPTED SUICIDE	7
111	ANXIETY/DEPRESSION	2
112	PSYCHOTIC VAGRANT	10
113	BIPOLAR MOOD DISORDER	2
114	SCHIZOPHRENIA	1
115	TYPHOID PSYCHOSIS	3
116	-----	
117	TOTAL	60

118 **TABLE 3: DISTRIBUTION OF SURGICAL DISEASE**

	SURGICAL DIAGNOSES	NO OF PATIENTS
119		
120	-----	
121	Gastric/ Duodenal/Colonic Carcinoma	4
122	Carcinoma of the Breast	7
123	Surgical morbidity resulting from attempted Suicide	7
124	Burns	1
125	Urological Disorders	5
126	Ruptured Appendicitis	2

127	Chronic Ulcers (sickle cell disease)	5
128	Road Traffic Accident	19
129	Intestinal Obstruction	4
130	Soft Tissue Sarcomas	2
131	Decubitus Ulcer	1
132	Typhoid Perforation	3
133	-----	
134	Total	60
135		

136 **TABLE 4: PSYCHIATRIC/SURGICAL CO-MORBIDITY**

137

PSYCHIATRIC DISORDER	CO-MORBID SURGICAL DISEASE
Delirium	Intestinal Obstruction(sepsis/electrolyte imbalance) Gastric /Duodenal Carcinoma (sepsis/electrolyte imbalance) Burns(sepsis) Fungating Soft tissue Sarcoma (sepsis) Urological Procedures (sepsis) Ruptured Appendicitis (sepsis) Decubitus Ulcer (sepsis) Cancer (?cerebral metastases)
Psychosis	RTA/Vagrant psychotic Typhoid Perforation RTA/Bipolar Disorder

	RTA/Schizophrenia
Drug Abuse	Sickle Cell Haemoglobinopathy/Chronic Ulcers from self injection of pentazocine.
Attempted Suicide	Cut-Throat Laceration of the wrist Oesophagitis/ Oesophageal stricture (Ingestion of corrosives)
Anxiety	Fribrosarcoma (Fear of operation/dying)

138

139 **TABLE 5: MORTALITY**

140	PSYCHIARIC DISORDER	NO OF DEATHS
141	-----	
142		
143	Delirium	21
144	Drug Abuse	--
145	Attempted suicide	1
146	Anxiety	1
147	Psychotic Vagrant	--
148	Bipolar Disorder	--
149	Schizophrenia	--
150	Typhoid Psychosis	1
151	-----	
152	Total	24

153

154 **TABLE 6: CHALLENGES OF MANAGEMENT**

155 SYMPTOMS NO OF PATIENTS

156 -----

157 DISTRUPTIVE BEHAVIOUR 4

158 AGRESSIVE BEHAVIOUR 2

159 ABSCONDERS 5

160 BIZZARE BEHAVIOUR 1

161 -----

162 TOTAL 12

163 **TABLE 7: AVERAGE BED STAY**

164 Psychiatric Co-morbidity Average Bed Stay (Days)

165 -----

166 Delirium 17

167 Drug Abuse 60

168 Attempted suicide 11.7

169 Anxiety 21

170 Psychotic Vagrant 60.9

171 Bipolar Disorder 60

172 Schizophrenia 18

173 Typhoid Psychosis 16.5

174

175

176

177 **DISCUSSION**

178 In this study, the prevalence of psychiatric disorders in our surgical in-patients
179 was low (2.3%), compared to 23 to 50% in other studies¹⁻⁴. Perhaps, many of
180 our cases were either over looked or unrecognised by our surgical staff,
181 ^{5,8,16,20,19} or were from the beginning taken to other centres within the
182 metropolis recognised for the care of the mentally ill patients. If the period of
183 study were longer, perhaps, more of these cases would have been unearthed.
184 Most of the reported incidences in other studies reviewed were from
185 prospective studies done by psychiatrists using varied psychological screening
186 instruments. Moreover, some of these studies included in-patients in the
187 surgical, and medical wards and sometimes gynaecological wards.^{19,20}.

188 We had a preponderance of males (67%) compared to females (33%) in this
189 study in contrast to other studies in which there were either equal numbers
190 of both genders or commonly more females than males^{1,2,8,9,19}. During the
191 period of study, the male admissions (1,711) in our hospital were double that
192 of the female (851). This may have accounted for the higher number of male
193 patients in this study. Again, close to a third (33%) of the cases under review
194 were victims of road traffic and burn accidents and a majority of them were
195 males (60%).

196 Trauma from road traffic accidents (32%) topped the list of surgical conditions
197 with co-morbid psychiatric illnesses in this study. 50% of the trauma cases
198 were vagrant psychotics rushed to our hospital either by good Samaritans, the
199 Police, and the Road Safety Corps or by staff of the Ministry of Women Affairs.
200 Being the only Government Hospital in the centre of town, many of these cases
201 were wont to be brought to our hospital. As vagrant psychotics, they wander
202 from place to place along busy roads, may have poor judgment, are commonly
203 under- fed and under-nourished, weak and perhaps have poor eye sights, and
204 might have become targets of careless drivers on the roads.

205 A prevalence rate of psychiatric illnesses in traumatically injured surgical in-
206 patients as high as 29% was found in another study¹⁷. Brandt and his
207 colleagues equally identified a greater incidence of psychiatric symptoms in
208 injured veterans of the Persian Gulf War compared to those veterans who
209 were not injured.¹⁰

210 In this series, 30% of the cases had abdominal and urologic procedures.
211 Abdominal operations have been recognised to be associated with psychiatric
212 morbidity. George Boras and co-workers analyzed the linked Hospital and
213 Primary Care Data Base in England and found a prevalence rate of 10.1% of
214 post-operative psychiatric morbidity in patients who had abdominal cancer
215 surgery¹⁴. In bariatric surgery, the overall prevalence of current psychiatric
216 disorder was found to be as high as 49%²¹. M Lo on the other hand, found a
217 prevalence rate of 5.1% of mental disorders in adult appendectomy patients
218 in Florida⁵.

219 In our centre, most of the abdominal surgical cases presented as acute
220 emergencies in advanced stages of electrolyte derangement and often times
221 with sepsis. Some of the procedures required long operating times under
222 anaesthesia, because of the complexities of the procedures. These operations
223 sometimes involved bowel or gastric resections and anastomosis or
224 appendectomy and drainage of peritoneal abscesses.

225 Some of these cases developed complications like septic shock, anastomotic
226 leaks, and extravasations of urine into the peritoneum, abdominal sepsis,
227 wound break down, multiple organ failure and delirium. One can confidently
228 argue that the burden of prolonged anaesthesia and complex surgery and
229 adverse post-operation complications, favour the development of acute
230 psychiatric illness.

231 25% of our cases developed psychiatric illness after surgery for malignancies of
232 the stomach, small bowel and colon, breast and soft tissues. Rate as high as
233 59.6% was found among cancer patients in General Hospital facilities in
234 Kenya²⁰. In Uganda, cancer was also found to be the most prevalent physical
235 illness with psychiatric co-morbidity¹⁹.

236 Most of the cancer cases in this study presented at advanced stages. The
237 breast cases already had metastases to the lungs with pleural effusion, to the

238 bones with fractures and to the pericardium with effusion. One of the breast
239 cases had Human Immune Deficiency virus/Acquired Immune Deficiency and
240 bilateral pleural effusion. The sigmoid colon carcinoma cases presented with
241 intestinal obstruction. The prostate carcinoma case had secondaries to the left
242 orbit, left clavicle as well as to the lumbar spine with associated proctosis,
243 fracture of the clavicle and paraplegia respectively. The burden of cancer on
244 the physical and well as the mental well being of these patients was
245 undoubtedly overwhelming and therefore the development of psychiatric
246 illness is perhaps understandable.

247 In our study, 12% (7) of the cases were attempted suicide cases with one
248 fatality (cut-throat). Five (71.4%) cases ingested corrosive (car battery acid),
249 and the remaining two (28.6%) cases had self-inflicted injuries (wrist slashing
250 and cut-throat). There were four males (57%) and three females (43%) in a
251 ratio of 1.3 to 1. The mean age for all the attempted suicide cases was 28.9
252 years while that for the male and female cases were 24.3 and 34.3 years
253 respectively. In a similar study by Jesus Alberdi-Sudupe and others, the
254 incidence of suicide attempts in patients admitted to hospital was 6.9%²². In
255 their study, they found more females (58.4%) than males (41.6 %) cases, a
256 reversal of the gender incidence in our study. In Ibadan, the incidence of
257 deliberate self harm in a six month study in three main hospitals was 0.16%
258 and 76.9% of the cases were under the age of 30 years with a male to female
259 ratio of 1.4 to 1. In the Ibadan study, 61.5% of the cases ingested chemicals
260 while 28.2% took psychotropic drugs²³. In Trinidad, the incidence of deliberate
261 self-harm in a General Hospital was 7.2%. There were more females than male,
262 in the ratio of 2.04 to 1. The mean age of the females was 22.3(SD 5.2) and the
263 mean age of the males was 43.1(SD 3.9) and 47%, 25%, 16% ingested tablets,
264 herbicide and bleach respectively while 8% of cases had self-inflicted injuries.²⁴

265 . Luis Jimenez-Trevino and others in Spain observed that the incidence of
266 attempted suicide in males and females was 2.4% and 1.1% respectively with
267 peak age at 35 to 44 years. 90% of the cases had drug overdose.²⁵ In doing a
268 comparative study; the incidence of attempted suicide in our series which
269 stood at 12% was higher than the findings in Trinidad (7.2%), Spain (6.9%) and
270 Ibadan (0.16). These differences may be attributable to geography, duration of
271 the study and the types of cohorts in the different studies. The down turn in
272 our economy, family pressures, unemployment, peer group influences,
273 progressive loss of extended family support and failure of Government social
274 services might have in no small measures contributed to the high rate of
275 attempted suicide in our study. In our study there were more males than
276 females like in Ibadan but unlike in Spain and Trinidad where there were more
277 females. This difference in the gender incidence could be ascribed to the
278 number of male admissions compared to the female admissions during the
279 period of study. Anecdotally, in our environment, the female appears to be
280 more resilient to psycho- social pressures. The mean age of 28.9 years is similar
281 to the finding in Trinidad (28.2).²⁴ A higher mean age was found in the study
282 done by Jesus et al which stood at 42.5±17.8. This was however a 10 year
283 cross-sectional study.²²

284 The modes of attempted suicide in our study were by ingestion of car battery
285 acid (71.4%) and by self-inflicted injuries (28.6%). Methods in other studies
286 mentioned above were by ingestion of chemicals (61.5%); drugs overdose
287 (28.2%), (47%), and (90%); herbicide (25%), bleach (16%) and deliberate self-
288 inflicted injuries (8%)^{23,24,26} . In our environment, we observed that battery acid
289 was the substance of choice for this act perhaps due to its easy availability and
290 affordability from car battery chargers, petrol stations and retail outlets
291 whereas psychotropic drugs are not easily bought across the counter without a

292 doctor's prescription. The act of slicing ones throat or wrist is not a common
293 finding in our culture. Herbicides are not commonly used in the city and the
294 knowledge of the use of bleach and other substances as vehicles to commit
295 suicide in our environment is perhaps lacking.

296 Pentazocine is commonly used in the management of bone pain crisis of sickle
297 cell anaemia and it is potentially addictive^{27,28}. In this study, five (8.3%) of the
298 patients were sickle cell anaemia patients who became dependent on
299 pentazocine, abused its use chronically and developed ulcers at the injections
300 sites on the upper thighs (80%), and right buttock (20%) regions. There were
301 four males and one female with a mean age of 26.7 years. These ulcers can be
302 described as large, poorly healing, and extending to the muscles and
303 surrounded by woody sclerotic skin and cutaneous tissues²⁸. In the study by
304 Iheanacho and others, 90.9% of sickle cell disease patients with a mean age of
305 34±6 years, who abused pentazocine, were found to have scars and ulcers²⁵.
306 These ulcers are usually indolent, difficult to treat and generally unsightly. The
307 mean age for our patients was lower than theirs probably because most of our
308 cases come from a single centre, the Sickle cell Centre in the City.
309 Axiomatically, the treatment of these ulcers is prolonged, frustrating and
310 requires phased debridement and antibiotic treatment and even with these
311 aggressive measures they remain quite indolent.

312 In our study, the commonest psychiatric co-morbidity was delirium which
313 constituted 50% of all the cases seen compared to the average incidence of
314 delirium of 40% seen in the intensive care unit in a study done by Brian
315 Meguire et al²⁹. In the study by Arott V and others, organic brain syndrome
316 (delirium), depressive disorders and alcoholism were the most prevalent
317 psychiatric co-morbidities in their study⁴. In another study by Clark et al, major

318 depressive illness (34.8%) was the commonest psychiatric disorder in surgical
319 inpatients¹. 14% of psychiatric morbidity in elderly surgical patients was due to
320 post-confusional state, while affective disorder constituted 5% in the study
321 done by Millar³.

322 In different general medical facilities in Kenya, 42% of these cases were found
323 to have mild and severe depression while 41% had bipolar mood disorder,
324 schizophrenia, and psychosis²⁰. In Uganda, depressive disorders which stood at
325 25.2% were the commonest psychiatric co-morbidity found in elderly patients
326 on non-psychiatric wards while depression was the commonest disorder
327 followed by organic disorders (delirium and dementia), adjustment and
328 generalised anxiety disorders in elderly patients admitted to non-psychiatric
329 wards in a general and teaching hospitals in Nigeria^{8,19}.

330 Delirium (organic brain syndrome or post-Confusional state) as the commonest
331 psychiatric co-morbidity in surgical in-patients was also the findings in other
332 studies^{1,3,4}. In this study 41.7% of the cases developed complications like
333 sepsis, severe electrolyte imbalance, bone and perhaps cerebral metastases
334 contributing to the high incidence of delirium in our study. It is important to
335 note that in big hospitals; very ill surgical patients are managed in the intensive
336 care units rather than in the wards as it is done in low resource centres like
337 ours.

338 In our study, we also observed untold nursing challenges from behavioural
339 problems in 12(20%) of the cases. Out this number, four (33.3%) had disruptive
340 behaviour, two (16.7%) aggressive behaviour, one (8.3%) bizarre behaviour
341 while five (41.7%) absconded from the hospital. In the study by Christos
342 Christodoulou and others, a similar incidence of behavioural abnormalities of
343 12% was found in patients admitted into the medical and surgical wards³⁰

344 These behavioural disturbances could be demanding, hostile, manipulating and
345 disturbing the ordered harmony of the ward. About 38% of doctors avoid them
346 while others have negative feelings towards them while others still may just be
347 tolerant, indifferent, and ambivalent or show overt or covert hostility^{14,31}.

348 In our cases, some of the patients forcefully removed inserted naso-gastric
349 tubes, intravenous cannulae and wound dressings. One of the cases was in
350 constant disagreement with co-patients, nurses and doctors and often times
351 engagement in shouting confrontations. A few were double incontinent
352 making nursing care difficult.

353 In our study, the length of bed stay of the patients ranged from 11.7 to 60.9
354 days with a mean of 33.1 days. The highest length of stay occurred in the drug
355 abuse (60), psychotic vagrant (60.9) and bipolar disorder (60) groups. In a
356 study, done in medical and surgical wards, the length of bed stay of surgical
357 patients with psychiatric co-morbidity was 19.8 ± 33.3 compared to 8.3 ± 13.2 of
358 surgical patients without psychiatric illness³². Other studies agree that
359 psychiatric illness delays recovery, increase morbidity and lengthens hospital
360 stay^{19,33}.

361 Many of our patients especially in the vagrant group were abandoned in the
362 hospital as no relations were forthcoming to discharge them. Road traffic
363 accident victims also had a long stay because of the fractures they sustained
364 needed time to heal. The chronicity and the indolent nature of the ulcers of
365 sickle cell anaemia cases who abused pentazocine injection, made
366 management difficult, frustrating, prolonged and time consuming.

367 We found a high in-patient mortality rate of 40% in this study. In the study
368 done by Thod E. Abrams and others an unadjusted 30-day mortality rate was
369 3.8% in patients with psychiatric co-morbidity and 4.0% in patients without
370 psychiatric illness in intensive care patients of All Veteran Health
371 Administration Hospital. In the adjusted 30-day mortality rate however, a
372 moderate increase in mortality was found in the patients who had psychiatric

373 co-morbidity when compared to patients without. These workers found the
374 highest in-patient mortality in depression, post-traumatic stress syndrome,
375 schizophrenia and bi-polar disorders¹⁰.

376 The highest mortality in our study occurred in the post-operation delirium
377 cases. Other studies identified higher mortality rates in patients with chronic
378 psychiatric illness with surgical co-morbidity than those surgical patients
379 without psychiatric disorder^{7,11}. Our patients were very ill patients in advanced
380 stages of sepsis, electrolyte derangement or overwhelming cancer burden, in
381 addition to the stress of anaesthesia and surgery they underwent. From this
382 scenario, the high fatality rate in our study is easy to comprehend when
383 compared to the lower mortality rates in other studies⁷. It has been argued
384 that severe mental illnesses fare poorly after surgery because of late
385 presentation of surgical diseases in these patients, life style factors and
386 multiple chronic medical conditions³⁴. This argument is not tenable in our
387 study because most of our mortalities arose in patients who hitherto were not
388 diagnosed mentally ill patients but developed acute organic factors after
389 surgery. Others claim that psychiatric complications undermine physical and
390 functional recovery and even affect survival, and cause modest increase in the
391 mortality for these patients¹⁹. High mortality has also been associated with
392 patients with carcinoma of the gastrointestinal tract and co-morbid psychiatric
393 illness who had surgery¹⁴.

394 **Conclusion**

395 We acknowledge the limitations of our study; its retrospective design, and the
396 few number of cases studied. Subtle psychiatric morbidity like mild depressive
397 disorder being that they were not disruptive perhaps would have been missed.
398 Hopefully, future research will focus on a prospective study.

399 This study unravelled the problems in the care of vagrant psychotics, who
400 abscond from hospital care and the sickle cell anaemia patient who abuse
401 drugs and develop indolent ulcers that are difficult to treat.

402 Our study showed a high mortality in surgical patients with psychiatric
403 morbidity especially in those with post-operative delirium.

404 Various types of psychiatric morbidity were seen in surgical admissions in
405 Central Hospital, Benin City. There were more males than females in this study
406 unlike in similar studies of this category. The highest number of these cases
407 came from vagrant psychotics who were involved in road traffic accidents.
408 These patients are wont to stay longer in the hospital.

409 A greater care of road traffic accident cases is needed and early identification
410 of mental illness in surgical patients and prompt treatment is essential. An
411 alternative route of pain control with non-addictive drug of sickle cell anaemia
412 patient is imperative. The management of delirium needs a concerted effort
413 and involves aggressive of control of sepsis, electrolyte imbalance,
414 dehydration, hypo-proteinaemia and vitaminosis. The management of these
415 cases in intensive care unit is also imperative.

416 Enhanced knowledge and awareness of psychiatric illnesses in the surgical
417 wards is needful.

418 CONSENT

419 Patients' consent was not sort in this study because they were not directly
420 involved and their identities were not revealed in the study.

421 ETHICAL APPROVAL

422 Necessary ethical approval was obtained by the authors from the Edo State
423 Hospital Management Board ethical committee.

424 REFERENCES

- 425 1. Clark D. M, Minas I.H, Stuart G.W. The Prevalence of Psychiatric
426 Morbidity in General Hospital in Patients. Aust N Z J Psychiatry. 1991
427 Sep;25(3):322-9.
- 428 2. Abiodun O.A , Ogunremi O.O. Psychiatric Morbidity in Medical and
429 Surgical Wards of a Nigerian General Hospital. J. Psychosom Res 1990;
430 34(4):409-14.
- 431 3. Millar H.R. Psychiatric Morbidity in Elderly Surgical Patients. Br J
432 Psychiatry 1981 Jan; 138:17-20.
- 433 4. Arott V, Driessen M, Bangert-Verleger A, Neubauer H,Schürmann A,
434 Siebert W. [Psychiatric Disorders in Hospitalised Internal Medicine and
435 Surgical Patients: Prevalence and Need for Treatment]. Nervenarzt 1995
436 Sep;66(9):670-7.
- 437 5. Richard Uwakwe. Psychiatric Morbidity in Elderly Patients Admitted to
438 Non-psychiatric Wards in a General/Teaching Hospital in Nigeria.
439 International Journal of Geriatric Psychiatry. 15(4): 346-54. May 2000.
- 440 6. Catherine Laurin, Kin L Lavole, Siman L. Bacon, Gilles Dapuis, Guillaume
441 lacoste, André Cartier, Manon Labrecque. Sex difference of Psychiatric
442 Disorders and Psychological Distress in Patients with Chronic Obstructive
443 Pulmonary Disease. Chest 2007; 132(1):148-155.
- 444 7. Levenson JL. Psychiatric Issues in Surgical Patients. Primary psychiatry.
445 2007;14(5):35-39.
- 446 8. Bouras G, Markar SR, Burns EM, Mackenzie HA, Bottle A, Athanasiou T,
447 Hanna GB, Darzi A. Linked Hospital and Primary Care Data Base Analysis of

- 448 the Incidence and Impact of Psychiatric Morbidity following Gastro-
449 Intestinal Cancer Surgery in England. *Ann Surg.* 2016 jul;264(1):93-
- 450 9. Abrams TE, Vaughan-Sarrazin M, Rosenthal GE. Influence of Psychiatric
451 Co-morbidity on Surgical Mortality. *Arch Surg.* 2010;145(10):947-953.
- 452 10. Liberson I, Abelson JL, Amdur RL, King AP, Cardneau JD, Henke P,
453 Graham LM. Increased psychiatric morbidity after abdominal aortic surgery:
454 risk factors for stress-related disorders. *J Vasc Surg.* 2006 May; 43(5):929-
455 34.
- 456 11. Brandt GT, Norwood AE, Ursano RJ, Wright k. Psychiatric morbidity in
457 medical and surgical patients evacuated from the Persian Gulf War.
458 *Psychiatric Services* Feb. 1997;48(1):102-4
- 459 12. Chie Ushijima, Koji Yamazaki, Hidenori Konso, Maskaza Katsurg, Ryo
460 Mori, Sadanori Takeo, Kensuke Ishikawa. Surgical treatment of Lung cancer
461 in patients with Psychological Disorders: A retrospective study. *Anal of*
462 *Cancer Therapy*, 2016; 7: 553-557.
- 463 13. Gagandeep Singh. Pre-surgical evaluation: 6 considerations. *Current*
464 *Psychiatry.* 2010 Oct;9(10):96-96.
- 465 14. Keller M, Sommerfeldt, Fischer C, Knight L, Riesbeck M, Löwe B,
466 Herfarth C, Lehnert T. Recognition of distress and psychiatric morbidity in
467 cancer patients: a multi-method approach. *Ann Oncol.* Aug 2004;
468 15(8):12439.
- 469 15. Hung-Yen L, Chih-Kun H, Chi-Ming T, Hung-Yu L, Yu-His K, Ching-Chung T,
470 Chin-Feng H, Su-Long L, Shu-Ching C, Yung-Chieh Y. Psychiatric disorders of
471 patients seeking obesity treatment. *BMC Psychiatry* 2013; 13:1.

- 472 16. Douglas F. Atzick, S.M Kang, Su Yeong, Kim, David Wishner. Patients
473 with Recognised Psychiatric Disorders in trauma. The Journal of trauma
474 October 2000;49(3): 487-95.
- 475 17.Noeline Nakasujja, Seggane Musisi, James Walugembe. Psychiatric
476 disorders among elderly on non-psychiatric wards in an African setting.
477 International Psychogeriatrics 2007 August 2007; 19(4): 691-704.
- 478 18. Yoshishiko Tsuji, Hiroaki Ohue, Hiroshi Ikata, Osamu Kinoshita, Fumio
479 Shibagaki. Surgical Treatment of Patients with Psychiatric Disorders: A
480 review of 21 Patients. Surgery Today. May 1997; 27(5): 387-391.
- 481 19.DavidbM. Ndetei, Lincoln I. Khasakhala, Victoria N. Mutiso, Francisca A.
482 Ongecha-Owuor, and Donald A. Kokonya. The Prevalence of Mental
483 Disorders in Adults in Different Medical Facilities in Kenya: A Cross-Sectional
484 study. Annals of General Psychiatry 2009; 8:1
- 485 20. Haldis Lier, Eva Biringer, Bjarte Stbhang, Tone Tangen. Psychiatric
486 Disorders and Participation in Pre- and Post-operation Counselling Groups
487 in Bariatric Surgery Patients. Obesity Surgery June 2011; 21(6):730-7.
- 488 21. M. Lo. Prevalence of Diagnosed Mental Disorders Among Florida Adult
489 Appendectomy Patients: Implications for Medical Cost Offset with Mental
490 Health Treatment. The Internet Journal of Mental Health. 2012 volume No
491 1.
- 492 22.Jesús Alberdi-Sudupe, Salvador Pita-Fernandez et al. Suicide Attempts
493 and Related Factors in Patients Admitted to a General Hospital: A Ten-year
494 Cross-sectional Study (1997-2007). BMC Psychiatry 2011; 11:51
- 495 23. Odejide A.O., Williams A.O.,Ohaeri J.U., Ikuesan. The Epidemiology of
496 Deliberate Self- harm: The Ibadan Experience. The British Journal of
497 Psychiatry. Dec.1986; 149(6):734-737
- 498 24. G. Hutchinson, C. Bruce, V. Simmons. Increasing Incidence of Admissions
499 to a General Hospital for Deliberate Self-harm in Trinidad. West Indian Med.
500 J 2008; 57(4): 346-351.

- 501 25.O.E. Iheanacho, I.P. Ezenawenyi, M.E. Enosolease. Pentazocine Abuse in
502 Sickle Cell Disease Seen at a Tertiary Hospital in Nigeria: A Chronic Menace.
- 503 26. Luis Jimenez-Trevino, Pilar A. Saiz, Paul Corcoran et al. The Incidence of
504 Hospital-Treated Attempted Suicide. *Crisis* 2012; 33: 46-53.
- 505 27. Alfred B. Makanjuola, Phillip O. Olatunji. Pentazocine Abuse in Sickle
506 Cell Anaemia Patients: A Report of Two Case Vignattes. *African Journal of
507 Drug and Alcohol Studies*, 2009; 8(2):59-64.
- 508 28. John B. Winifield, Kenneth Greek. Cutaneous Complications of
509 Parenterally Administered Pentazocine. *JAMA* 1973; 226(2):189-190.
- 510 29. Brian Meguire, C.J. Basten, Christopher James Ryan, J. Gallagher.
511 Intensive Care Unit Syndrome: A Dangerous Misnomer. *Archives of Internal
512 Medicine*, May 2000; 160(7): 906-9.
- 513 30. Christos Christodoulou, Katerina Fineti, Athanasios Douzenis, George
514 Moussas, Loannis Michopoulos, Lefteris Lykouras. Transfer to Psychiatry
515 through Consultation-Liason Psychiatry Service: 11 years of experience.
516 *Annals of General Psychiatry* 2008; 7:10 doi10.1186/1744-859X-7-10.
- 517 31. Wyas J.N, Niraj Ahuja. Text-Book Post-Graduate Psychiatry; Jayee
518 Publishers (P) LTD (Pg: 74-89) 2nd Edition: 426-427.
- 519 32. Fulop G., Vita J.J, Hammer J.S. Impact of Psychiatric Co-morbidity on
520 length of Hospital Stay for Medical/Surgical Patients: A Preliminary report.
521 *Am J Psychiatry* 1987 July; 144(7): 878-82.
- 522 33.Sveinsson IS. Postoperative psychosis after heart surgery. *J. Thorac
523 Cardiovasc Surg.* 1975 Oct; 70(4): 717-26.
- 524 34.Laurel A. Copeland, John E. Zeber, Velerie A. Laweren, et al. Serious
525 mental Illnesses Associated with Receipt of Surgery in Retrospective
526 Analysis of Patients in the Veterans Health Administration. *BMC sugery*
527 2015; 15:74.