

Original Research Paper

ASSESSMENT OF THE MICROBIAL POPULATIONS AND PHYSICOCHEMICAL COMPONENTS OF PARAQUAT TREATED SOIL

ABSTRACT

This study assessed the microbial population and physicochemical components of paraquat treated soils. The soil samples were treated with the low, recommended and high doses of paraquat respectively for a period of eight weeks at two weeks interval. Isolation and identification of the soil microbes as well as the determination of the physicochemical components (soil pH, organic carbon and organic matter) were carried out using standard procedures. The results of the study showed that, microbes such as *Staphylococcus* sp., *Micrococcus* sp., *Pseudomonas* sp., *Escherichia coli*, *Bacillus* sp., *Actinomycetes bovis*, *Actinomycetes Israeli*, *Streptomyces* sp., *Aspergillus fumigatus*, *Aspergillus niger*, *Aspergillus flavus*, *Absidia corymbifera* and *Rhizopus stolonifer* were found present in the paraquat treated soils of different doses with the control soil samples having most of the fungal species isolated. The organic matter and carbon contents of the paraquat treated soils of different doses were at the 4th weeks after application (WAA) significantly ($P < 0.05$) the highest whereas the pH of the soils was at 2nd WAA the highest compared to those of the other soils at different WAA. Different doses of paraquat at different WAA affected soil microbial populations as well as physicochemical components of the soil. So, the effects of paraquat on soil microbial population and physicochemical components depended on the concentrations used and the duration of application.

Key words: Paraquat, soil, microbial populations, physicochemical components

1.0 INTRODUCTION

Herbicides are chemical compounds that are used to kill unwanted plants. They are used to clear waste grounds, industrial sites, railways and farm lands (Kellogg *et al.*, 2000). Prior to the wide spread use of chemical herbicides, cultural controls such as altering soil pH and fertility levels were used to control weeds. Mechanical control (including tillage) was also (and still is) used to control weeds (Kellogg *et al.*, 2000). Today, chemical herbicides is the indispensable instruments of reliable and intensive plant production and without their help, hunger and societal tensions would be more serious than the Ecotoxicological effect which they are intended (Bandala *et al.*, 2007). Numerous experiments have been conducted that demonstrated the potential positive impacts of the use of herbicides by farmers; for instance; maize yields doubled and production costs fell by 61% in Nigeria (Benson, 1982). Shave and Avav, (2004) also reported in Nigeria that the cost of weed control in rice with herbicides was 50% lower than hoe weeding.

In achieving the optimization of agricultural resources thus satisfying human needs and at the same time maintaining the quality of the environment and sustaining natural resources, the soil microbial community composition is of great importance, because they play a crucial role in carbon flow, nutrient cycling and litter decomposition, which in turn affect soil fertility and plant growth (Chauhan *et al.*, 2006; Tripathi *et al.*, 2006; Pandey *et al.*, 2007), and hence occupy a unique position in biological cycles in terrestrial habitat. The soil microbial population is considered as active nutrient pool to plants and plays an important role in nutrient cycling and decomposition in ecosystem (De-Lorenzo *et al.*, 2001). A healthy population of soil microorganisms can stabilize the ecological system in soil (Chauhan *et al.*, 2006) due to their ability to regenerate nutrients to support plant growth. Any change in their population and activity may affect nutrient cycling as well as availability of nutrients which indirectly affect productivity and other soil functions (Wang *et al.*, 2008).

However, despite the benefits of using these herbicides by farmers as noted by Bandala *et al.* (2007) and Shave and Avav (2004), the increased use of herbicides in agricultural soils causes the contamination of the soil with toxic chemicals, thus may exert certain effects on non-target organisms including soil microorganisms (Stanley *et al.*, 2013). In Nigeria, especially Mubi local government area of Adamawa State as well as most of the northern part of Nigeria where farming is the major occupation, strict observance of manufacturer's instructions on how to make use of these herbicides are not being adhered to as they are ignorant of its danger to the environment. Natural ways of controlling pests and pathogens of crops which are friendly to the environment have been forfeited by farmers for synthetic herbicides as the synthetic herbicides provide quick solutions to pests or pathogens problems as noted by Bandala *et al.* (2007). Therefore, there is need to occasionally assess the effects of herbicides on micro flora and other soil organisms as reported by Zaid *et al.* (2014) so as to know the measures to be taken. The use of paraquat as herbicide of choice in this study was as a result of it being the most widely used herbicide to control broad-leaf weeds and grasses especially in the study area.

2.0 MATERIALS AND METHODS

2.1 Study area

The study was carried out at Modibbo Adama University of Technology Yola, in the Department of Plant Science Botanical garden, Girei Local Government Area of Adamawa state, Nigeria.

2.2 Experimental design

Completely randomized block design (CRBD) was used to design the experimental field. For each treatment, there was one block which was replicated three times.

2.3 Preparation of herbicides

The paraquat used was purchased from Jimeta Modern Market Yola, Adamawa State of Nigeria from the recommended dealer of herbicides. The preparation/concentration of the paraquat for application was prepared according to the manufacturer's recommendations. One-half and double of the recommended concentration for the herbicide was prepared also with a little modification for application at the rates of 0.3ml, 0.6ml, and 0.9ml (at 150 ml 10L) 0.6 L/ha (at 30 ml in 16 L sprayer). Each of the concentrations had three replications.

2.4 Soil Treatment

The application of the herbicide with the concentrations prepared above was done at two weeks interval over a period of eight weeks. The control blocks were not treated.

2.5 Sample Collection

The top soil samples were collected at the depths of up to 0-1 cm, 1-3 cm, and 3-6 cm respectively from each block of the field during each collection. It was then put into a sterilized polythene bags and were taken to the laboratory for isolation. The soil samples were then made free of large stones and plant debris using 2.0 mm mesh sieve and stored at 4°C. The collection was done at two weeks interval for a period of eight weeks.

2.6 Isolation of Microorganisms

2.6.1 Isolation of Bacteria

Isolation of bacteria was performed by making serial dilution of the taken samples and the dilution that was used for studies were 10^{-2} and 10^{-4} . The dilution was spread in freshly prepared Nutrient agar medium plates. The media was weighed out and prepared according to the manufacture's specification, with respect to the given instructions and directions. The inoculated plates were then incubated at 37°C for 24 – 48 hrs after cooling.

After the incubation period, the pure cultures was obtained based on the color of colony and the pure cultures of the bacterial isolates was subjected to morphological and characterization tests to determine the identity of the bacteria isolates with reference to Bergey's Manual of Determinative Bacteriology as described by Cheesbrough (2006).

2.6.2 Isolation of Fungi

The isolation of fungi was performed by making pour plate method using Potato Dextrose Agar (PDA.) medium. The medium was weighed at 39g/L of distilled water and prepared according to the manufacture's specification, with respect to the given instructions and directions. The plates were then incubated at room temperature for 5 days. After incubation period, the pure cultures were obtained and the fungi were identified (firstly by staining a pinch of the pure culture placed on a slide with lacto-phenol stain and secondly, observing it under the microscope using 10X and 40X objective lens) through its micro and macro morphological structures (Guy, 2006).

2.6.3 Isolation of Actinomycetes

About 2.0ml, 5.0ml, 15ml of glycerol yeast agar and 1.0 g of potassium hydrogen phosphate (K_2HPO_4) were used for the isolation of actinomycetes and after incubation period of 7-14 days. The isolates were characterized based on colony.

2.7 Determination of soil pH

The pH of soil sample was determined using pH meter 3150 Jenway model according to the method described by Onyeike and Osuji (2003).

2.8 Determination of organic matter and organic carbon

117 Organic carbon and organic matter in the soil was determined using Walkly and Black (1934)
118 method.

119 3.0 RESULTS



120 3.1 Fungi, Bacteria and Actinomycetes species identified from paraquat treated and 121 control (untreated) soil samples at two (2) weeks after application (WAA)

122 The results of the fungi, bacteria and actinomycetes identification in paraquat treated and
123 untreated (control) soil samples at two weeks after application (WAA) showed that in
124 paraquat low dose (PLD), paraquat recommended dose (PRD), paraquat high dose (PHD) and
125 the control (CON), fungi, bacteria and actinomycetes such as *A. fumigatus*, *Pseudomonas* sp.,
126 and *A. israelis*; *A. flavus*, *E.coli* and *Streptomyces* sp.; *A. corymbifera*, *Pseudomonas* sp. and
127 *A. bovis*; *A. fumigatus*, *A. flavus*, *A. corymbifera*, *A. niger*, *R. stolonifer* and *Micrococcus* sp.
128 and *Streptomyces* sp. respectively were identified (Table 1).

129 3.2 Fungi, Bacteria and Actinomycetes identified from paraquat treated and untreated 130 (control) soil samples at four (4) weeks after application (WAA)

131 At four (4) weeks after application, the fungi, bacteria and actinomycetes identification in
132 paraquat treated and control soil samples showed that, in PLD, PRD, paraquat high PHD and
133 the CON, fungi, bacteria and actinomycetes such as *A. flavus*, *Bacillus* sp. and *A. bovis*; *A.*
134 *niger*, *E. coli* and *A. Israeli*; *R. stolonifer*, *Pseudomonas* sp. and *Streptomyces* sp.; *R.*
135 *stolonifer*, *A. flavus*, *A. niger*, *A. corymbifera*, *A. fumigatus* and *E. coli* and *A. bovis*
136 respectively were identified (Table 1).

137 3.3 Fungi, Bacteria and Actinomycetes identified from paraquat treated and control soil 138 samples at six (6) weeks after application (WAA)

The result of the identification revealed the presence of fungi, bacteria and actinomycetes such as *A. niger*, *Staphylococcus* sp. and *A. bovis*; *A. flavus*, *Pseudomonas* sp. and *Streptomyces* sp.; *R. stolonifer*, *E. coli* and *A. israeli*; *A. niger*, *A. flavus*, *R. stolonifer*, *A. corymbifera*, *A. fumigatus* and *Bacillus* sp. and *Streptomyces* sp. in PLD, PRD, PHD and CON soil samples respectively (Table 1).

3.3 Fungi, Bacteria and Actinomycetes identified from paraquat treated and control soil samples at eight (8) weeks after application (WAA)

Identified from the PLD, PRD, PHD and CON soil samples were fungi such as *R. stolonifer*, *A. flavus*, *A. niger*, and *R. stolonifer*; *A. flavus*, *A. niger*, *A. corymbifera* and *A. fumigatus* respectively. *Bacillus* sp, *Pseudomonas* sp., *Bacillus* sp. and *E. coli* and *A. bovis*, *Streptomyces* sp., *A. Israeli* and *A. bovis* respectively were the bacterial and actinomycetes species identified from the four soil samples (Table 1).

Table 1: Fungi, Bacteria and Actinomycetes species identified from paraquat treated and untreated soil samples

Week Interval	Treated Soil	Fungal Species	Bacterial Species	Actinomycetes Species
2 nd WAA	Paraquat Low Dose	<i>Aspergillus fumigatus</i>	<i>Pseudomonas</i> sp.	<i>A. Israeli</i>
	Paraquat Recommended Dose	<i>Aspergillus flavus</i>	<i>E. coli</i>	<i>Streptomyces</i> sp.
	Paraquat High Dose	<i>Absidia corymbifera</i>	<i>Pseudomonas</i> sp.	<i>A. bovis</i>
	Control (untreated soil sample)	<i>A. fumigatus</i> <i>A. flavus</i> <i>A. corymbifera</i> <i>Aspergillus niger</i> <i>Rhizopus stolonifer</i>	<i>Micrococcus</i> sp.	<i>Streptomyces</i> sp.
4 th WAA	Paraquat Low Dose	<i>A. flavus</i>	<i>Bacillus</i> sp.	<i>A. bovis</i>
	Paraquat Recommended	<i>A. niger</i>	<i>E. coli</i>	<i>A. Israeli</i>

6th WAA	Dose			
	Paraquat High Dose	<i>R. stolonifer</i>	<i>Pseudomonas</i> sp.	<i>Streptomyces</i> sp.
	Control (untreated soil sample)	<i>R. stolonifer</i> <i>A. flavus</i> <i>A. niger</i> <i>A. corymbifera</i> <i>A. fumigatus</i>	<i>E. coli</i>	<i>A. bovis</i>
	Paraquat Low Dose	<i>A. niger</i>	<i>Staphylococcus</i> sp.	<i>A. bovis</i>
	Paraquat Recommended Dose	<i>A. flavus</i>	<i>Pseudomonas</i> sp.	<i>Streptomyces</i> sp.
	Paraquat High Dose	<i>R. stolonifer</i>	<i>E. coli</i>	<i>A. Israelis</i>
	Control (untreated soil sample)	<i>A. niger</i> <i>A. flavus</i> <i>R. stolonifer</i> <i>A. corymbifera</i> <i>A. fumigatus</i>	<i>Bacillus</i> sp.	<i>Streptomyces</i> sp.
	Paraquat Low Dose	<i>R. stolonifer</i>	<i>Bacillus</i> sp.	<i>A. bovis</i>
	Paraquat Recommended Dose	<i>A. flavus</i>	<i>Pseudomonas</i> sp.	<i>Streptomyces</i> sp.
	Paraquat High Dose	<i>A. niger</i>	<i>Bacillus</i> sp.	<i>A. Israelis</i>
8th WAA	Control (untreated soil sample)	<i>R. stolonifer</i> <i>A. flavus</i> <i>A. niger</i> <i>A. corymbifera</i> <i>A. fumigatus</i>	<i>E. coli</i>	<i>A. bovis</i>

WAA: Week After Application

3.4 Effect of Paraquat Treatment on Fungal Count at 2 – 8 Weeks after Application (WAA)

At the 2nd WAA of PLD, PRD and PHD, the result showed that the fungal count was significantly higher ($P < 0.05$) compared to those of the other WAA with 4.00, 3.00 and 4.00 x 10⁴ cfu/mg respectively. However, that of PRD was not significantly different ($P < 0.05$) from those of 4th – 8th WAA. The CON on the other hand, showed the highest fungal count at 6th

160 and 8th WAA with 5.00 and 5.00 x 10⁴ cfu/mg respectively which were significantly different
161 (P<0.05) from those of 2nd and 4th WAA (Table 2).

162 **3.5 Effect of Paraquat Treatment on Soil Bacterial Population at 2–8 WAA**

163 The bacterial population was significantly the highest (P<0.05) at the 4th – 8th WAA of PLD,
164 PRD and PHD with 33.20, 32.00 and 32.30 x 10⁴ cfu/mg respectively. The control, however,
165 was at 2nd and 6th WAA recorded the highest population of bacterial with 35.00 and 35.00 x
166 10⁴ cfu/mg respectively (Table 2).

167

168 **3.6 The Effect of Paraquat Treatment on Actinomycetes Population at 2 – 8 WAA**

169 The population of actinomycetes at the 2nd WAA of PLD and PRD and even the CON were
170 significantly the highest (P<0.05) with 2.50, 1.40 and 3.48 x 10⁸ cfu/mg respectively. The
171 same thing was observed for PRD at 6th WAA with 6.80 x 10⁸ cfu/mg (Table 2).

172 **3.7 The Effect of Paraquat Treatment on Soil Organic Carbon at 2 – 8 WAA**

173 In Table 3, the application of paraquat doses recorded the most effective action on the soil
174 organic carbon only at the 4th WAA with 1.34 %, 1.12 % and 1.12 % respectively except for
175 its high dose at the 6th WAA which also had the highest effect (1.10) that was not
176 significantly different (P<0.05) with that at the 4th WAA. The control, on the other hand,
177 recorded the most effective action at the 6th and 8th WAA with 1.42 % and 1.43 %
178 respectively (Table 3).

179 **3.8 The Effect of Paraquat Treatments on Soil Organic Matter at 2 – 8 WAA**

180 In Table 3, the application of the different doses of paraquat significantly (P<0.05) showed
181 the most effective action at the 4th WAA with 1.93%, 1.93% and 1.84% respectively. The

182 control, however, was only effective on the soil organic matter only at the 8th WAA with
183 2.46%.

184 **3.9 The Effect of Paraquat Treatment on Soil pH at 2nd - 8th WAA**

185 The application of paraquat doses was at the 2nd WAA recorded the most effective action
186 with 7.39 %, 7.49 % and 7.49 % respectively. Similar action was also observed at 6th WAA
187 for PHD with 7.49 % which was significantly the same ($P < 0.05$) with that at the 2nd WAA.
188 The control, also was at the 2nd WAA had the most effective action with 7.52 % (Table 3).

189

190

Table 2: Effect of paraquat on soil fungi, bacteria and actinomycetes populations

Week	Fungal count (10 ⁴ cfu/mg)				Bacterial Population (10 ⁴ cfu/mg)				Actinomycetes (10 ⁸ cfu/mg)			
	PLD	PRD	PHD	CON	PLD	PRD	PHD	CON	PLD	PRD	PHD	CON
2 nd WAA	4.00 ^a	3.00 ^a	4.00 ^a	2.00 ^b	24.00 ^b	16.00 ^c	9.80 ^c	35.00 ^a	2.50 ^a	2.70 ^d	1.40 ^a	3.48 ^a
4 th WAA	3.00 ^{ac}	2.00 ^a	1.00 ^b	3.00 ^b	33.20 ^a	15.20 ^c	12.30 ^b	16.40 ^b	0.55 ^c	4.00 ^c	0.12 ^c	0.15 ^c
6 th WAA	3.00 ^{ac}	2.00 ^a	2.00 ^b	5.00 ^a	21.40 ^{bc}	32.00 ^a	32.30 ^a	35.00 ^a	1.32 ^b	6.80 ^a	0.17 ^c	2.39 ^b
8 th WAA	2.00 ^{bc}	3.00 ^a	2.00 ^b	5.00 ^a	19.00 ^c	20.30 ^b	9.00 ^c	9.50 ^c	0.82 ^c	5.30 ^b	1.23 ^b	2.76 ^b

191

Key: Means with the same superscript(s) along the column are not significantly different at P<0.05.

192

PLD: Paraquat low dose; **PRD:** Paraquat recommended dose; **PHD:** Paraquat high dose; **CON:** Control; **WAA=** Week After Application

193

194 **Table 3: Effect of paraquat treatment on soil organic carbon, Organic matter and soil pH**

Week	Organic carbon (%)				Organic matter (%)				Soil pH			
	PLD	PRD	PHD	CON	PLD	PRD	PHD	CON	PLD	PRD	PHD	CON
2nd WAA	0.38 ^d	0.55 ^d	0.38 ^c	0.95 ^c	0.65 ^c	0.95 ^d	0.62 ^d	1.63 ^d	7.39 ^a	7.49 ^a	7.49 ^a	7.52 ^a
4th WAA	1.34 ^a	1.12 ^a	1.12 ^a	1.07 ^b	1.93 ^a	1.93 ^a	1.84 ^a	2.33 ^c	6.49 ^b	6.81 ^{ac}	6.84 ^{ac}	6.83 ^{ac}
6th WAA	1.01 ^b	0.99 ^c	1.10 ^a	1.42 ^a	1.62 ^b	1.89 ^b	1.57 ^c	2.44 ^b	5.73 ^{cd}	6.35 ^{bc}	7.49 ^a	6.55 ^{bc}
8th WAA	0.95 ^c	1.06 ^b	0.94 ^b	1.43 ^a	1.63 ^b	1.82 ^c	1.62 ^b	2.46 ^a	6.23 ^{bd}	6.94 ^{ac}	6.36 ^{bc}	6.74 ^{bc}

195 **Key:** Means with the same superscript(s) along the column are not significantly different at P<0.05.

196 **PLD:** Paraquat low dose; **PRD:** Paraquat recommended dose; **PHD:** Paraquat high dose; **CON:** Control; **WAA=** Week After Application

197 **4.0 DISCUSSION**

198 In this study, assessment of fungal species in paraquat untreated (control) soil samples from
 199 the 2 – 8 WAA presents fungal species such as *A. fumigatus*, *A. flavus*, *A. niger*, *A.*
 200 *corymbifera* and *R. Stolonifer* which were found common in all the control soil samples.
 201 Similarly observation was made by Dagze *et al.* (2016) in their control soil samples when
 202 they determined the effect of atrazine on soil microbes. This occurrence could be attributed to
 203 the fungi being well adapted to the soil conditions or probably the health status of the soil as
 204 similarly reported by Zain *et al.* (2013) who associated soil microbial activity and content to
 205 soil health. Also, the similarity in fungal species isolated in the control soil samples of
 206 paraquat (of this study) and atrazine (of Dagze *et al.*, 2016) treated soils could be as a result
 207 of these studies being carried out in the same study area with probably the same
 208 environmental conditions as such factor according to Bolter *et al.* (2006) affect soil microbes.
 209 On the other hand, fungal species such as *A. fumigatus*, *A. flavus*, *A. corymbifera*, *A. niger*
 210 and *R. stolonifer* were found present in the paraquat treated soils with *A. niger*, *A. flavus* and
 211 *R. stolonifer* being the commonest especially from the 4-8th WAA. This finding was
 212 supported by the findings of Ayansina and Oso (2006) who reported the presence of *A. niger*,
 213 *A. flavus* and *Penicillium* sp. in their herbicides treated soil. However, their findings differ in
 214 one of the fungal species identified probably as a result of the differences in the type of
 215 agrochemicals, concentrations, mode of applications, environmental conditions and
 216 herbicides combination used as these according to Subhani *et al.* (2000) and Zain *et al.*
 217 (2013) affect soil's microbes content. Apparent from the assessment of the type of fungal
 218 species present in the untreated (control) soils of this study was the fact that fungal species, *A.*
 219 *corymbifera*, was not present in the paraquat treated soils especially from 4th – 8th WAA. This
 220 could be that the paraquat had a fungicidal effect on the fungi species

221 The results of the identification of bacteria and actinomycetes present in the treated and control
 222 (untreated) soil samples showed that, bacteria such as *Staphylococcus* sp., *Bacillus* sp., *Streptococcus*
 223 sp., *Pseudomonas* sp., *E. coli* and *Micrococcus* sp. were present in both the treated and untreated
 224 (control) soil samples with *Pseudomonas* sp. and *E. coli* being the most common in occurrence in the
 225 soil samples from the 2nd – 8th WAA. On the other hand, actinomycetes such as *A. bovis*, *A. israelis*,
 226 *Streptomycetes* sp. were identified and  the commonest in all the WAA.

227 These bacteria and actinomycetes species identified, diff  cross the number of WAA of the
 228 treatments as well as the doses probably as a result of the effect of the chemicals on them. This was so
 229 because herbicides could adversely affect soil microbes depending upon the application rate/dose and
 230 the type of herbicide used (Ayansina and Oso, 2006; Sebiomo *et al.*, 2011). The bacterial species
 231 identified in the treated and control soil samples of this study were similarly reported by Stanley *et al.*
 232 (2013) in the control and paraquat treated soils. However, some of the bacterial species identified in
 233 this study such as *Staphylococcus* sp., *Streptococcus* sp., and *E. coli* were not reported in their
 234 findings probably as a result of the differences in concentration used, period of application and
 235 collection of samples for assessment and the depth at which the soil samples were collected.

236 The effect of paraquat treatments on soil fungal count at the 2nd WAA of PLD, PRD and PHD showed
 237 a higher fungal count compared to those of other WAA. Similar report was given by Stanley *et al.*
 238 (2013) who reported a decrease in the fungal count of the soil treated with butachlor, pyrazosulfuron
 239 and paraquat with passage of time from 7th to 28th day after treatment. The decrease of the fungal
 240 count observed in this study could be due to the adverse effect of the herbicide on the fungal
 241 component of the soil. This was proven by the fungal count of the control as the fungal count was
 242 higher especially at 6th and 8th WAA.

243 Bacterial population was low for most of the paraquat doses at the 2nd WAA, but increased at other
 244 WAA. Ayansina and Oso (2006) discovered that higher concentrations of herbicide treatments
 245 resulted in much lower microbial counts when compared to soils treated with recommended doses.
 246 This study somehow agrees with the above statement because the bacterial population of the control
 247 soil samples had the highest compared with those of paraquat treated soils.

Actinomycetes population was high at 2nd WAA of PLD, PRD and the control as well as at the 6th WAA of PHD. This could be influenced by the factors such as temperature, pH, organic carbon content, aeration and moisture content of the soils as reported by Arifuzzaman *et al.* (2010). Similar result was published by Stanley *et al.* (2013) who reported a significant increase in actinomycetes population from 7th to 28th day as a result of herbicide application.

The behavior of most herbicides is generally influenced by the content of organic carbon (Torrents and Jayasunda, 1997). The phenomenon of herbicide leaching was primarily related to the adsorption of herbicides in the different soils, mainly due to their contents of fulvic-acids, humus in the organic fraction of the soils, followed by parameters such as the cat-ion exchange capacity, acid dissociation constant (Alister *et al.*, 2011). Variation in the effect of paraquat treated soils on organic carbon content was observed with respect to the doses of the herbicides and the weeks after application. Same was observed for the effect of paraquat on the soil organic matter content as well as pH of the soils. This could be due to vigorous microbial activities in the soil (Greaves *et al.*, 1976).

5.0 CONCLUSION

Different dose of paraquat at different WAA affected soil microbial populations as well the physicochemical components of the soil. So, the effects of paraquat on soil microbial populations and physicochemical components depended on the concentrations used and the duration of application. It also determines the type of microbes present in the soil by getting rid of those that cannot withstand its effects, in this way allowing only the resistible species.

REFERENCES

- Alister. CA and Kogun, G (2011). Effect of physicochemical properties of five agricultural soils on herbicidesoil adsorption and leaching. *Journal of agricultural and crop production Santiago*. **1**(2): 1-8.
- Arifuzzama, M, Kwatun, MR and Rahman, H (2010). Isolation and screening of actinomyletes from sundarbans soil for antibacterial activity. *African Journal of Biotechnology*. **9** (9): 4615-4619.
- Ayansina, ADV; Oso, BA (2006). Effect of two commonly used herbicides on soil micro flora at two different concentrations. *African Journal Biotechnology*. **5**(2): 129-132.

- 276 Bandala ER, Gelover, S, Leal, MT, Bulnes, CA, Jonenez A, and Estrad, CA (2007). Solar
277 photocatytic degradation of Aldrincatal . *Journal of Environmental Research*. **76**: 189 -199.
- 278 Benson, JM (1982). Weed in tropical crops. Review of Abstracts on constraints in Production
279 caused by weeds in maize, rice, sorghum, millet, groundnut and cassava. Food and
280 agriculture organization (FAO) plant production paper; 32 (1).
- 281 Bolter, M, Bloem, J, Meiners, K and Mover, R (2006). Enumeration and bivolume
282 determination of microbial cells: CABI Publishing, Wallingford, UK. Pp.93-107.
- 283 Chauhan, AK, Das, A, Kharkwal, H, Kharkwal, AC and Varma, A (2006). Impact of
284 microorganisms on environment and health, In: A. K. Chauhan and A. Varma (ed.)
285 Microbes: Health and environment.
- 286 Cheesbrough, M (2006). District Laboratory Practice Part II. Cambridge. London, 78-92
- 287 Dagze, JK, Chimbekujwo, IB and Tizhe, TD (2016). Determination of microbial population
288 and physicochemical components of atrazine treated soil. *International Journal of*
289 *Research*, 3(14): 765-783.
- 290 De-Lorenzo, ME, Scott, GI and Ross, PE (2001). Toxicity of pesticides to aquatic
291 microorganisms: a review. *Environmental Toxicology and chemistry*, 20:84-98.
- 292 Greaves, MP, Davies, HA, Marsh, JAP and Wingfield, GL (1976). Herbicides and soil
293 microorganism, *CRC Crit. Rev, microbiology* **5**:1-38.
- 294 Guy, E (2006). Learning to learn the fourth generation: making sense of personalized learning
295 FLO: Bristol 2006 (in Press).
- 296 Kellogg, RI, Nebring, R, Grube, A, Gross, DW and Plotkin, S (2000) Environmental
297 indicators of pesticides leaching and run- of from foram fields. U.S Department of
298 Agriculture and National conservation. Retrived 14th march 2014. From service,
299 <http://www.nrcs.usda.gov/technical/land/pubs/eip.paphtml>.
- 300 Onyeike, EN and Osuji, JO (2003). Research technique in Biological and Chemical Sciences.
301 Springfield Publishers Limited. Pp.369-402.
- 302 Pandey, RR, Sharma, G, Tripathi, SK and Singh, AK (2007). Litter fall, litter decomposition
303 and nutrient dynamics in subtropical natural forest and managed plantations in
304 northeastern India. *Forest Ecology Management*, 240:96-106.
- 305 Sebiomo, A, Ogundero, VW, and Bankole, SA (2011). Utilization and soil minerals .
306 *Research journal of environmental and Earthsciences*. **4** (6). 617-62.
- 307 Shave, PA and Avav, T (2004). Economic of need management in late planted minimum tillage rice
308 in Benue Nigeria. *IRAJ* **29**(1) 73-74.
- 309 Stanley, H O, Maduiké, E M and Okerentugba, PO (2013). Effect of herbicide (atrazine and paraquat)
310 application on soil bacterial population. *Sky Journal of Soil science and Environmental*
311 *Management*, **2**(9):101-105.
- 312 Subhani A, M, Ayam, EC, Huang, JX (2000). Effect of pesticides (Herbicides) on soil microbial
313 biomass review, *Pakistan Journal of Biological Science*. **3**(5)

- 314 Torrents, A, and Jayasunda, S (1997). The sorption of non-tonic pesticides onto clays and the
315 influence of natural organic carbon chemosphere. **35**:1549-1565.
- 316 Walkly, A and Black, IA (1934). An examination of the Degjareff method for determining
317 organic carbon in soil: Effect of variation in digestion conditions and of inorganic soil
318 constituents. *Soil Science*, 63:251-263.
- 319 Wang, QK, Wang, SL and Liu, YX (2008). Responses to N and P fertilization in a young *Eucalyptus*
320 *dunni* plantation, microbial properties, enzyme activities and dissolved organic carbon.
321 *Applied Soil Ecology*, 40:484-490.
- 322 Zaid, AM, Mayouf, M, and Faroiy, YS (2014). The effect of pre-emergent herbicides soil microflora
323 and N. fixing bacteria in Pea field. *International journal of science: basic and Applied*
324 *Research*. **15**(1):131 -138.
- 325 Zain, NMM, Mohammed, RB, Siyam, K, Morshed, MM, and Awang, Y (2013). Effects of selected
326 herbicides on soil microbial populations in oil palm plantation of Malaysia: a microcosm
327 experiment, *African Journal of Microbiology Research*. **5**: 367-374.