

Original Research *Article*

Influence of organic fertilizer (Nomau®) on soil, leaf nutrient content, growth and yield of Physic nut (*Jatropha Curcas*) in Makurdi, North Central, Nigeria.

ABSTRACT

Aim: The aim of investigation was to evaluate the influence of organic fertilizer on soil, leaf nutrient content, growth and yield of Physic nut in Makurdi, North Central, Nigeria.

Study design: The experiment was laid out in a randomized complete block design and replicated **three times trice**.

Place and Duration of the Study: Field experiment was carried out at Teaching and Research Farm of the University of Agriculture, Makurdi, Nigeria during 2009 and 2010 cropping seasons.

Methodology: The trial consisted of four treatments: organic fertilizer (Nomau®) at rate of 0, 0.50, 0.75 and 1.00 t ha⁻¹. Soil samples were collected from the plough layer (0 – 20 cm) at the beginning of the experiment as

Formatted: Highlight

Formatted: Not Highlight

composite. After harvest of the first and second cropping seasons, soil composite samples were collected on the basis of treatments for routine chemical analysis. The Nomau® manure was incorporated into the soil two weeks before planting physic nut. Parameters measured at three weeks intervals number of primary branches, plant height and leaf area and yield at the end of cropping season.

Results: Organic fertilizer is an important contribution of organic matter that improves the physical and chemical characteristics of the soil. The soil at the experimental site had soil pH which was adequate for producing most crops in the tropics, low in organic matter, available phosphorus, cation exchange capacity, exchangeable calcium, potassium and Magnesium. It increased SOM, N, P, Mg, K, Na and CEC. Its application significantly increased number of primary branches, plant height, leaf area, leaf N, P and K concentrations of physic nut and yield relative to control.

Conclusion: The findings from the experiment shows that maximum production of physic nut and soil sustainability can be achieved from treatment 1.00 t ha⁻¹ if properly managed in Markudi, Nigeria.

Key words: Organic fertilizer, physic nut, nutrient content, soil chemical properties, yield.

1. INTRODUCTION

Agriculture is known to be the oldest industry in the world. Its purpose in the growing of Crops and rearing of animals, all geared towards production of food and feed for man and his live stocks. Over the years, grain yields have depreciated drastically due to the degrading nature

60 of soils, poor fertility management and low import
61 technology to improve the fertility of the soil. The use of
62 organic manure could be adopted [1]. Following
63 geometric increase in population, especially in Nigeria,
64 adequate attention should be directed towards massive
65 and cheapest way of food production. In order to
66 achieve this, emphasis should be laid on the easiest
67 means of enriching our soil which organic manure is
68 one. The organic production system aims at supporting
69 and sustaining healthy ecosystems, soil, farmers, food
70 production, the community, and the economy.

71
72 Reduction and elimination of the adverse effects of
73 synthetic fertilizers and pesticides on human health and
74 the environment is a strong indicator that organic
75 agriculture is gaining worldwide attention [2,3]. Organic
76 fertilizers are environmentally friendly, since they are
77 from organic sources [4]. The current global scenario
78 firmly emphasizes the need to adopt eco-friendly
79 agricultural practices for sustainable food production.
80 The cost of inorganic fertilizers is increasing
81 enormously, to the extent that they are out of reach for
82 small and marginal farmers. The organic fertilizers
83 provide nutritional requirements, suppress plant pest
84 populations, and increase the yield and quality of
85 agricultural crops in ways similar to inorganic fertilizers
86 [1,5,6,7]. Organic fertilizers promote microorganisms
87 growth and help to reduce the application of inorganic
88 fertilizers and contribute to improving soil fertility and
89 reducing a negative environmental impact [8]. Organic
90 fertilizers contain macro-nutrients, essential micro-
91 nutrients, vitamins, growth-promoting indole acetic acid
92 (IAA), gibberellic acid (GA) and beneficial

microorganisms [9]. The sustainability of conventional agriculture in world is under threat from the continuous degradation of land and water resources and from declining yields due to the indiscriminate use of agrochemicals. Inorganic fertilizer application has obviously not provided the much needed sustainability of crop production under continuous system [10]. Vanlawe [11] stated that organic matter management remains a corner stone for successful farming in many areas of the tropics.

Physic nut (*Jatropha Curcas*) is a bio-diesel crop belonging to the family of *Euphorbiaceace* which comes from the genus *Jatropha*. It is a large shrub native of Central America and Mexico co-traditionally grown as a hedge in countries of Sub-Saharan Africa, tropical and other sub tropical regions of the world. The name (*Jatropha curcas*) is derived from the Greek words (jatos) meaning “physician” and (trophe) meaning “nutrition” hence the common name physic Nut. It is a small tree or shrub with smooth gray bark, which exudes a whitish coloured watery, later when cut. Normally, it grows between three and five metresmeters in height, but can attain a height of up to eight or ten meters under favourable conditions [12].

Physic Nut is a drought resistant perennial that can grow in wastelands even on gravelly, sandy and saline soils, flowering usually once a year but, under certain conditions, can flower almost throughout the year like in permanently humid regions or under irrigation and fertilization. The ideal climate for *Jatropha* can be summarized as an annual rainfall not exceeding 600mm

Comment [GS1]: Which format of citation??
Use one format in the text.

Comment [SSD2]: This is the recommended format.

Formatted: Font: Italic, Highlight

Formatted: Font: Italic

in moderate climate conditions, 1200mm in hot climatic zones and soil pH less than 9 [13]. The atmospheric temperature is 20⁰c as the plants are sensitive to ground frost that may occur in winter. *Jatropha* grows in soils that are quite infertile, usually found at lower elevations sometimes below 500mm [14] and it can also be grown in deserts due to its resistance to high degree of aridity.

Physic nut leaves are smooth, large and green to pale green in color counting **measuring** 4-6lobes and measuring 10-15cm in length and width. The ratio of male to female flowers averages 29:1 but this is highly variable and may range from 25 to 93 male flowers to 1 to 5 female flowers produced on each inflorescence [15]. It also has been reported that the male-to-female flower ratio declines as the plant ages [16], suggesting that fruiting capacity may increase with age. The unisexual flowers of *Jatropha* depend on pollination by insects, including bees, flies, ants and thrips. One inflorescence will normally produce 10 or more fruits. Fruit set generally results from cross pollination with other individual plants, because the male flowers shed pollen before the female flowers on the same plant are receptive. In the absence of pollen arriving from other trees, *Jatropha* has the ability to self pollinate, a mechanism that facilitates colonization of new habitats [15]. This study was conducted to evaluate the effects of organic fertilizer on soil properties, growth performance, leaf nutrient content and yield of Physic nut in Makurdi, North Central, Nigeria.

2. MATERIALS AND METHODS

2.1 EXPERIMENTAL SITE.

The experiments were conducted at the Teaching and Research farm of the University of Agriculture, Makurdi (Latitude 7° 41¹N and Longitude 8° 35¹E), Benue State during the 2009 and 2010 cropping seasons. The site was not used for any cropping activities for the past three years. The site is located at an elevation of about 98 meters above sea level. This **L**ocation falls within the Southern Guinea Savanna Agroecological zone of Nigeria. The soil of the experimental site was classified as Typic Ustropepts and Typic Haplustults (USDA) or Eutric Cambisols and Haplic Lixisols (FAO),[17].

2.2 EXPERIMENTAL TREATMENTS AND DESIGN

There were four levels of the industrial organic fertilizer (Nomau®). The treatments were replicated three times in a Randomized Complete Block Design (RCBD) giving a total of 12 plots with each plot measuring 4 x 3 m with an alley of 0.5 m between plots and 1m between replications. The Nomau® manure was incorporated into the soil two weeks before planting physic nut seeds. Treatments were: 0.00, 0.50, 0.75 and 1.00 t ha⁻¹.

2.3 LAND PREPARATION AND PLANTING

The land was manually cleared and ridged. The organic fertilizer was incorporated. Physic nut plant was used as a test crop. Planting was done in July, 2009. Physic nut plants were monitored for two cropping seasons. The seeds were manually drilled on rows to 3 cm depth after two weeks of incorporation of organic manure into the soil. The seedlings were later thinned to one per hole at 1 m plant –to –plant spacing three weeks after planting to give a plant population of 10,000 plants per hectare.

Weedings wereas done manually at four weeks interval.

2.4 DATE COLLECTION

Comment [G53]: How many weedings have you done??

Comment [SSD4]: Three weedings were done in each year.

The following growth parameters were measured at three weeks intervals (from three to twelve weeks after planting in the first season and continues from forty nine to fifty eight weeks after planting in the second season) number of primary branches, plant height and leaf area and yield.

2.5 SOIL ANALYSIS

Soil samples were collected from the plough layer (0 – 20 cm) at the beginning of the experiment as composite. After harvest of the first and second cropping seasons, soil composite samples were collected on the basis of treatments. The soil samples were bulked, air-dried and sieved using a 2-mm sieve for routine chemical analysis, as described by Carter [18]. Particle-size analysis was carried out for textural class using the hydrometer method [19]. Soil pH was determined in a soil/water (1: 2) suspension using a digital electronic pH meter. Soil organic carbon was determined by the Walkley and Black procedure by wet oxidation using chromic acid digestion [20]. Total N was determined using micro-Kjeldahl digestion and distillation techniques [21], available P was determined by Bray-1 extraction followed by molybdenum blue colorimetry [22]. Exchangeable K, Ca and Mg were extracted with a 1 M NH_4OAc , pH 7 solution. Thereafter, K was analysedanalyzed with a flame photometer and Ca and Mg were determined with an atomic absorption spectrophotometer [23]. Cation exchange capacity (CEC) was determined using the procedures described by IITA [24] and modified by Anderson and Ingram [25].

2.6 MANURE ANALYSIS

The organic fertilizer (Nomau®) was analyzed for organic matter, nitrogen, phosphorus, Potassium,

Magnesium, Calcium and Sodium. Total N was determined using macro-Kjeldahl method. Available phosphorus was determined by Bray-1 method. The exchangeable K and Na were determined on a flame photometer while Mg and Ca were determined on Atomic Absorption Spectrophotometer (AAS)

2.7 STATISTICAL ANALYSIS

Data collected were subjected to analysis of variance (ANOVA) and significant means were compared using Duncan New Multiple Range Test (DNMRT) at 5% probability level.

3. RESULTS AND DISCUSSION

The mean rainfall distribution for the two cropping seasons was presented in Table 1. In 2009 rainfall was highest in the month of August followed by October and June. In 2010, September recorded highest rainfall of 335.5 mm and was followed by July (196.9 mm), August (178.1 mm) and May (133.1 mm).

Table 1: mean rainfall distribution (mm)

Month	2009	2010
Jan	2.3	0
Feb	0	0
March	3	12.6
April	73.8	31.4
May	196.3	133.1
June	235.6	113
July	84.6	196.9
August	334	178.1
September	141	335.5
October	284.1	121.3

November	1.2	24
December	0	0
Total	1355.9	1145.9

Sources: Federal Ministry of Aviation, Meteorological
Department, Makurdi, Benue State

3.1 SOIL PROPERTIES

The nutrient concentration of organic manure used is presented in Table 2. The organic manure was quite high in organic matter and Ca had more than Mg, N, K, and Na. Table 3 presents initial and after harvest soil analysis data for 2009 and 2010 cropping seasons. The data indicated that the soils were loamy sand with high sand particle. The soils pH were adequate for producing most crops in the tropics, low in organic matter (OM), available phosphorus (P), cation exchange capacity (CEC), exchangeable calcium (Ca), potassium (K) and Magnesium (Mg).

Table 2: Chemical analysis of packaged organic manures used (Nomau)

Nutrient element	Nomau
Organic matter (%)	42.01
Nitrogen (%)	2.91
Phosphorus (mg kg ⁻¹)	12.19
Potassium (cmol kg ⁻¹)	1.62
Magnesium (cmol kg ⁻¹)	3.13
Calcium (cmol kg ⁻¹)	15.91

Sodium (cmol kg⁻¹) 0.69

Table 3: Physicochemical properties of soil before planting and after harvests.

	2009												Formatted: Font: 12 pt	
	Exch.												Formatted Table	
	Cations													
Sample	pH	Org. M	Total N	Avail. P	(cmol/kg)	CEC	Exch. A	Sand	Silt	Formatted: Font: 12 pt				
	H ₂ O	%	mg kg ⁻¹	Na ⁺	K ⁺	Ca ²⁺	Mg ²⁺	cmol/kg	%	Formatted: Font: 12 pt				
Pre	Formatted Table													
plantin	Formatted: Font: 12 pt													
g	6.14	1.53	0.04	7.00	0.15	0.03	2.38	0.56	2.24	1.02	78.00	12.80	9.20	
0.5 t	8.08												Formatted: Font: 12 pt	
ha ⁻¹	6.21	2.17	0.08		0.24	0.05	3.36	0.98	2.81	0.99	72.96	18.40	8.64	
0.75t	9.24												Formatted: Font: 12 pt	
ha ⁻¹	6.25	2.29	0.08		0.29	0.06	3.84	1.14	3.12	0.98	70.96	19.12	9.92	
1.00 t	10.21												Formatted: Font: 12 pt	
ha ⁻¹	6.31	2.62	0.09		0.33	0.07	4.03	1.21	3.09	0.96	70.24	21.84	7.92	
0.00 t	6.83												Formatted: Font: 12 pt	
ha-1	6.02	1.42	0.03		0.16	0.04	2.36	0.49	2.23	1.02	74.24	16.84	8.92	
	2010												Formatted: Font: 12 pt	
0.50 t	8.02												Formatted: Font: 12 pt	
ha-1	6.20	2.11	0.05		0.21	0.04	3.34	0.95	2.74	0.96	72.55	19.01	8.44	
0.75t	9.13												Formatted: Font: 12 pt	
ha-1	6.25	2.22	0.06		0.24	0.04	3.67	1.09	3.05	0.95	71.08	19.12	9.80	
1.00 t	10.07												Formatted: Font: 12 pt	
ha-1	6.32	2.39	0.06		0.29	0.05	3.98	1.15	3.01	0.95	71.03	20.87	8.10	
0.00 t	6.11												Formatted: Font: 12 pt	
ha-1	6.00	1.38	0.02		0.11	0.03	2.31	0.46	2.16	0.99	74.09	16.54	9.37	

Soil chemical analysis results at the end of 2009 and 2010 cropping seasons showed increase among all treatments, for soil pH except the control. The highest pH values 6.31 and 6.32 were obtained at the plot that received 1.00 t ha⁻¹ of nomau while the control has the lowest pH values 6.02 and 6.00. This could be due to the fact that the organic manure had a relatively high buffering capacity based on its high organic matter content (42.01 %) and can fix any change in its pH during organic matter decomposition. In support of this, Ano and Agwu [26] had found that organic manure increased soil pH and macronutrients of soil in southern Nigeria.

The increase in soil organic matter in both cropping seasons by all organic fertilizer treated plots particularly at the higher rates is an indication of the quality of the organic manures and that soil organic matter increases with the amount incorporated. Treatment 1.00.t.ha⁻¹ was the highest rate and recorded the highest organic matter 2.62 and 2.39 % in 2009 and 2010 cropping seasons respectively and the control recorded the lowest values 1.42 and 1.38 %. This increase in soil organic matter with increased rates of organic manure has been reported by Okpara and Mbagwu [27] that increase in organic matter with poultry manure application on vegetable crops; Osundare [28] who reported increase in soil organic matter from the application of organic manure such as sorted city refuse.

Addition of organic fertilizer brought about improvement in most of the soil chemical properties. Total nitrogen, available phosphorus and exchangeable cations were

improved in both cropping seasons. The organic fertilizer decomposed to release organic matter, N, P, K, Ca and Mg. Hence soil OM, N, P, K, Ca and Mg tended to increase with level of organic fertilizer. The plots that received 1.00 t ha⁻¹ of organic fertilizer (nomau) had the highest of these nutrients and the control obtained the lowest in the both cropping seasons. Improvement in nutrient status of organic fertilizer amended plots implies that organic fertilizer could be used for soil management and sustainable production of physic nut. In support of this, Ano and Agwu [26] had found that animal manure increased soil pH and macronutrients of soil in southern Nigeria. Also, Bahl and Torr [29]; Salako [30] reported that poultry manure improved surface P and other major nutrients and yield of maize. These findings confirmed earlier report by Lombin *et al* [31] that animal manure improved soil productivity in two ways; through improvement of the physical conditions of the soil and through the nutrient it supplies to the soil. The nutrient contents of 2010 cropping season was higher in all plots that received nomau than the control. This could be as a result of residual effect of organic manure. This result is in conformity with the studies of IITA [32] in which who it was reported that organic manure incorporated into the soil increase plant nutrients such as nitrogen, phosphorus, potassium, magnesium, calcium and sodium. Nutrients contained in organic manures are released more slowly and are stored for a longer period in the soil ensuring longer residual effects, improved root development and higher crop yields [33].

The results of organic fertilizer on cation exchange capacity for both cropping seasons are presented in

Tables 3. The organic fertilizer treated plots produced highest cation exchange capacity. Among plots treated with organic fertilizer, treatment 1.00 t ha⁻¹ had the highest cation exchange capacity (CEC) .The control had the least value of cation exchange capacity. The increase in cation exchange capacity for all plots that received organic fertilizer in both cropping seasons was the product of high organic matter content that resulted from the applied manures. This increase in cation exchange capacity is responsible for the increase and retention of exchangeable cations as observed in this experiment. This result agreed with Lombin *et al.*, [31] who reported that organic manure is a major contributor to the cation exchange capacity of the soil. Lal and Kang [34] have also reported that the higher the organic matter contents of the soil, the higher the cation exchange capacity. This contribution of organic manure to increase nutrient holding capacity of the soil is consistent with the findings of Davis *et al.*, [35] and Duriugbo *et al.*, [36].

The decrease in exchangeable acidity observed in this study was as a result of these organic fertilizer treatments. The initial exchangeable acidity (1.02 cmol kg⁻¹) was higher than the plots amended with organic fertilizer. The plot with the least exchangeable acidity was 1.00 t ha⁻¹ of nomau which recorded 0.06 and 0.05 cmol kg⁻¹ in 2009 and 2010 cropping season respectively. This could be the effect of calcium released in the soil that resulted in increased soil pH. These results is in conformity with Naramabuye and Haynes [37] who reported that at lower pH values, where aluminium is highly soluble, addition of organic manure

from poultry, pig and cattle caused aluminium concentration to be reduced compared with the control.

This is attributable to the binding of aluminium to solid phase organic matter resulting in its reduction. Ano and Agwu [38] also reported the reduction in soil exchangeable acidity due to application of organic fertilizer.

Comment [GS5]: Very good observations

3.2 GROWTH PARAMETERS

The result of the effect of nomau organic fertilizer on the mean number of primary branches of physic nut is presented in Table 4. Plot treated with 1.00 t ha⁻¹ of organic fertilizer produced the highest mean number of primary branches (3.33) at 3 weeks after planting but were the same statistically with other treatments. At 6, 12, 49, weeks after planting 1.00 t ha⁻¹ of organic fertilizer recorded the highest number of primary branches followed by 0.75 t ha⁻¹ which was statistically similar to other rates except the control that had least performance on number of primary branches. Physic nut height was not significantly affected by organic fertilizer from 3 to 12 weeks after planting (Table 5). At From 49 to 58 weeks after planting, 1.00 t ha⁻¹ of organic fertilizer had the highest mean plant height that was the same statistically with plant height obtained from plots treated with 0.75 t ha⁻¹ and significantly different from the control. The shortest plant height was recorded from plots **treated** with no manure (control). Leaf area recorded under the various rates of nomau showed significant differences ($P = .05$) at 6, 9 and 12 weeks after planting as shown in table 6. Treatment 1.00 t ha⁻¹ amended plot recorded the highest mean value of leaf area, whilst the least was observed under the control.

theise finding are also confirmed by Shah *et al.*[39] who found greater leaf area with the application of organic manure. Michael *et al.*, [40] reported that poultry manure has been found to enhance the number of leaves in lettuce by providing sufficient amount of nutrients that accelerate the growth of leaves.

Table 4: Effect of nomau on primary branches of *Jatropha*

Treatments	weeks after planting							
	3	6	9	12	49	52	55	58
0.50 t ha ⁻¹	2.0 0a	6.00 ab	6.6 7b	7.00 ab	8.33 ab	9.33 ab	10.0 0a	10.3 3b
0.75t ha ⁻¹	3.0 0a	7.00 ab	7.6 7b	8.00 a	9.33 a	10.0 0a	10.6 7a	12.0 0a
1.00 t ha ⁻¹	3.3 3a	8.00 a	8.3 3a	8.67 a	10.6 7a	10.6 7a	11.3 3a	12.3 3a
0.00 t ha ⁻¹	2.0 0a	5.67 b	6.3 3b	6.67 b	8.33 b	8.67 b	9.67 a	10.3 3b

Values with different letters in columns are significantly ($P = .05$) different.

Effect of organic fertilizer treatments on some growth parameters in Tables (4. 5 and 6) revealed significant positive impacts of the different treatments on the studied growth parameters, i.e., primary branches, plant height (cm), and leaf area (cm²) with superiority for the treatment 1.00 t ha⁻¹ which produced the highest values for all the previous characters. These findings are in harmony with those obtained by Kumar *et al.*, [41]. The

superiority of plots treated with organic fertilizer could be because manure contains all the essential micro and macro elements required for plant growth. The significant difference in growth parameter obtained in this study can therefore be attributed to the increased nutrient availability and release of mineral nutrients by organic fertilizer treated plots as observed by Ano and Agwu [38]. The obtained results agreed with the finding of Ajari *et al.*, [42] in okra production in which they reported that organic manure could increase plant height and branches of crops. The results also agreed with the Balyeri *et al.*, [43]; their studies showed that organic manure influenced growth, yield and nutritional quality of containerized aromatic pepper (*Capsicum annuum* L., var 'Nsukka Yellow'). They concluded that the poultry manure increased the leaf area due to sufficient nitrogen availability which in turn improve the vegetative growth of the crop.

Table 5: Effect of nomau on plant height (cm) of Jatropha

Treatments	Weeks after planting							
	3	6	9	12	49	52	55	58
0.50 t ha ⁻¹	29.93a	36.73a	52.10a	56.43a	119.80ab	146.10b	151.73bc	160.50bc
0.75t ha ⁻¹	34.87a	43.43a	55.13a	58.73a	130.63a	156.47a	162.37ab	167.97ab
1.00 t ha ⁻¹	33.40a	41.47a	55.13a	60.73a	130.57a	160.23a	164.33a	171.47a
0.00 t ha ⁻¹	28.20a	36.10a	49.23a	55.27a	114.07b	143.30b	149.40c	157.c

Comment [GS6]: It would be good to put the table in landscape

441 Values with different letters in columns are significantly
442 ($P = .05$) different.

443

444

445 Table 6: Effect of nomau on leaf area (cm²) of Jatropha

446

nts	Weeks after planting							
	3	6	9	12	49	52	55	58
a ⁻¹	120.60a	129.37b	131.20b	133.97c	141.43a	140.80a	137.40a	134.40a
a ⁻¹	137.47a	148.10ba	150.43a	151.50a	141.03a	140.80a	141.53a	140.07a
a ⁻¹	138.00a	146.03a	147.30a	148.67ab	143.10a	141.70a	142.00a	143.53a
a ⁻¹	127.70a	137.63ab	140.70ab	141.17b	141.53a	141.40a	140.10b	139.30a

447 Values with different letters in columns are significantly
448 ($P = .05$) different.

449

450 3.3 NUTRIENT UPTAKE

451 The nitrogen uptake by physic nut as affected by organic
452 fertilizer treatments is presented in Table 7. At 3, 6 and
453 49 weeks after planting nitrogen uptake was highest in
454 plots treated with 1.00 t ha⁻¹ of organic fertilizer (2.01,
455 2.32 and 2.49 % respectively) which was statistically
456 higher than all other treatments except 0.75 t ha⁻¹. At 9
457 weeks after planting, all the plots that received organic
458 fertilizer were statistically similar and higher than the
459 control. At week 55 and 58 after planting the plots that
460 received 1.00 t ha⁻¹ of organic fertilizer had the highest
461 value of nitrogen uptake and were statistically higher
462 than all other treatments. Control had the least value of
463 nitrogen uptake by physic nut and was significantly
464 lower compared to other plots that received organic
465 fertilizer. **application.**

466

Table 7: Effect of nomau on nitrogen uptake (%)

Treatments	Weeks after planting							
	3	6	9	12	49	52	55	58
0.50 t ha ⁻¹	1.84b	2.27b	2.31a	2.34b	2.45b	2.46b	2.46b	2.46b
0.75 t ha ⁻¹	1.98a	2.31a	2.35a	2.36ab	2.48a	2.49ab	2.48b	2.47b
1.00 t ha ⁻¹	2.01a	2.32a	2.35a	2.38a	2.49a	2.51a	2.52a	2.52a
0.00 t ha ⁻¹	1.72b	2.10c	2.11b	2.12c	2.28c	2.31c	2.33c	2.33c

Comment [G57]: Landscape could help to adjust data in tables

Values with different letters in columns are significantly ($P = .05$) different.

Phosphorus uptake at 3 weeks after planting showed no significant difference among the treatments although plot that received 1.00 t ha⁻¹ of organic fertilizer has the highest P value of 0.28 % (Table 8). All the plots treated with organic fertilizer were significantly higher than the control in P uptake. A significant difference was observed on potassium uptake in physic nut due to application of different rates of organic fertilizer (Table 9). The highest content of K was recorded in treatment 1.00 t ha⁻¹ followed by 0.75 t ha⁻¹. All plots treated with manure were higher significantly than the control. The control recorded the least values of K content in physic nut.

Table 8: Effect of nomau on phosphorus uptake (%)

489

Treatments	Weeks after planting							
	3	6	9	12	49	52	55	58
0.50 t ha ⁻¹	0.27 a	0.28 a	0.31 b	0.34 a	0.42 a	0.43 a	0.44 a	0.44 a
0.75t ha ⁻¹	0.26 a	0.28 a	0.32 ab	0.36 a	0.41 a	0.43 a	0.45 a	0.45 a
1.00 t ha ⁻¹	0.28 a	0.31 a	0.35 a	0.37 a	0.43 a	0.44 a	0.45 a	0.46 a
0.00 t ha ⁻¹	0.21 a	0.22 b	0.23 c	0.24 b	0.28 b	0.31 b	0.32 b	0.33 b

490 Values with different letters in columns are significantly
 491 ($P = .05$) different.

492

493

494 Table 9: Effect of nomau on potassium uptake (%)

495

Treatments	Weeks after planting							
	3	6	9	12	49	52	55	58
0.50 t ha ⁻¹	2.60 b	2.8 9c	2.9 5c	3.0 4b	3.3 5b	3.38 b	3.39 b	3.4 1a
0.75t ha ⁻¹	2.90 ab	3.2 1b	3.2 4b	3.2 6a	3.4 1a	3.42 ab	3.42 ab	3.4 2a
1.00 t ha ⁻¹	3.00 a	3.2 5a	3.2 9a	3.1 8a	3.4 1a	3.43 a	3.44 a	3.4 4a
0.00 t ha ⁻¹	2.11 c	2.1 8d	2.2 2d	2.2 3c	2.3 2c	2.35 c	2.36 c	2.3 7b

496 Values with different letters in columns are significantly
 497 ($P = .05$) different.

498

499 The increased availability of nutrient in soil observed for
500 nomau treatments led to increased uptake of N, P and K
501 as indicated by leaf analysis. Ojo *et al.*, [44] reported
502 increases in the levels of P and N in maize plant tissues
503 due to incorporation of organic manure into the soil. This
504 was linked to high content of N and P in nomau organic
505 manure which are released upon its decomposition.
506 Nahar *et al.* [45] and Mandal *et al.*, [46] reported that
507 organic manures incorporated into soil decompose and
508 increase soil nutrient status thereby enhancing nutrient
509 uptake by plants. Earlier, Ojeniyi and Adegboyega [47]
510 found that organic manure increased leaf N, P, K, Ca
511 and Mg contents of Celosia. Also Smith and Ayenigbara
512 [48] found that GD increased leaf N, K and Ca status of
513 Indian Spinach. This observation is in agreement with
514 Isitekhale and Osemwota [49]. Isitekhale and Osemwota
515 [49] advocated the use of manure since it enhances the
516 release, availability and absorption of nutrients when
517 compared to NPK fertilizer even after the first year of
518 application.

519

520 **3.4 PHYSIC NUT YIELD**

521 The seed weight has a direct bearing on the final
522 economic yield of a crop. The differences in the seed
523 weight are generally related to the period between
524 anthesis and maturation of seeds. During this period
525 supply of assimilates to the fruits and formative seeds is
526 crucial and plants with better supply of balanced
527 nutrients are at greater advantage than those under the
528 low nutrient supply [50]. The plot that received 1.00 t ha⁻¹
529 of organic fertilizer (Table 10) produced the highest
530 total yield per hectare (0.74 t ha⁻¹) which was statistically

similar to plots that received organic fertilizer and significantly differed from the control treatment. The control treatment performance was the least among the treatments with the physic nut yield of 0.29 t ha⁻¹ in 2009 cropping season. The yield of physic nut in 2010 showed significant difference among the treatments. The highest yield (1.54) was recorded in treatment 1.00 t ha⁻¹ which was significantly higher than all other rates of organic fertilizer. In general, physic nut crop undergoing various doses of organic fertilizer treatments exhibited increased yield of seeds in comparison to the control; the maximum increase being recorded in plants undergoing organic fertilizer treatment where the seed yield was significantly higher than untreated plants Onwu *et al.*, [51] reported that application of higher doses of nitrogen resulted in higher seed yield of *Jatropha* plant. Patolia *et al.*, [52] recorded significantly higher yields in physic nut plants that received N and P (through urea, single super phosphate, farm yard manure and *jatropha* cake) over the controls.

Table 10: Effect of nomau on plant yield

Treatment	2009	2010
0.50 t ha ⁻¹	0.70a	1.26c
0.75t ha ⁻¹	0.70a	1.38b
1.00 t ha ⁻¹	0.74a	1.54a
0.00 t ha ⁻¹	0.29b	1.15d

Values with different letters in columns are significantly ($P = .05$) different.

4. CONCLUSION

The beneficial effect applications of organic fertilizer on the soil physico-chemical properties favorably influenced pH, CEC, organic carbon, total nitrogen, available phosphorus and potassium. The increases in soil pH, total nitrogen, available phosphorus, exchangeable cations and the reduction in exchangeable acidity are indicative of the facilitative effect of organic fertilizer on soil nutrient release for good crop performance.

Physic nut growth as demonstrated by number of primary branches, plant height (cm) and leaf area (cm²) were significantly influenced by the application of organic fertilizer. Treatment 1.00 t ha⁻¹ has been observed to significantly increase physic nut growth when compared to the others. Although physic nut has the ability to flourish under any condition even without fertilizer application, **but** greater yield is obtained when fertilizer is applied as observed from the research work.

The increased availability of nutrient in soil observed for organic fertilizer treatments led to increased uptake of N, P and K as indicated by leaf analysis.

The observations and findings obtained from the experiment on the Influence of packaged organic fertilizer (Nomau®) on soil, leaf nutrient content, growth and yield of Physic nut (*Jatropha Curcas*) shows that maximum production of physic nut and soil sustainability can be achieved from treatment 1.00 t ha⁻¹ if properly managed in Markudi, Nigeria.

REFERENCES

1. Chiefez, B., Hatcher, P., Hadar, Y. and Chen, Y. Chemical and Biological Characterization of Organic and Municipal solid waste. Environmental Quality. 1996 25:776-785.
2. Abam, P. O., I. N. Onyekwere and K. I Nwosu. Properties and Management of OG Oj soils Cross River State, Nigeria for increased cassava yield, Proceedings of the 40th conference of Agriculture Society of Nigeria. 2006 pp346- 349.
3. Agbogidi, O. M, and Okonmah, C. U. Growth and Yield of Maize as Influenced by Organic Manure type in a Niger Delta Environment International Journals of Agriculture and Rural Development (IJARD) 2012 15 (1) 818 – 824.
4. Anon “Agriculture issue on compost controversy”. Acres U.S.A, June, 2002 p 20.
5. FCAI, “Year Bulletin, Meteorology and Weather”. Federal College of Agriculture, Ishiagu, Ebonyi State Nigeria. 2003
6. FMANR. Literature on soil fertility investigation in Nigeria. A bulletin produced by the Federal Ministry of Agriculture and Natural Resources. Lagos. 1990 p40
7. FAO (Food and Agriculture Organization, Production Year book vol. 50 Rome, Italy. 1995.
8. Mangel, M. Principles of Plant Nutrient”. International Potash Institute, Borne, Switzerland. 1978
9. Olatunji, U. S., Ayuba, A. and Oboh V. U. Growth and yield of Okra and Tomatoes as affected by pig dung and other organic manure. American Jorunal of Plant Physiology 2006.1 (2):78-85

10. Osiname, O.A. Soil fertility in Nigeria: My experience. Agronomy in Nigeria. Agronomy Reunion day, University of Ibadan, Nigeria, ed 4th. October 2000 University of Ibandan Nigera
11. Vanlauwe, B. Soil Organic Matter and Crop Production in West Africa Context. In: Agronomy in Nigeria Agronomy Re-union Day 4th October 2000 University of Ibandan Nigera.
12. Jatropha Plant Source. Source for Elite Planting Material.
http://www.satophaplantsource.com/Jatropha_curcas.html 2006.
13. Foldi N, M. Sanchez, Mittleback M and Hackle S. *Jatropha curcas* Linnaeus as a source for the production of biofuel in Nicaragua, Biological resources Technology. 1996. 58:77-82.
14. World Bank.. World Bank in Jatropha concept. 2002
15. Raju, A.J.S. and V. Ezradanum. *Current Science*, 2002. 83; 11.
16. Achten, W.M.J., L. Verchot, Y.J. Franken, E. Mathijs, V.P. Singh, R. Aerts and B. Muys. *Jatropha* bio-diesel production and use. Biomass Bioenergy. 2008 32: 1063-1084.
17. F.D.A.L.R. Soil Map of Nigeria. Soil Survey Division. Federal Department of land Resources. FMANR – Abuja. 1990
18. Carter M R. Soil sampling and methods of analysis. Canadian Society of Soil Science. Boca Raton (FL): Lewis. 1993. 823 pp.
19. Sheldrick B, Hand Wang C. Particle-size distribution. In: Carter MR, editor. Soil sampling and

- 655 methods of analysis. Ann Arbor (MI): Canadian
656 Society of Soil Science/Lewis. 1993. p. 499–511.
- 657 20. Nelson DW, Sommers LE. Total carbon,
658 organic carbon and organic matter. In: Sparks DL,
659 editor. Methods of soil analysis. Part 3. 2nd ed.
660 SSSA Book Series No. 5. Madison (WI): ASA and
661 SSSA. 1996.p. 961–1010.
- 662 21. Bremner J M. Nitrogen-total. In: Sparks DL,
663 editor. Methods of soil analysis. Part 3. Chemical
664 methods. 2nd ed. SSSA Book Series No. 5.
665 Madison (WI): ASA and SSSA. 1996 p. 1085–1121.
- 666 22. Frank K, Beegle D, Denning J. Phosphorus. In:
667 Brown JR, editor. Recommended chemical soil test
668 procedures for the North Central Region. North
669 Central Regional Research Publication No. 221
670 (revised). Columbia (MO): Missouri Agric. Exp. Stn.
671 1998. p.21–26.
- 672 23. Okalebo JR, Gathua KW, Woomer PL.
673 Laboratory methods of soil and plant analysis. A
674 working manual. 2nd ed. Nairobi (Kenya): TSBF-
675 CIAT, SACRED Africa, KARI, SSEA. 2002. 128 pp.
- 676 24. IITA. International Institute for Tropical
677 Agriculture; Selected methods for soil analysis.
678 Manual series No. 1. Revised edition, IITA Ibadan,
679 Nigeria. 1979 70pp.
- 680 25. Anderson, J. M. and J.S.I Ingram. Tropical
681 Soils Biology and Fertility. A Hand book of Methods,
682 2nd ed. CAB International 1996 pp 57 – 74.
- 683 26. Ano, A.O and Agwu, J.A Effect of animal
684 manures on selected soil properties: II. Nitrogen,
685 potassium and phosphorus. Nigerian Journal of Soil
686 Science (2006)16: 145- 150.

27. Okpara, M.I and J.S.C. Mbagwu. Effectiveness of cattle dung and swine waste as bio fertilizers on an Ultisol of Nsukka. Land Degradation, Agricultural productivity and rural poverty. Environmental implication, proceedings of the 28th Annual conference of the Soil Science Society of Nigeria, Umudike, Nigeria 2003. 71 – 80
28. Osundare, B. Effect of different companion crops and fertilizer types on soil nutrient dynamics and performance of cassava. Nigeria Journal of Soil Science. 2004. 14: 13-17
29. Bahl, G.S and Toor, G.S Influence of poultry manure on phosphorus availability and the standard phosphate requirement of crop estimated from quantity –intensity relationship in different soils. Bioresource Technol 2002. 85:317-322.
30. Salako, F.K. Effect of tillage, mucuna pruiens and poultry manure on maize growth on physically degraded alfisols in Abeokuta, southwestern Nigeria. Nigeria Journal of Soil Science. 2008. 18:10-21.
31. Lombin, I.G; Adepetu, I.A and 16.Ayotade, A. Complementary use of organic manures and inorganic fertilizer in arable production. Proceedings of National Organic Fertilizer Seminar Held at Durbar Hotel, Kaduna, Nigeria. 1991. Pg 146.
32. IITA.. International Institute for Tropical Agriculture; Alley Farming Results Turn to Farmers. IITA Annual Report 1994 pp 107 – 110.
33. Sharma, A.R. and B.N. Mittra. Effect of different rates of application of organic manure. *Journal of Agricultural Science* (Cambridge) 1991. 117:313 – 318

34. Lal, R and B. T. Kang. management of organic matter in soils of the Tropical In: soil management Abstracts. 1982. 302 – 400 (2) 152 – 178.
35. Davis, A S., D. F Jacob, K E Wight. Organic matter amendment of fallow forest tree seedling nursery soils influence soil properties and biomass of sorghums cover crop. Purdue University, West Lafayette, IN 2006 112-118
36. Duruigbo, C L.,J. C Obiefuna and E. U. Onweremadu. Effect of poultry manure rates on the soil acidity in an Ultisol. International Journal of Soil Science 2007. 2 (2) 154-158
37. Naramabuye F. X and R J Haynes. Short-term effects of three animal manures on soil pH and Al solubility. Australian Journal of Soil Research. 2006. 15pp.
38. Ano, A.O. and J.A Agwu. Effect of animal manures on selected soil properties (1) *Nigeria Journal of Soil Science*. 2005. 15: 14-19
39. Shah, S.T., F. Ghafoor, N. Khan, M. Sajid, N. Amin, Z. Shah, S. Bibi, T. Ahmad and F. Haq. Organic fertilizer affect the growth attributes of weeds and swiss chard. Pak. J. Weed Sci. 2016. 22(3): 463-470.
40. Michael T., D. Phiwokwakhe, P.K. Wahome and T.O. Oseni. Study the Effects of organic fertilizers on growth, yield, quality and sensory evaluation of red lettuce.Abjna. .2010 1(6): 1319-1324.
41. Kumar, G.P., S.K. Yadav, P.R. Thawale, S.K. Singh, A.A.Juwarkar: Growth of *Jatropha curcas* on heavy metal contaminated soil amended with industrial wastes and *Azotobacter* – A greenhouse

study. Bioresource Technology. 2008 99 : 2078–
2082

42. Ajari, O., L. E. K. Tsado., J. A. Oladiran., E. A. Salako. Plant height and fruit yield of okra as affected by field application of fertilizer and organic matter in Bida, Nigeria. *The Nigerian Agricultural Journal*. 2003 34: 74-80.
43. Balyeri, P.K., G.T. Otitoju, N. E. Abu and S. Umeh. Poultry manure influenced growth, yield and nutritional quality of containerized aromatic pepper. *Afr. J. Agric. Res.* 2016. 11(23): 2013-2023
44. Ojo, A. O., Adetunji, M. T., Okeleye, K. A., & Adejuyigbe, C. O. Soil fertility, phosphorus fractions, and maize yield as affected by poultry manure and single superphosphate. *International Scholarly Research Notices*, 2015, 8.
<https://doi.org/10.1155/2015/616213>
45. Nahar, Z., Mistry, K. K., Saha, A. K., & Khaliq, Q. A. Response of nitrogen levels on yield of sesame (*Sesamum indicum* L). *South Asian Association for Regional Cooperation Journal of Agriculture*. 2008 6(1), 1-7.
46. Mandal, A., Ashok, K. P., Singh, D., & Rginald, M. Effects of long-term organic and chemical fertilization on N and P in wheat plants and in soil during crop growth. *Agrochimica, Pisa*. 2009. 53, 79-91.
47. Ojeniyi, S.O. and A.A. Adegboyega, Effect of combined use of urea and goat manure in Celosia. *Nigerian Agric. J.*, 2003. 54: 87-90.
48. Smith, M.A.K. and E.A. Ayenigbara, Comparative growth and nutrient composition of Indian spinach in an enricjed humid

tropical environment. Afr. Crop Sci. Conf. Proc., 2001. 5: 1007-1013.

49. Isitekhale H H E, Osemwota I O. Residual effects of poultry manure and NPK fertilizer on nutrient contents and uptake by tomato in the forest and derived savanna soils of Edo State. Nig.J.Soil Sci 2010 20(2):17-25

50. Scott, R. K., Ogunremi, E. A., Ivins, J. D. and Mendham, N. J. The effect of fertilizers and harvest date on growth and yield of oilseed rape sown in autumn and spring. J. Agric. Sci. (Camb.) 1973. 81:287-293

51. Onwu, A.C., Tanimu J. and Waizah Y. Effect of poultry manure on soil properties, nutrient contents and yield of *Jatropha Curcas*. *FUW Journal of Agriculture and Life Sciences*. 2017. 1(2): 97-103

52. Patolia, J.S., Ghosh, A., Chikara, J., Chaudhary, D.R., Parmar, D.R. and Bhuva, H. M. Response of *Jatropha curcas* grown on wasteland to N and P Fertilization Discipline of Phytosalinity, Central Salt and Marine Chemicals Research Institute, G.B. Marg, Bhavnagar 2007.364 002, India.

Formatted: Font: 22 pt