

Growth Performance and Yield Response of Sweetcorn as Influenced by Fertilizer Application Grown Under Bukidnon, Philippines Condition

ABSTRACT

Aims: To determine the influence of fertilizer on the growth and yield of sweetcorn grown under Bukidnon, Philippines condition.

Study design: Field experiment laid out in Randomized Complete Block Design (RCBD)

Place and Duration of Study: Research Station of IPB-UPLB (7° 51' 31.788" N and 125° 3' 40.4568" E), Central Mindanao University, Musuan, Bukidnon, Philippines on February 2016 to May 2016.

Methodology: Soil samples were taken from the site for initial characterization. Six treatments were employed; T₁- no fertilizer, T₂- Recommended rate of inorganic fertilizer (RRIF) based on soil analysis of the experimental area (70 – 50 – 0 N, P₂O₅, K₂O kg ha⁻¹), T₃- 2 tons ha⁻¹ Vermicompost, T₄- ½ RRIF (35 – 25 – 0 N, P₂O₅, K₂O kg ha⁻¹) + 1 ton ha⁻¹ Vermicompost, T₅- ½ RRIF (35 – 25 – 0 N, P₂O₅, K₂O kg ha⁻¹) + 2 tons ha⁻¹ Vermicompost and T₆- RRIF (70 – 50 – 0 N, P₂O₅, K₂O kg ha⁻¹) + 1 ton Vermicompost. Harvesting proceeded at 70 days after sowing (DAS).

Results: The application of Full RRIF + 1 ton Vermicompost ha⁻¹ significantly influenced the plant height of sweetcorn at 20 DAS. Soil's negative logarithm of hydrogen ions present or pH was greatly affected by the application of inorganic fertilizer alone. Moreover, the application of ½ RRIF + 2 tons of Vermicompost ha⁻¹ caused significant effects towards the organic matter content (%) of the soil at harvest. On the other hand, the yield of sweet corn measured by the number of ears expressed in per hectare basis shows to be highly affected by the application of Full RRIF along with 1 ton Vermicompost ha⁻¹.

Conclusion: The combined application of the recommended rate of inorganic fertilizer and Vermicompost are possible ways that may be undertaken in order to yield sweetcorn in higher portion under Bukidnon condition as well as maintaining the quality of the soil of Bukidnon, Philippines.

Keywords: Growth; Yield; Sweetcorn; Fertilizer; Bukidnon.

1. INTRODUCTION

Sweet corn scientifically known as *Zea mays* L. var *Saccharata* is a variety of maize with a high sugar content. Sweet corn is the result of a naturally occurring recessive mutation in the genes which control conversion of sugar to starch inside the endosperm of the corn kernel. Unlike field corn varieties, which are harvested when the kernels are dry and mature (dent stage), sweet corn must be picked when immature (milk stage) and prepared and eaten as a vegetable, rather than a grain. Since the process of maturation involves converting sugar to starch, sweet corn stores poorly and must be eaten fresh, canned, or frozen, before the kernels become tough and starchy [1].

The key to high quality sweet corn is rapid growth, adequate soil moisture and nutrients, and harvesting the ears at optimum maturity. Sweet corn requires rich soil with ample nitrogen and moisture. Soil moisture is found critical for the germination of sweet corn, as it absorb more water than other types for germination to occur [2]. A wide variety of soils is suitable, moreover, it is important that the soil be well drained and well supplied with organic matter. The optimum range of pH for this crop is 5.8 to 7.0.

Fertilizer, natural or artificial substance containing the chemical elements that improve growth and productiveness of plants. Fertilizers enhance the natural fertility of the soil or replace the chemical elements taken from the soil by previous crops [3].

The province of Bukidnon is considered to be the food basket of Mindanao, being the major producer of rice and corn in the region. Products from plantations in the province also include pineapples, bananas and sugarcane. Two types of climate prevail between the northern and southern sections of Bukidnon, The northern part is classified as belonging to Type III, that is, there is no pronounced rain period but relatively dry during the months of November to May. In the southern portion of the province, the climate is classified as Type IV with no dry season [4].

2. MATERIAL AND METHODS

2.1 Location

The field experiment was conducted at the Research Station of IPB-UPLB (7° 51' 31.788" N and 125° 3' 40.4568" E), Central Mindanao University, Musuan, Bukidnon, Philippines.

2.2 Collection, Preparation and Characterization of Soil Samples

Surface soil samples at 0-20 cm depth were collected randomly from the experimental area following a zigzag direction prior to the land preparation. The collected soil samples were placed in cellophane bags and then brought to the Soil and Plant Analysis Laboratory (SPAL), Department of Soil Science, College of Agriculture, Central Mindanao University, Musuan, Bukidnon, Philippines wherein laboratory analyses were conducted. Prior to analysis, the collected soil samples were air-dried at room temperature for about a week, and passed through a 2-mm sieve and were stored in a clean plastic containers. Soil samples were also collected from each experimental plot after harvest of sweet corn. The chemical and physical properties of the soil were determined and analyzed at the Soil and Plant Analysis Laboratory (SPAL). The properties tested include; soil pH in 0.01 M CaCl₂ at a soil to solution ratio of 1:5 [6]; organic matter content by the Walkley and Black method [7]; extractable P using the Bray 2 method [8] and exchangeable K using 1N NH₄OAc buffered at pH 7.0 using a Flame photometer [5].

2.3 Land Preparation and Lay-outing

The total land area used in the experiment was 463.75 m² (35 m x 13.25 m). It was divided into three (3) blocks and each block had a dimension of 131.25 m². A one meter space was provided between blocks and experimental plots as alleyways. The field was plowed using an animal-drawn moldboard plow. Plowing was done twice at one week interval to destroy the emerging weeds. Harrowing was done after plowing to further pulverize larger soil aggregates. Furrows were made at the time of planting at a distance of 75 cm between rows and 25 cm between hills.

2.4 Fertilizer Application

The Vermicompost was sourced out from one of the Vermi farms in Valencia City, Bukidnon, Philippines. The Vermicompost was applied in those plots assigned with organic fertilizer as treatment following the rate of two (2) tons ha⁻¹. It was carefully broadcasted within each plot before the seeding operation. While basal application of inorganic fertilizer was done in treatments assigned to inorganic fertilizer. Inorganic fertilizers were placed in a hole in the furrow covered with a thin layer of soil then followed by the sowing of seeds and then covered again with soil to have a close contact between the seed and the soil, thus, would facilitate uniform germination.

2.5 Care and Management

Care and management immediately started right after seeding up to the harvesting period. Weed population was closely monitored to avoid possible competition of nutrients. Moreover, disease monitoring was also done. Application of pesticides was also employed due to the evident infestation of insect pests. Due to adverse climatic condition during the conduct of the experiment, irrigation was done once a week to sustain the water need of the crop. Irrigation ceased when the experimental plants were about to be harvested at 65 DAS.

2.6 Tagging of Data Plants

Ten (10) sample plants were randomly selected from data rows in each experimental plot. A sheet of white paper was stapled to each data plants to serve as marker and guide during data collection.

2.7 Statistical Analysis

Statistical analysis was done after tabulating the gathered data through the Statistical Tool for Agricultural Research (STAR) software. Moreover, some parameters were found significant as manifested in the F computed value, comparison of means then proceeded using Honestly Significance Difference (HSD) test as the Post hoc test undertaken [9].

3. RESULTS AND DISCUSSION

3.1 Characteristics of Soil in the Experimental Area

Table 1 shows that the soil samples collected from the experimental area has a pH value of 5.52 and is classified as strongly acidic [11]. The soil has organic matter content of 3.90% which is considered marginal [5]. For the extractable phosphorus, it has a value of 17.37 mg kg⁻¹ and is classified as medium in amount [10]. On the other hand, exchangeable potassium was found high in amount because of its value 1.11 cmol kg⁻¹ [10]. Hence, the fertilizer recommendation for the experimental site was 70-50-0 kg ha⁻¹ of N, P₂O₅ and K₂O.

3.2 Growth of Sweet Corn as Affected by Fertilizer Applications

The mean values of plant height at 20, 40, 60 DAS and ear height in plots treated with different fertilizers are presented and discussed in this section.

3.2.1 Plant Height at 20 DAS

Table 1. Chemical properties of soil in the experimental soil (0-20 cm)

Properties	Value	Methods
pH	5.52	0.01 M CaCl ₂
Organic Matter Content, %	3.90	Walkley-Black
Extractable Phosphorus, mg kg ⁻¹	17.37	Bray P ₂
Exchangeable Potassium, cmol kg ⁻¹	1.11	1N NH ₄ OAc / Flame photometer

Table 2 shows the mean plant heights of sweetcorn measured at 20, 40 and 60 DAS. Based on statistical analysis, it was found out that height of sweet corn at 20 DAS was significantly affected by the fertilizers applied. Where plots treated with Full RRIF + 1 ton Vermicompost ha⁻¹ (T₆) got the tallest plants, however, post hoc analysis would say that T₆ has no significant difference with T₂, T₃, T₄ and T₅. But significantly different with that of T₁ (no fertilizer). These results are in agreement with the findings of other researchers [12, 13] who reported that there is a significant increase in growth parameters including plant height

Table 2. Plant height at 20, 40, 60 DAS and ear height of sweetcorn as affected by fertilizer application

TREATMENTS		PLANT HEIGHT, cm			Ear height, cm
CODE	DESCRIPTION	20 DAS [†]	40 DAS	60 DAS	
T ₁	No fertilizer	37.48 b	117.22	195.30	85.23
T ₂	Full RRIF	43.20 ab	135.18	207.87	95.40
T ₃	2 tons Vermicompost ha ⁻¹	43.27 ab	137.65	209.67	95.30
T ₄	½ RRIF + 1 ton Vermicompost ha ⁻¹	40.50 ab	121.85	188.90	89.53
T ₅	½ RRIF + 2 tons Vermicompost ha ⁻¹	41.41 ab	136.00	202.60	94.30
T ₆	Full RRIF + 1 ton Vermicompost ha ⁻¹	44.58 a	135.10	198.70	90.97

† Means followed by the same letter(s) are not significantly different at 5% level of significance based on HSD

and number of leaves of corn plants when applied with NPK fertilizers. At 40 and 60 DAS, no significant difference was noted among plants treated with different fertilizer materials, however, this result is contradictory to the report of [14] that application of fertilizer particularly nitrogen could cause increase in height of corn as it will promote more cell division.

3.2.2 Ear height of plants

Ear heights of sweetcorn plants were measured and is presented in Table 2. Analysis of variance declares no significant influence was observed among the sweetcorn plants by the imposed treatments.

3.3 Yield components and Yield of Sweetcorn as Affected by Fertilizer Application

The mean values of ear diameter, ear length and yield (number of ears per hectare) in plots treated with different fertilizers are presented and discussed in this section.

3.3.1 Ear diameter and ear length of Sweetcorn

The ear diameter of sweetcorn is presented in Table 3. Fertilizer treatments gave no significant effects towards the ear diameter of sweetcorn. Largest ear diameter was observed in those plots applied with 2 tons ha^{-1} of Vermicompost (T_3) with a value of 5.13 cm while smallest was observed in those plots with no fertilizer application (T_1) with a value of 4.73 cm. These results were opposite to the report of [15] who reported that application of amendments like fertilizer with NPK can lead into an increase in plant height, stem girth, number of leaves, leaf area, number of cobs, ear diameter and length, weight of cob, 100-grain weight, and grain yield of maize.

Ear length of sweetcorn gave no significant response on the influence of fertilizers applied. Ear length in plots treated 2 tons ha^{-1} of Vermicompost gave the longest length with 20.47 cm. Shortest length was observed in those plots treated with $\frac{1}{2}$ RRIF + 2 tons Vermicompost ha^{-1} (T_5). Results were conflicting to the results of [12,13,14,15] who reported that application of fertilizers could cause improvements and increase in corn growth and yield performance as it will supply the nutrients needed by the planted crop.

Table 3. Ear diameter, ear length and yield of sweetcorn as affected by fertilizer application

TREATMENTS		Ear diameter, cm	Ear length, cm	Yield [†] (number of ears) ha^{-1}
CODE	DESCRIPTION			
T_1	No fertilizer	4.73	19.93	43757 b
T_2	Full RRIF	5.00	19.43	45197 b
T_3	2 tons Vermicompost ha^{-1}	5.13	20.47	43596 b
T_4	$\frac{1}{2}$ RRIF + 1 ton Vermicompost ha^{-1}	4.93	19.40	44406 b
T_5	$\frac{1}{2}$ RRIF + 2 tons Vermicompost ha^{-1}	4.93	19.20	42222 b
T_6	Full RRIF + 1 ton Vermicompost ha^{-1}	4.97	19.83	48820 a

† Means followed by the same letter(s) are not significantly different at 5% level of significance based on HSD

3.3.2 Yield (number of ears) of Sweetcorn per hectare

Highest number of ears yielded was observed in those plots applied with Full RRIF + 1 ton Vermicompost ha^{-1} (T_6) with a value of 48820 ears per hectare. Moreover, lowest yield was noted in those plants that were applied with $\frac{1}{2}$ RRIF + 2 tons Vermicompost ha^{-1} having a value of 42222 ears per hectare. Statistical analysis declares that the yield of sweetcorn is significantly affected by the treatments. Moreover, post hoc test reveals that T_6 (Full RRIF + 1 ton Vermicompost ha^{-1}) is

significantly the highest among all the treatments imposed. Reports of [16,17] show that there was an increase in yield among sweetcorn plants treated with inorganic fertilizers, organic fertilizers and their combinations. It was known that application of inorganic fertilizer would lead to an increased yield due to the readily available and mineralized nutrients present in inorganic fertilizers along with constant release of nutrients by organic fertilizers. Thus, these reports were confirmations to the results and findings of this study, that fertilizers can cause increase in the yield of sweetcorn.

The experimental area has a marginal amount of organic matter [7] making sweetcorn plants still very productive amidst no application of fertilizer in T_1 . Thus, showing no significant difference as observed in terms of yield and other growth parameters for treatments T_1 , T_2 , T_3 , T_4 and T_5 .

3.4 Soil Chemical Properties at Harvest as Affected by Fertilizer Application

The mean values of soil pH, organic matter content (%), extractable P (mg ka^{-1}) and exchangeable K (cmol kg^{-1}) in plots treated with different fertilizers are presented and discussed in this section.

3.4.1 Soil chemical properties at harvest

The negative logarithm of hydrogen ions present in the soil or commonly known as pH was significantly affected by the imposed fertilizer treatment based on soil analysis conducted [6] after harvest as presented in Table 4. Plots with no fertilizer application (T_1) had the highest pH value of 5.85 which was significantly higher with those plots treated with Full RRIF + 1 ton Vermicompost ha^{-1} (T_6). However, post hoc analysis using HSD at 5% level of significance revealed that T_1 pH value has no significant difference with of T_2 , T_3 , T_4 and T_5 . Results presented by [18] is opposite to the findings of the study.

Table 4. pH, organic matter content, extractable P and exchangeable K of soil at harvest as affected by fertilizer application

TREATMENTS		Some Soil Chemical Properties at Harvest			
CODE	DESCRIPTION	pH [†]	Organic Matter Content, % [†]	Extractable P, mg kg^{-1}	Exchangeable K, cmol kg^{-1}
T_1	No fertilizer	5.85 a	3.93 b	11.00	1.24
T_2	Full RRIF	5.59 ab	3.93 b	14.33	1.20
T_3	2 tons Vermicompost ha^{-1}	5.84 a	4.11 ab	14.17	1.21
T_4	$\frac{1}{2}$ RRIF + 1 ton Vermicompost ha^{-1}	5.72 ab	4.05 ab	10.17	1.13
T_5	$\frac{1}{2}$ RRIF + 2 tons Vermicompost ha^{-1}	5.65 ab	4.15 a	16.33	1.26
T_6	Full RRIF + 1 ton Vermicompost ha^{-1}	5.54 b	4.00 ab	13.33	1.23

[†] Means followed by the same letter(s) are not significantly different at 5% level of significance based on HSD

The reason is due to the short period of time that sweetcorn stays in the field. Sweetcorn plants are harvested in less than 3 months which would cause incomplete reactions in the soil. Leading to a change in pH.

Organic matter content of the soil was found significantly affected by the imposed treatments based on statistical analysis. Highest organic matter content was observed in plots applied with $\frac{1}{2}$ RRIF + 2 tons Vermicompost ha^{-1} (T_5) followed by those plots treated with 2 tons Vermicompost ha^{-1} (T_3), $\frac{1}{2}$ RRIF + 1 ton Vermicompost ha^{-1} (T_4), Full RRIF + 1 ton Vermicompost ha^{-1} (T_6) and lastly T_1 (no fertilizer) and T_2 (Full RRIF). Post hoc test reveals that T_5 value was not significantly different with that of T_3 , T_4 and T_6 . But significantly higher with that of T_1 and T_2 . Application of organic fertilizer like Vermicompost can readily increase and improve the amount of organic matter in the soil as reported by [19].

The extractable P measured in mg kg^{-1} was not significantly affected by fertilizer treatments. However, highest value was obtained by those plots applied with $\frac{1}{2}$ RRIF + 2 tons Vermicompost ha^{-1} (T_5). Exchangeable K was also not significantly affected by the imposed treatments of fertilizer. Highest value was also obtained by those plots treated with $\frac{1}{2}$ RRIF + 2 tons Vermicompost ha^{-1} (T_5). Treatment 5 got the highest values for extractable P and exchangeable K at harvest.

4. CONCLUSION

Results of this study show that the application of inorganic fertilizer and Vermicompost toward Bukidnon soil planted to sweetcorn may cause productivity as revealed in the statistical analysis undertaken. The combined effects of both type of fertilizer is promising towards the productivity and maintenance of soil quality and fertility. Greater yield, being the common objective of any farmer particularly in Bukidnon area, may consider the combined influence of inorganic fertilizer and Vermicompost in sweetcorn production. Thus, the combined application of fertilizers caused promising growth performance towards sweetcorn plants planted under Bukidnon, Philippines condition.

REFERENCES

1. en.wikipedia.org/wiki/Sweet_corn
2. COX, R. 2010. Growing sweet corn in the backyard garden. 888 E. Iliff Avenue, Denver, CO80210. Accessed 21 May 2018. Available: <http://www.colostate.edu/Dept/CoopExt/4dmg/VegFruit/corn.htm>
3. <https://www.britannica.com/topic/fertilizer>
4. en.wikipedia.org/wiki/Bukidnon
5. PCARRD. Standard Method of Analysis for Soils, Plants Tissue, Water and Fertilizers. Los Baños, Laguna, Philippines. 1991.
6. Mclean EO. Soil pH and lime requirement. In: Page AL, Miller RH, Keeney DR. Methods of Soil Analysis, Part 2: Chemical and Microbiological Properties, 2nd Edition. ASA, SSA, Madison, Wisconsin, USA; 1982.
7. Nelson DW, Sommers LE. Organic carbon. In: Page AL, Miller RH, Keeney (eds). Methods of Soil Analysis, Part 2: Chemical and Microbiological Properties, 2nd Edition. ASA, SSA, Madison, Wisconsin, USA; 1982.
8. Bray RH, Kurtz LK. Determination of total, organic and available forms of phosphorus in soil. Soil Science. 1945; 59: 39–45.
9. STAR. Statistical Tool for Agricultural Research version 1.0 <http://bbi.irri.org/>
10. Landon JR. Booker Tropical soil manual: a handbook for soil survey and agricultural land evaluation in the tropics and subtropics. (Booker Tate: Longman Scientific & Technical, Harlow, UK). 1984.
11. USDA. Soil taxonomy: a basic system of soil classification for making and interpreting soil surveys. 1998.
12. Adiaha MS, Agba OA. Influence of different methods of fertilizer application on the growth of maize (*Zea mays* L.). for increase production in South Nigeria. WSN 54 (2016) 73-86 EISSN 2392-2192. Accessed 21 May 2018.
Available: <http://www.worldscientificnews.com/wp-content/uploads/2016/01/WSN-54-2016-73-86-1.pdf>

- 290 13. Makinde EA, Ayoola OT. Growth, yield and NPK uptake by maize with complementary organic
291 and inorganic fertilizers. African Journal of Food, Agriculture, Nutrition and Development. Volume
292 10 No. 3. March 3, 2010. Accessed 21 May 2018.
293 Available: <https://www.ajol.info/index.php/ajfand/article/view/54078>
294
- 295 14. Anjorin FB. Comparative growth and grain yield response of five maize varieties to nitrogen
296 fertilizer Application. Greener Journal of Agricultural Sciences, 3(12): 801-808. Accessed 21 May
297 2018.
298 Available: <http://doi.org/10.15580/GJAS.2013.12.060913818>.
299
- 300 15. Ndukwe OO, IA Ekesiobi, NC Uzundu and ELC Nnabuife. Growth and yield responses of maize
301 (Zea mays) to poultry manure and NPK 15-15-15 fertilizer in Igbariam, Anambra State,
302 Southeastern Nigeria. Inter J Agri Biosci, 3(6): 261-265. 2014. Accessed 21 May 2018.
303 Available: www.ijagbio.com 2014
304
- 305 16. Lazcano C, Revilla P, Malvarb RA. Dom'inguez J. Yield and fruit quality of four sweet corn
306 hybrids (Zea mays) under conventional and integrated fertilization with Vermicompost.
307 (wileyonlinelibrary.com) DOI 10.1002/jsfa.4306. Accessed: 21 May 2018.
308 Available:<https://pdfs.semanticscholar.org/f7a2/9ac300c6f37e8971e0890171143744613f8b.pdf>
309
- 310 17. Mariusz S, Jacek P. Effect of different rates of nitrogen fertilizer on growth and yield of sweet corn
311 cobs. TEKA. COMMISSION OF MOTORIZATION AND ENERGETICS IN AGRICULTURE –
312 2013, Vol. 13, No. 1, 197–200. Accessed 21 May 2018.
313 Available:agro.icm.edu.pl/agro/element/bwmeta1.element.agro...4dfd.../197_Teka_13_1.pdf
314
- 315 18. Arsova A. Effect of fertilizer application and soil ph on the acidic and sorption properties of maize
316 leaves and stems. BULG. J. PLANT PHYSIOL., 1995, 21(1), 52–57. Accessed: 21 May 2018.
317 Available: http://www.bio21.bas.bg/ipp/gapbfiles/v-21/95_1-52_57.pdf
318
- 319 19. Canatoy R. Growth and yield response of sweet corn (Zea mays l. var. saccharata) as affected by
320 tillage operations and fertilizer applications. International Journal of Education and Research Vol.
321 6 No. 4 April 2018. Accessed: 21 May 2018.
322 Available: <http://www.ijern.com/journal/2018/April-2018/22.pdf>
323