

## Original Research Article

### Geostatistical Evaluation of Spatial Variability of Selected Soil Physical Properties under Different Landuse Systems in Ado Ekiti, southwest Nigeria

#### ABSTRACT

The characterization of spatial variability of soil physical and chemical characteristics is very important for precision farming and managing agricultural soils. Therefore, the objectives of this study therefore were to evaluate some selected soil physical properties of a cultivated field in Ado Ekiti, southwest Nigeria and quantify the spatial characteristics of the evaluated properties using classical statistical and geostatistical techniques. The field was planted to cowpea, sole maize and maize/cassava intercrop. A total of one hundred and eighty-four (184) georeferenced surface samples were collected for analysis of texture, bulk density (BD), particle density (Pd), porosity (Pt) and saturated hydraulic conductivity (Ksat). The soil properties showed varying degree of spatial variability, with Ks highly variable. There was weak correlation between Ksat versus BD and Pt but the correlation was significant with sand content. The variability of these properties revealed weak to strong spatial dependence. The BD, Pd, Pt and Ksat could be well described using either Gaussian or spherical models while sand and clay content gave pure nugget effect. The range of spatial dependence values indicated that future sampling could be done within a distance between 214 and 511 m. The kriged maps further showed the spatial distributions of these soil physical properties across the three different land use systems. The documentation of these physical properties in field scale distribution maps will allow derivation of zones of physical and mechanical sensitivity. This can further help define management zones, which can be combined with minimum soil samples to provide a more accurate prediction of spatial variability of soil properties for site-specific soil management under different agricultural land use systems.

29 Keywords: Spatial variability, classical statistics, geostatistics, soil management, soil physical  
30 properties

## 31 **1.0 INTRODUCTION**

32 Soils are known to vary across landscapes and so do their properties. Significant  
33 within – field variability attributable to natural factors of soil formation and crop management  
34 practices has also been reported [1]. Under similar management practices, soils in agricultural  
35 fields have shown highly variable properties [2]. In view of this within –field variability in  
36 soil properties, applying uniform management treatments, such as blanket fertilizer  
37 application or excessive tillage, often result in over – application of such inputs in low-  
38 yielding areas and over application of inputs in high-yielding areas [3].

39 Quantifying the spatial variability of soil properties therefore becomes appropriate in  
40 farm planning and management for developing a more productive and efficient crop  
41 management systems [1]. Traditionally, the spatial variability of soil properties has been  
42 evaluated through classical descriptive statistics and geostatistical techniques that verify  
43 relationships among several soil samples of a specific area or field, using the study of  
44 regionalized variables [4]. While classical statistics uses the measure of central tendency to  
45 quantify only the degree of spatial variability of soil properties within the field, geostatistical  
46 analysis methods of variography and kriging have been proven to be more useful for  
47 characterizing and mapping spatial variation of soil properties and have also received  
48 increasing interest by soil scientists and agricultural engineers [5, 6, 7, 8]. In quantitative  
49 evaluation of within – field spatial variability, geostatistical technique has been successfully  
50 applied by various authors [e.g. 9, 10, 1, 11]. Nigeria agricultural soils are also characterized  
51 by the variability of soil properties in space and thus the variability of crop yield within field,  
52 however field management has remain uniform such as blanket application of fertilizer. This

53 practice portends danger to the environment as well as increased cost of production.  
54 Elsewhere, the study of spatial variability of soil properties has been used to generate  
55 information to mitigate these problems through precision farming. Therefore, the objectives  
56 of this study therefore were to evaluate some selected soil physical properties of a cultivated  
57 field and quantify the spatial characteristics of the evaluated properties using classical  
58 statistical and geostatistical techniques.

59

## 60 **2.0 MATERIALS AND METHODS**

### 61 **2.1 Description of study site**

62 The study site is a 3-ha field cultivated to arable crops (cowpea, sole maize and  
63 maize/cassava intercrop) located on the SIWES Training Farm of the Teaching and Research  
64 Farm, Ekiti State University, Ado Ekiti, Ekiti State. The site is located on latitude 70 41' 57.9'  
65 N, longitude 50 05' 0" E and 406 m above the mean sea level. The land has been previously  
66 used for the cultivation of yam and cowpea and was left fallow for about 3 years before the  
67 SIWES students started cultivating on it for training on crop production.

68

### 69 **2.2 Field procedure and soil sampling**

70 Of the 3-ha field, 1 ha planted to cowpea, 1 ha to sole-maize and only about 0.7 ha to  
71 maize/cassava inter-crop were used for the study. Grids were set up on the field within the  
72 three land use. Ninety-four (94) grids (10 m x 10m) were set up in cowpea plot, fifty (50)  
73 grids (20 m x 10 m) in sole maize and forty-four (44) grids (15 m x 10 m) in maize/cassava  
74 intercrop, giving a total of one hundred and eighty-four (184) grids (Figure 1). The center of  
75 each grid was geo-referenced with the aid of GPS (Garmin model) for soil sampling.  
76 Disturbed and undisturbed soil samples were collected from the 0-20 cm surface layer at the  
77 center of each grid. Thus, a total of one hundred and eighty-four (184) samples were

78 collected altogether. The samples collected were neatly packed and transferred to the  
79 laboratory for analysis.

### 80 **2.3 Evaluations**

81 *Soil texture.* The granulometric analysis was determined using the modified hydrometer  
82 method following the procedure described in [12] from disturbed air-dried soil samples after  
83 passing through 2-mm sieve.

84 *Bulk density.* After preparation in the laboratory, the undisturbed core samples were oven-  
85 dried at 105°C for 48 h and the weight of dry soil was determined. The bulk density was  
86 determined using the equation according to [13]:

$$87 \qquad \qquad \qquad BD = \frac{M_s}{V_s}$$


88 1

89

90 where  $BD$  is bulk density,  $\text{g cm}^{-3}$ ;  $M_s$  is weight of dry soil, g;  $V_s$  is volume of soil,  $\text{cm}^3$ .

91 *Particle density.* Particle density was determined using volumetric bottle method following  
92 the procedure described in [14] from disturbed air-dried soil samples after passing through 2-  
93 mm sieve and then oven-dried for 24 h.

94 *Total porosity.* It was determined using the relation:



95

96 Figure 1. (a) Map of Nigeria showing (b) Ekiti State and (c) the study site

97

98 
$$Pt = 1 - \frac{BD}{Pd}$$

99 2

100

101 where  $Pt$  is the total porosity,  $\text{cm}^3 \text{cm}^{-3}$ ;  $BD$  is the bulk density,  $\text{g cm}^{-3}$ ;  $Pd$  is the particle  
 102 density,  $\text{g cm}^{-3}$ .

103 *Soil saturated hydraulic conductivity.* Soil saturated hydraulic conductivity ( $K_{sat}$ ) was  
 104 determined by the constant-head permeameter [15] on undisturbed soil samples collected in  
 105 metal cylinders (of known volume) after saturation by capillarity in a water bath for 48 hours.  
 106 The determination of  $K_{sat}$  was performed by collecting and measuring the amount of water

107 that percolates through the soil sample under a constant hydraulic head of about 3 cm in the  
 108 water column, according to the methodology described by [12]. From the data, soil Ksat was  
 109 calculated according to Equation 3.

110

$$111 \quad K_{sat} = \frac{Q * L}{A * H * t}$$

112 3

113

114 where Ksat is saturated hydraulic conductivity, cm/hr; Q is volume of water that flow through  
 115 the soil column in a given time, cm<sup>3</sup>; L is length of the soil column, cm; H is length of soil  
 column + water head above the soil column, cm; A is area the soil column, cm<sup>2</sup>; t is time, h.

116

## 117 **2.4 Data analysis**

### 118 *2.4.1 Descriptive statistics of soil properties*

119 Descriptive statistics of minimum, maximum, average, standard deviation (SD),  
 120 skewness, kurtosis and coefficient of variation (CV) of data on sand, clay, bulk density,  
 121 saturated hydraulic conductivity, particle density and total porosity. The saturated hydraulic  
 122 conductivity data that did not follow normal distribution (Shapiro-Wilk test) was log  
 123 transformed for further analysis. In addition, the frequency distribution graph was plotted for  
 124 each variable. All classical statistical analyses were carried out using SPSS (IBM version 20).

125

### 126 *2.4.2 Geostatistical analysis*

127 Geostatistical analysis was done using the GS+ (Gamma Design Software, Version  
 128 5.2, 2005) to determine the spatial dependency and estimation of the soil properties  
 129 evaluated. Isotropic semivariograms of linear, power, spherical, exponential and Gaussian,  
 130 were tested from omnidirectional semivariances,  $\hat{\gamma}(h)$ , of a set of spatial observations,  $Y_{xi}$ ,  
 131 expressed as [16]:

132

$$133 \quad \hat{\gamma}(h) = \frac{1}{2N(h)} \sum_{i=1}^{N(h)} (Y_{x+h} - Y_x)^2$$

134

4

135

136 where  $\hat{\gamma}(h)$  is the covariance;  $h$  is the spatial separation distance, known as the time lag;  
 137  $N(h)$  is the number of pairs of observations separated by a distance;  $Y_x$  is soil variable  
 138 observed at point  $x$  while  $Y_{x+h}$  soil variable observed at point  $x + h$ .

139

To characterize the spatial covariance structure of the variables, the best model was  
 140 selected based on the coefficient of determination,  $R^2$ . From the models, basic spatial  
 141 parameters such as nugget (Co), sill (C+Co) and range (Ao) were determined. The nugget to-  
 142 sill ratio expressed as the structural variance was calculated for each soil physical property  
 143 and used to evaluate the degree of spatial dependence associated with each soil property.  
 144 Structural variance values were categorized into one of three classes of spatial dependence as  
 145 proposed by [17]. For structural variance less than 0.25, the variable is considered strongly  
 146 spatially dependent; if the structural variance is greater than 0.25 and less than 0.75, the  
 147 variable is considered moderately spatially dependent; and if the structural variance is greater  
 148 than 0.75, the variable was considered weakly spatially dependent [17, 18]. In addition, a  
 149 structural variance value close to zero indicates continuity in the spatial dependence.

150

After selecting the best fit semivariogram model for each variable, contour maps were  
 151 created through ordinary kriging of the Geostatistical Analyst extension in ArcGIS v. 10.1<sup>®</sup>  
 152 (Esri, Redland, CA, USA). Cross-validation of the kriged results was made using validation  
 153 statistics of mean absolute error (MAE) and mean square error (MSE) as:

154

$$MAE = \frac{\sum_{i=1}^N |z^* - \bar{z}|}{N}$$

155 5

156 
$$MSE = \frac{\sum_{i=1}^N (z^* - \bar{z})^2}{N}$$

157 6

158 where  $z^*$  is the predicted soil variable;  $\bar{z}$  is the mean of measured soil variable;  $N$  is the total  
159 number of sampling locations. The predicted values for each soil variable were obtained from  
160 the cross-validation procedure in the GS<sup>+</sup>.

161

## 162 **3.0 RESULTS AND DISCUSSION**

### 163 ***3.1 Descriptive statistics***

164 The descriptive statistics of soil variables of the SIWES Training Farm is presented in Table  
165 1. The sand content ranged between about 51 and 68% (mean = 64.3%) while clay content  
166 was low, ranging between 2 and 11% (mean = 7.04%). The soil had BD ranging from 1.10 to  
167 1.73 g cm<sup>-3</sup> (mean = 1.43 g cm<sup>-3</sup>) while the particle density ranged from 2.02 to 2.97 g cm<sup>-3</sup>  
168 (mean = 2.51 g cm<sup>-3</sup>). For total porosity (Pt), the values were between 0.27 and 0.056 cm<sup>3</sup>  
169 cm<sup>-3</sup> (mean = 0.43 cm<sup>3</sup> cm<sup>-3</sup>). The saturated hydraulic conductivity (Ksat) ranged from 2.35  
170 to 326.20 cm h<sup>-1</sup>, with an average value of 48.74 cm h<sup>-1</sup>. For Ksat, the results are in agreement  
171 with the findings of [19]and [20] who from different studies reported high variability in Ksat.  
172 The relatively low values of BD and clay content obtained from the study could have led to  
173 increase in the value of Ksat. Low Ksat also indicated low level of compaction and presence  
174 of large number of macropores which allow water to percolate through the soil. The least  
175 varied physical property was found to be particle density. For instance, the spatial distribution  
176 of water retention properties closely followed the distribution pattern of sand and clay  
177 content. This indicates a differential water retention capacity of different textured soils across  
178 the field. The relatively high variability of Ksat may be attributed to differences in soil pore  
179 geometry as a result of soil disturbance. Increase in porosity could be as a result of low bulk

180 density i.e. degree of compaction and granulation is very low and also increase in organic  
181 matter.

182 Table 1. Descriptive statistics of soil physical properties of the field.


Property	N	Min.	Max.	Mean	SD	CV	Skewness	Kurtosis
Sand, %	184	51.29	67.65	64.30±0.170	2.35	0.037	-1.85±0.18	6.04±0.36
Clay, %	184	2.32	11.32	7.04±0.110	1.49	0.211	-0.13±0.18	0.27±0.36
BD, g cm <sup>-3</sup>	184	1.10	1.73	1.43±0.098	0.13	0.093	-0.07±0.18	-0.56±0.36
Pd, g cm <sup>-3</sup>	184	2.02	2.97	2.51±0.011	0.13	0.050	-2.24±0.18	14.04±0.36
Pt, cm <sup>3</sup> cm <sup>-3</sup>	184	0.27	0.56	0.43±0.004	0.06	0.137	-0.31±0.18	-0.33±0.36
Ksat, cm h <sup>-1</sup>	94	2.35	326.20	48.74±5.928	57.50	1.179	2.61±0.25	8.14±0.49

183 BD: bulk density; Pd: particle density; Pt: total porosity; Ksat: saturated hydraulic conductivity  
184 N: number of samples; Min.: minimum value; Max.: maximum value; SD: standard deviation; CV: coefficient  
185 of variation  
186

187 According to the classification proposed by [21], a parameter is considered to be low  
188 in terms of variability if the CV<12%, moderately variable when 12% < CV<60% and highly  
189 variable when CV>60%. In this study, the CVs for sand, BD, and Pd were less than 12%,  
190 indicating that these variables had low variability within the field. On the other hand, Clay  
191 and Pt, had CV between 12 and 60%, indicating moderate variability while Ksat had  
192 CV>100%, indicating very high variability. Similar studies have also reported low CV for  
193 sand [10] and BD [10, 11]. [10] found moderate CV for clay content. For Ksat, the result  
194 agrees with the findings of [19] and [20] who reported high variability of Ksat. In this study,  
195 the high variability of Ksat may be attributed to differences in soil pore geometry as a result  
196 of variable soil disturbance during land preparation. Certain sampling points may be  
197 characterized by biopores created by soil organisms and plant roots, thus increasing the water  
198 movement.

199 The frequency and normal distribution curves for the variables are shown in Figure 2.  
200 Only the logarithm transformed Ksat (LnKsat) had positive skewness, showing skewness to

201 the right, while other variables sand, clay, BD, Pd and Pt had negative skewness (Table 1),  
 202 showing skewness to the left (Figure 2). [22] stated that where a variable shows symmetry to  
 203 either right



204  
 205 Figure 2. Frequency and normal distribution curve of the selected soil physical properties of  
 206 the field.

207

208 or left, there is the tendency of high frequency of values below or above mean, respectively.  
 209 In this study, sand, clay, BD, Pd and Pt had high frequency of values above the mean. [11] in  
 210 a study on spatial variability of physical properties under land use change reported negative  
 211 and positive skewness for BD and Pt, respectively. According to [23], for a normal  
 212 distribution, the kurtosis coefficient must be zero, and values between +2 and -2 are accepted.  
 213 In this study, only the kurtosis values for clay, BD and Pt were within the acceptable limit. In  
 214 addition, the negative kurtosis for BD and Pt (Table 2) indicates that the curves were  
 215 platykurtic, showing the distribution was flatter than normal. Whereas the positive kurtosis

216 for clay indicates that the data was leptokurtic, that is, the distribution was narrower than  
217 normal (Figure 2). Other researchers [e.g. 24, 11] have also reported this behavior.

### 218 ***3.2 Relationships between soil physical properties***

219 The relationships between sand, clay, BD, Pd, Pt and LnKsat are presented in Table 2.  
220 There was significant positive correlation between Ks and sand content. Total porosity (Pt)  
221 had negative and significant correlation with BD whereas the correlation was positive with  
222 Pd. Sand had negative and significant correlation with clay content. The basis of the positive  
223 relationship between soil Ksat and sand content is direct; that is, higher Ksat values are  
224 associated with coarser rather than finer textured soil. In addition, high sand content indicates  
225 more macropore or transmission pores, hence increased water conductivity. Total porosity  
226 has an inverse relationship with bulk density, thus the confirmation obtained here. On the  
227 other hand, an increase in particle density indicates more pores, especially micropores and  
228 hence contributes to total pores. A soil having a more of sand will definitely have low content  
229 of clay which is a function of parent material from which the soil is formed.

230

### 231 ***3.3 Spatial variability and mapping of soil physical properties***

232 Table 3 and Figure 3 show the results of the geostatistical analysis of the measured  
233 soil physical properties. Sand and clay showed pure nugget effect (Figure 3 a and b); BD, Pd,  
234 and Pt were fitted to Gaussian model (Figure 3c, d and e) while LnKsat was fitted to spherical  
235 model (Figure 3f), with the coefficient of determination ( $R^2$ ) ranging from 0.104 (sand) to  
236 0.947 (LnKs). Other researchers [e.g. 10, 24, 25, 26, 11] have reported these models for soil  
237 physical properties. The nugget effect or the semivariance at separation distance of zero ( $h =$   
238  $0$ ) ranged between  $0.00 (\text{cm}^3 \text{cm}^{-3})^2$  (from Pt) and  $5.6 (\%)^2$  (from sand). According to [27],  
239 the range is a function of field and experimental variability, or random variability that is  
240 undetectable at the scale of sampling. Except for sand and clay, the close to zero nugget from

241 other variables is an indication of very smooth spatial continuity between neighbouring  
 242 points. The sand and clay content that had high nugget effect compared to other variables  
 243 indicates high discontinuity

244 Table 2. Results of Pearson correlation test between the soil physical properties.

Property	LnKs	BD	Pd	Pt	Sand	Clay
LnKs	1	0.122	-0.054	-0.138	0.215*	-0.100
BD		1	-0.097	-0.879**	0.071	-0.131
Pd			1	0.555**	0.027	0.103
Pt				1	-0.044	0.151
Sand					1	-0.310**
Clay						1

245 BD: bulk density; Pd: particle density; Pt: total porosity; LnKs: log transformed saturated hydraulic conductivity

246 \*\*Correlation is significant at the 0.01 level (2-tailed).

247 \*Correlation is significant at the 0.05 level (2-tailed).

248

249 among samples. [28] stated that the higher the nugget effect, the greater the discontinuity in  
 250 samples. As the separation distance ( $h$ ) increases, the semivariance increases to a more or less  
 251 constant value, known as the sill or total semivariance. The sill values ranged from 0.02 ( $\text{cm}^3$   
 252  $\text{cm}^{-3}$ )<sup>2</sup> (Pt) and 5.60 (%)<sup>2</sup> (sand). The ranges of spatial dependencies vary between 214 and  
 253 511 m, indicating that the optimum sampling interval varies greatly among the different soil  
 254 properties [10]. The sand and clay content that showed small range (214 m) of spatial  
 255 dependence indicates that spatial continuity diminishes rapidly over a short distance. The  
 256 value of semi-variogram range of the soil physical properties obtained in this study were not  
 257 in agreement with the range obtained in previous studies [e.g. 25, 26, 29]. Differences in soil,  
 258 land use type, cropping and management systems in the different regions may account for  
 259 these contrasting results.

260 Furthermore, the resulting semivariograms indicate strong spatial dependencies (SSD)  
 261 for BD, Pd and Pt. The structural variance also showed moderate spatial dependence for Ksat  
 262 and weak spatial dependence for sand and clay. These results indicate that the distribution of  
 263 the soil properties in space is not random. Strong spatial dependent in soil properties is an  
 264 indication that such properties are controlled by variability in intrinsic soil properties such as  
 265 geology, soil  
 266 Table 3. Fitted models and estimated parameters of the experimental semivariograms of soil  
 267 physical properties of the field.


Var.	Model	$C_o$	$C_o+C$	$A_o$	$C_o/(C_o+C)$	Spatial dependence	$R^2$	MAE	MSE
Sand	Nugget effect	5.600	5.60	214.3	1.00	WSD	0.104	0.620	0.553
Clay	Nugget effect	2.170	2.17	214.3	1.00	WSD	0.596	0.304	0.139
BD	Gaussian	0.020	0.07	510.9	0.23	SSD	0.833	0.046	0.003
Pt	Gaussian	0.003	0.02	510.9	0.13	SSD	0.900	0.020	0.001
Pd	Gaussian	0.013	0.05	510.9	0.25	SSD	0.560	0.021	0.001
LnKs	Spherical	0.768	1.83	410.9	0.42	MSD	0.947	0.498	0.306

268 BD: bulk density,  $g\ cm^{-3}$ ; Pd: particle density,  $g\ cm^{-3}$ ; Pt: total porosity,  $cm^3\ cm^{-3}$ ; LnKs: log transformed  
 269 saturated hydraulic conductivity,  $cm\ h^{-1}$   
 270  $C_o$ : nugget effect;  $C_o+C$ : sill;  $A_o$ : spatial range, m; SSD: strong spatial dependence; MSD: moderate spatial  
 271 dependence; WSD: weak spatial dependence  
 272  $R^2$ : coefficient of determination; MAE: mean absolute error; MSE: mean square error.  
 273

274 forming factors, texture and so on [30], whereas moderate and weak spatial dependence could  
 275 be due to management such as land use, tillage, cropping system, irrigation, among others.

276 By using the kriging algorithm of the geospatial analyst tool in ArcGIS, the contour  
 277 maps of the individual soil property are shown in Figures 4-8. The visualization of the  
 278 distribution maps showed that the soil varies in terms of physical properties, that is  
 279 heterogeneity, indicating that the distribution of the variables are strongly influenced by both  
 280 factors including geology, management practices, soil texture, among others. Figure 4 shows  
 281 the kriged contour map of the spatial variability and classification of the sand content of the


282 field. For the cowpea plot, it was observed that there was slightly high sand content. Also for  
 283 sole maize plot, there was slightly high sand content. For maize/cassava intercrop, there was  
 284 moderately high sand content. Figure 5 shows the kriged contour map of the spatial  
 285 variability and classification of the clay content. For the cowpea plot, the kriged contour map  
 286 showed that there was very low to low clay content in the northeastern region of the map. It  
 287 was noted that for sole maize plot, there was low clay




288  
 289 Figure 3. Semivariogram of a) sand content, b) clay content, c) soil bulk density (BD), d)  
 290 total porosity (Pt), e) particle density (Pd), and f) log transformed saturated hydraulic  
 291 conductivity (LnKs) of the field.

292


293 content due to inherent soil factors such as soil type and environmental factor. For  
294 maize/cassava, it was observed that there was very low clay content in this area of the field.  
295 The differences in the sand and clay contents are attributed to geology and intrinsic soil  
296 forming factors and the differences in these textural properties have implications in terms of  
297 pore space, water and nutrient retention and availability. Figure 6 shows the kriged contour  
298 map of the




299  
 300 Figure 4. Kriged contour map showing the spatial variability and classification of the sand  
 301 content of the field.




302  
 303 Figure 5. Kriged contour map showing the spatial variability and classification of the clay  
 304 content.



305  
 306 Figure 6. Kriged contour map showing the spatial variability and classification of the soil  
 307 bulk density (BD) of the field.



308  
 309 Figure 7. Kriged contour map showing the spatial variability and classification of the soil  
 310 total porosity (Pt) of the field.



311

312 Figure 8. Kriged contour map showing the spatial variability and classification of the soil  
 313 saturated hydraulic conductivity (Ksat) of sole maize and maize/cassava intercrop area of the  
 314 field.

315 variability and classification of the soil bulk density (BD) of the field. For the cowpea plot, it  
316 shows that there was low BD. Also from the sole maize plot, it was observed that there was  
317 low BD. The low bulk density indicates that the degree of compaction is low due to recent  
318 ploughing, harrowing and ridging operations conducted on the soil. For maize/cassava  
319 intercrop, the bulk density (BD) was medium (a bit higher) compared to cowpea and maize  
320 plots which may be attributed to crop intensification. The higher sand content in this region is  
321 also an avenue for the increased BD as more pore volume is available for compression.  
322 Figure 7 shows the kriged contour map of the spatial variability and classification of the soil  
323 total porosity (Pt) of the field. For both cowpea and sole maize plots, the total porosity (Pt) is  
324 classified as high. The high Pt observed may be as a result of low bulk density which is  
325 attributed to better aggregation and improved pore space. Conversely, maize/cassava  
326 intercrop had Pt classified as medium to low. This may be attributed to the relatively higher  
327 BD due to crop intensification.

328 Figure 8 shows kriged contour map of the spatial variability and classification of the  
329 soil saturated hydraulic conductivity (Ksat) for sole maize and maize/cassava intercrop only.  
330 For sole maize plot, the Ks is classified as moderate to high while it was classified as high for  
331 maize/cassava intercrop. The high Ks observed in maize/cassava intercrop may be due to  
332 high volume of macropore due to high sand content. The saturated hydraulic conductivity is a  
333 dynamic property of soil and its behavior is determined by the degree of compaction that the  
334 soil offers [31] as well as the quantity and continuity of pores, mainly macro spores.

335 The results of test of cross-validation of the kriging procedure checked using  
336 performance parameters of MAE and MSE are shown in Table 3. While the MAE indicates  
337 the bias, the MSE determines the prediction accuracy (Utset et al. 2000). Both the MAE and  
338 MSE values are very low, indicating that the kriging procedure was acceptable. Regardless of  
339 what factors caused the spatial variability observed, the magnitude of the soil properties may

340 be expected to influence the spatial distribution of crop growth and yield, thus having  
341 considerable implications regarding the implementation of soil sampling schemes and site-  
342 specific management practices.

343

## 344 **CONCLUSIONS**

345 The geostatistical methods showed spatial variability of the soil physical properties  
346 across the field. The variability of these properties is not random, revealing weak to strong  
347 spatial dependence.

348 The BD, Pd, Pt and Ksat could be well described using either Gaussian or spherical  
349 models. The semivariogram for sand and clay contents shows a small range of spatial  
350 dependence and purely nugget effect.

351 Crop intensification of maize/cassava intercrop influenced soil physical properties.

352 Spatial variability of soil physical properties across the field is attributed to a  
353 combination of previous sand mining activities and farming practices, parent material, and  
354 weather conditions.

355 The documentation of these physical properties in field scale distribution maps will  
356 allow derivation of zones of physical and mechanical sensitivity. This will further help define  
357 management zones, which can be combined with minimum soil samples to provide a more  
358 accurate prediction of spatial variability of soil properties for site-specific soil management.

359

## 360 **REFERENCES**

- 361 [1] Jung, W.K., Kitchen, N.R., Sudduth, K.H., Anderson, S.H. (2006). Spatial characteristics  
362 of claypan soil properties in an agricultural field. *Soil Sci. Soc. Am. J.*, 70: 1387-1397.  
363 [2] Babalola, O. (1978). Spatial variability of soil water properties of tropical soils of Nigeria.  
364 *Soil Science*, 126: 269-279.  
365 [3] Fergurson, R.B., Hergert, G.W., Schepers, J.S., Gotway, C.A., Cation, J.E., Peterson,  
366 T.A. (2002). Site-specific nitrogen management of irrigated maize: Yields and soil  
367 residual nitrogen effects. *Soil Sci. Soc. Am. J.*, 66: 544-553.

- 368 [4] Davis, J.C. (1986). *Statistics and Data Analysis in Geology*, 2nd Ed. Wiley & Sons, New  
369 York, NY.
- 370 [5] Webster, R., Oliver, M.A. (2001). *Geostatistics for Environmental Scientists*. Wiley &  
371 Sons, New York, NY.
- 372 [6] Corwin, D.L., Lesch, S.M., Shouse, P.J., Sopper, R., Ayers, J.E. (2003). Identifying soil  
373 properties that influence cotton yield using soil sampling directed by apparent soil  
374 electrical conductivity. *Agron. J.*, 95: 352-364.
- 375 [7] Mueller, T.G., Hartsock, N.J., Stombaugh, T.S., Shearer, S.A., Cornelius, P.L., Barnhise,  
376 R.I. (2003). Soil electrical conductivity map variability in limestone soil overlain by  
377 loess. *Agron. J.*, 95: 496-507.
- 378 [8] Corwin, D.L., Lesch, S.M. (2005). Survey protocols for characterizing soil spatial  
379 variability with apparent soil electrical conductivity: I. survey protocols. *Comput.  
380 Electron. Agric.*, 46: 103-133.
- 381 [9] Traugmar, E.B, Yost, R.S., Uehara, G. (1985). Application of geostatistics to spatial  
382 variability of soil properties. *Advances in Agronomy*, 38: 45-94.
- 383 [10] Jabro, J.D., Stevens, B.W., Evans, R.G. (2006). Spatial relationships among soil physical  
384 properties in a grass-alfalfa hay field. *Soil Science*, 171(9): 719-727.
- 385 [11] Awe, G.O. (2017). Land use change and crop production: Spatial variability of some soil  
386 physical and hydraulic properties of an Alfisol grown to cowpea (*Vigna unguiculata*).  
387 *American-Eurasian J. Agric. & Environ. Sci.*, 17(1): 85-98.
- 388 [12] EMBRAPA. (2011). Centro Nacional de Pesquisa de Solos (Rio de Janeiro, RJ). Manual  
389 de  
390 métodos de análise de solo. 2. Ed. Reviewed. Rio de Janeiro, 230pp.
- 391 [13] Blake, G.R., K.H. Hartge (1986). Bulk density, in: Klute, A. (Ed.), *Methods of Soil  
392 Analysis: Part 1, Physical and Mineralogical Methods*, 2<sup>nd</sup> edn, AS. SSSA, Madison,  
393 USA, 363-375.
- 394 [14] Gubiani, P.I., Reinert, D.J., Reichert, J.M. (2006). Metodo alternative para determinacao  
395 da densidade de particulars do solo – exatidao, precisao, e tempo de processamento.  
396 *Ciencia Rural*, 36(2): 664-668. (Abstract in English).
- 397 [15] Klute, A., Dirksen, C. (1986). Hydraulic conductivity and diffusivity: laboratory  
398 methods, in: Klute, A. (Ed.), *Methods of Soil Analysis. Part 1. Physical and  
399 Mineralogical Methods*, 2nd ed. *Agron. Monogr.* 9. ASA, Madison, WI, pp. 687–734.
- 400 [16] Nielsen, D.R., Wendroth, O. (2003). *Spatial and temporal statistics: sampling field soils  
401 and their vegetation*. Catena Verlag, Reiskirchen, p 398.
- 402 [17] Cambardella, C.A., Moorman, T.B., Novak, J.M., Parkin, T B., Karlen, D L., Turco,  
403 R.F.,  
404 Konopka, A.E. (1994). Field scale variability of soil properties in Central Iowa soils.  
405 *Sci. Soc. Am. J.*, 58: 1501-1511.
- 406 [18] Iqbal, J., Thomasson, J.A., Jenkins, J.N., Owens, P.R., Whisler, F.D. (2005). Spatial  
407 variability analysis of soil physical properties of alluvial soils. *Soil Sci. Soc. Am. J.*, 69:  
408 1338-1350.
- 409 [19] Shukla, M.K., Slater, B.K., Lal, R., Cepuder, P. (2004). Spatial variability of soil  
410 properties and potential management classification of a chernozemic field in lower  
411 Austria. *Soil Sci.*, 169: 852–860.
- 412 [20] Wang, Y.Q., Shao, M.A. (2013). Spatial variability of soil physical properties in a region  
413 of the Loess Plateau of PR China subject to wind and water erosion. *Land Degrad.  
414 Develop.*, 24: 296-304.

- 415 [21] Warrick, A.W., Nielsen, D.R. (1980). Spatial variability of soil physical properties in the  
416 field. In application of Soil Physics. Ed. D. Hillel. 319-324.
- 417 [22] Souza, L.C. de (1999). Variabilidade especial da salinidade de um solo aluvial no semi  
418 arido paraibano. 77p
- 419 [23] Ortiz, J.L. (2003). Emprego do geoprocessamento no estudo da relação entre potencial  
420 produtivo de um povoamento de eucalipto e atributos do solo e do relevo. 2003. 205pp.  
421 Dissertação (Mestrado) - Escola Superior de Agricultura "Luiz de Queiroz",  
422 Universidade de São Paulo, Piracicaba, Brasil. (Abstract in English).
- 423 [24] Carvalho, L.A.I, Meurer C.A, Silva J., Centurion, J.F. (2010). Spatial variability of soil  
424 properties in two managements in sugarcane crop. Eng. Agric., 32(1): 60-68.
- 425 [25] Cavalcante, E.G.S., Alves, M.C., Souza, Z.M., Pereira, G.T. (2011). Variabilidade  
426 especial dos atributos físicos do solo sob diferentes usos e manejos. R. Bras. Eng. Agri.  
427 Amb., 15(3): 237-243 (Abstract in English).
- 428 [26] Santos, D., Souza, E.G., Nobrega, L.H.P., Bazz, C.L., Goncalves Junior, A.C. (2012).  
429 Variabilidade especial de atributos físicos de um Latossolo Vermelho após cultivo de  
430 soja. R. Bras. Eng. Agri. Amb., 16(8): 843-848 (Abstract in English).
- 431 [27] Webster, R., Oliver, M.A. (1992). Sample adequately to estimate variograms of soil  
432 properties. J. Soil Sci., 43: 177-192.
- 433 [28] Vieira, S.R. (2000). Geoestatística em estudos de variabilidade especial do solo. In:  
434 Novais, R.F., Alvarez, V.H., Schaefer, C.E.G.R. (Eds). Tópicos em ciência do solo.  
435 Viçosa: Sociedade Brasileira de Ciência do Solo, 1: 1-53. (Abstract in English).
- 436 [29] Schaffrath, V.R., Tormena, C.A., Fidalski, J., Goncalves, A.C.A. (2008). Variabilidade e  
437 correlação especial de propriedades físicas de solo sob plantio direto e preparo  
438 convencional. R. Bras. Ci Solo, 32: 1369-1377 (Abstract in English).
- 439 [30] Yasrebi, J., Saffari, M., HamedFathi, N., Emadi, M., Baghernejad, M. (2008). Spatial  
440 variability of soil fertility properties for precision agriculture in southern Iran. J. Appl.  
441 Sci., 8: 1642-1650.
- 442 [31] Reichert, J.M., Suzuki, A.S., Reinert, D.J., Horn, R. and Hakasson, I. (2009). Reference  
443 bulk density and critical degree of compactness for no-till crop production in subtropical  
444 highly weathered soils. Soil and Tillage Research. 102: 242-254.