*Original Research Article*

# Effect of the foliar spray of some growth regulators with the fertilization by Fe and Zn on the yield and quality of Lentil grown in sandy soil


## ABSTRACT

**Aims:** A field study was carried out under the sandy soil conditions about the effect of some plant growth regulators (PGR) on the yield and quality of lintel (*Lens culinaris Medikus*).
**Study design:** a complete randomized block design.
**Place and Duration of Study:** winter seasons of 2016/2017 – 2017/2018 at the Ismailia Agricultural Research Station, Egypt.
**Methodology:** Lintel grains were inoculated by efficient strained *Rhizobium Leguminosarum*. Sowings were performed in Dec. 2016, 2017. Foliar spray of the PGR treatments at the recommended doses included the gibberellic acid ($GA_3$, 100 ppm), salicylic acid (SA, 100 ppm), K-humate (6000 ppm) and K-silicate ($K_2SiO_3$, 200) accompanied by the fertilization by EDTA chelated Fe (60 ppm) and/or Zn (100 ppm). Foliar spray was carried out 30, 45, and 60 days after sowing.
**Results:** For the estimated yield components and growth parameters, the effect of Zn treatments was more pronounced than Fe except for the plant height and the number of branches/plant. The Fe and Zn foliar spray significantly increased the lentil seed yield by 20.8% and 28.8%, respectively, compared with control. All treatments significantly increased the protein (%) and N (g/kg) in seeds compared with the control. The humate combined with the Fe or Zn (Fe + H and Zn + H) showed the most significant increase in the Fe in the seeds (by 54.89 and 100.16%, respectively). Combination between the silicate with Fe (Fe + Si) increased the Zn content seed more than the silicate with Zn (Zn + Si).
**Conclusion:** The seed yield (kg/ha) was significantly increased compared to the control in the order: Zn + $GA_3$ > Zn + Si > Zn + H > Fe + $GA_3$ > Fe + H > Fe + Si > Zn (Cont.) > Fe (Cont.) > Fe + SA > Zn + SA.



***Keywords:*** *Gibberellic; Humate; Lentil; Salicylic; Sandy Soil; Silicate*


## 1. INTRODUCTION

The reclaimed sandy soils are promising for the agricultural expansion. Improving their nutritional statue increase the total production through a good agronomic management. The area grown by lentil (*Lens culinaris Medikus*) is being increased in the reclaimed soils. It is one of the most important leguminous winter crops grown in Egypt. Lentil growing in the arid and semi-arid zones of the Mediterranean region experiences multiple environmental stresses such as drought, salinity, and heat. The proper nutrients supply and the water management are of the major constraints in increasing the lentil yield [1,2].

Lentil has the third-highest level of protein, after soybean and hemp. It contains dietary fibre, vitamin B1, and minerals. It is also a good source of iron (Fe) [3]. Lentil production fluctuates mainly due to the cultivation of low yielding and environment sensitive genotypes [4]. Seed germination is the most sensitive stage to stress and affects the final crop yield. The reduction of growth under stress conditions affects the cell division, restricts the amount of the plant available water, and changes the balance of the plant endogenous levels of phytohormones [5].

**Comment [F1]:** Insert the exact GPRS location of the study site.

**Comment [F2]:** What are your recommendations from your study.

**Comment [F3]:** Re-write this sentence to bring out its meaning clearly.

**Comment [F4]:** Insert citation.

**Comment [F5]:** Insert citation.

**Comment [F6]:** Insert citation.

**Comment [F7]:** Insert citation.

**Comment [F8]:** Insert citation.

UNDER PEER REVIEW

Plant hormones are involved in the induction of plant stress responses. A decreased cytokinin and gibberellic acid (GA3) and increased abscisic acid contents have been observed in salt-stressed plants [6]. To mitigate the adverse effect of such environmental stresses, the exogenous application of the plant growth regulators (PGRs) has obtained a substantial interest. The PGRs like auxins and gibberellins are organic substances that regulate intracellular processes to improve the plant growth [7]. The choice of the compound depends on the environmental condition, plant species and facilities available. The seed priming with the gibberellic acid (GA3), salicylic acid (SA) and ascorbic acid (ASC) has been found to increase the germination and seedling characteristics of lentil aged seeds [3].

The exogenously applied PGRs protect the plant against the water stress injury and retard the proline synthesis and transformations [8]. It may be a management tool improves the agricultural crops tolerance against stress and accelerates a potential crop yield in near future [9].

Gibberellins (especially GA3) are the most important valuable compound for enhancing the productivity of commercial crops and improving most of the yield components. They are natural penta-cyclic di-terpenoid carboxylic acids. They are associated with various plant growth and development processes such as seed germination, stem elongation, leaf expansion, floral organ development, reduced time to flowering, increased flower number and size and induction of some hydrolytic enzymes in the aleurone of cereal grains. Genes participating in both GA biosynthesis and response pathways may be involved [10].

The cereal aleurone layer constitutes an ideal system with a single cell type, a well characterized hormone signal (GA3). During germination, the aleurone layer synthesizes and secretes hydrolases to the starchy endosperm to start and sustain reserve mobilization. Almost all of the aleurone membrane systems participate in elaborating the response to GA3. There is evidence that suggests that GA3 perception occurs at the plasma membrane [10].

The endogenous GA content in maize, poplar, and sorghum, might be promoted by the exogenous GA3 that contribute to the increased plant growth [11]. Lentil is well adapted to semiarid plains regions where the soil profile is often not fully recharged with water. Lentil has a relatively shallow root system 0.6 m according to the genotypes. Increasing the concentrations of the phytohormones GA3 could reverse the adverse effects of water and salt stress on lentil [5,12].

Salicylic acid (SA) is a common hormone-like endogenous PGR and a natural signalling molecule. It is involved in the plant defence response against pathogen infection, biotic and environmental stress by influencing various physiological processes and biochemical reactions. The foliar application of the SA on sunflower has been enhanced the biomass production, photosynthetic pigments and the photosynthetic process [13].

Salicylic acid (SA) is distributed in the whole plant kingdom and has been characterized in 36 plants. In the plants, such as rice, crabgrass, barley and soybean the level of salicylic acid is ~1 µg/g fresh mass. It shows diverse regulatory roles in the plant metabolism of plant growth, development, seed germination, fruit yield, ion uptake and transport, photosynthetic rate, stomatal conductance and transpiration. It may contribute in regulating the antioxidant enzyme activity and gene expression [4].

Pre-treatment with SA improved the majority of physiological and morphological parameters in capsicum seedling and plants [14]. The foliar application of SA with farmyard manure during the vegetative and reproductive stages gave the highest lentil grain yield [12]. The proline metabolism in lentil was significantly altered maintaining the accumulation of higher levels of free proline and supporting its protection from salt stress [13].

The use of PGRs must be under strict control because they can provoke several diseases. The use of a silicon adjuvant improved growth regulator uptake by cotton and reduced the need of reapplication by at least 50% [15]. Foliar spray of Silicon alone at 50-ppm may result a bad influence on the human body [16]. The humic acids play a role similar to the PGRs [17,18].

Additionally, legumes have high Fe requirement for the heme components of the haemoglobin, nodule formation, enzymes like nitrogenase complex and hydrogenases and for ferredoxin electron carrier. Caixian has found that a reduced rate of nitrogenase activity has been observed in the Fe limited peanut nodules, indicating a possible direct limitation by Fe deficiency on nodule function [19]. Similar results were obtained previously [20-22].

Zinc plays a role in the photosynthesis in leaves and the carbohydrate metabolism and their conversion of sugar to starch [23]. Zinc is constituent of several plant enzymes that contribute to its rate of metabolism and protein synthesis [24]. Zinc deficient plant produced lower seed yields and reduced the availability of seeds by ~ 50% compared to plant with sufficient levels of Zn [20,25].

The present study aims to study the variation in the yield and quality of lintel (*Lens culinaris Medikus*) in sandy soil under the effect of the foliar spray of gibberellic acid (GA3), salicylic acid (SA), K-humate and K-silicate with fertilization by Fe and Zn.

**Comment [F9]:** Insert citation

**Comment [F10]:** Insert citation.

**Comment [F11]:** Insert citation.

**Comment [F12]:** Insert citation.

**Comment [F13]:** Insert citation.

**Comment [F14]:** Delete.

**Comment [F15]:** Insert citation.

**Comment [F16]:** Insert citation.

**Comment [F17]:** Insert citation.

**Comment [F18]:** Delete.

**Comment [F19]:** Insert citation.

**Comment [F20]:** Insert citation.

**Comment [F21]:** Ensure the citation numbers are serialized from the lowest to highest.

**Comment [F22]:** Insert citation.

UNDER PEER REVIEW

## 2. MATERIAL AND METHODS

A field experiment was carried out during the winter seasons of 2016/2017 – 2017/2018 under the conditions of sandy soil (Typic Torripsamment; Entisol [Arenosol AR] [26]) of the Ismailia Agricultural Research Station, Agricultural Research Center (ARC) - Egypt. Some of the physical and chemical properties of the experiment soil are shown in Table 1.

**Comment [F23]:** Provide the exact GPRS location of the study site.

**Table 1 Some characteristics of the experiment soil before cultivation**

| Character | | |
|---|---|---|
| **Particle size distribution (%)** | **Coarse sand** | 72.12 |
| | **Fine sand** | 14.32 |
| | **Silt** | 3.22 |
| | **Clay** | 10.34 |
| **Texture class** | | Sandy |
| **$CaCO_3$ (%)** | | 0.36 |
| **OM (%)** | | 0.23 |
| **pH (1:2.5 soil : water suspension)** | | 8.01 |
| **SP** | | 25.00 |
| **EC (dS/m) (1:5 soil : water extract)** | | 0.30 |
| **Available nutrients (mg/kg)** | **N** | 20.50 |
| | **P** | 2.01 |
| | **K** | 50.13 |

### 2.1. Planting

Many investigations were conducted in Egypt to develop promising lentil cultivars using local and exotic accessions, which were evaluated under variable conditions of Egyptian environments. Several cultivars such as Giza 9, Sinia 1 (Precoz) and Giza 370 were recommended due to multi-locations stability [1].

**Comment [F24]:** Refer to comment 9

Lintel grains Giza 9 variety were obtained from Field Crop Research Department, Field Crops Institute, Agriculture Research Centre/Egypt. Grains without visible defect, insect damage, and malformation were selected.

Lintel grains were inoculated by efficient strained *Rhizobium Leguminosarum* (supplied by the Department of Microbiology at SWERI/ARC) as follows: an amount of starch as adhesive glue material suitable for the grains quantity was wetted by 50 mL of warm water then mixed thoroughly with grains. The bacteria was mixed with the starch-wetted grains and dried for 90 min.

Sowings were performed in Dec. 2016, 2017 where two grains/hill were hand sown with 5 cm apart between hills. Eight treatments listed in Table 2 were applied in a complete randomized block design with three replicates in 24 plots (9 $m^2$ each). Fertilization by foliar spray was carried out 30, 45, and 60 days after sowing using the following treatments: 60 and 100 ppm of EDTA chelated Zn and/or Fe, respectively, 6000 ppm of K-humate, 200 ppm of K-silicate ($K_2SiO_3$), 100 ppm of SA and/or 100 ppm of GA3. Planting and the other agronomic practices were applied as recommendations of the Ministry of Agriculture.

**Comment [F25]:** Write in full.

Fertilizers were applied during the seedbed preparation as follows: 40 N units/ha as ammonium sulphate (20% N), 100 $P_2O_5$ units/ha as mono-calcium superphosphate (15.5% $P_2O_5$) and 100 $K_2O$ units/ha as potassium sulphate (48% $K_2O$).

At harvest time, (May 2017, 2018 - 150 days from sowing) a random sample of 10 plants from each plot was collected and air-dried. Some of the growth parameters such as the plant height (cm), number of branches/plant and the number of pods/plant were recorded. Yield components such as the weight of 1000 seeds (g), the seed yield (ton/ha) and the straw yield (kg/ha) have been calculated according to the total seed yield per plot area and the mean of the two seasons was recorded.

**Comment [F26]:** Which samoling procedure did you use and which formular did you use to calculate sample size.

UNDER PEER REVIEW

**Table 2 The treatments used in the experiment**

| Ser. No. | Treatment | Symbol |
|---|---|---|
| 1 | Control | Cont. |
| 2 | Fe (Control) | Fe (Cont.) |
| 3 | Fe + K-humate | Fe + H |
| 4 | Fe + Salicylic Acid | Fe + SA |
| 5 | Fe + Silicate | Fe + Si |
| 6 | Fe + Gibberellic Acid | Fe + GA3 |
| 7 | Zn (Control) | Zn (Cont.) |
| 8 | Zn + K-humate | Zn + H |
| 9 | Zn + Salicylic Acid | Zn + SA |
| 10 | Zn + Silicate | Zn + Si |
| 11 | Zn + Gibberellic Acid | Zn + GA3 |

### 2.2. Analysis of plant and soil samples

Lentil grains and straw were dried at 70ºC for 48h and ground. A half gram of ground grains or straw was wet digested using acid mixture (1:1 $H_2SO_4/HClO_4$) [27]. After harvesting, soil available N, P, K were extracted by 1% $K_2SO_4$, 0.5 N $NaHCO_3$, and 1 N $NH_4OAc$ (pH 7.0), respectively. Total percentage of N, P and K in digested plant samples and the available in the soil extracts were estimated by distillation using Kjeldahl apparatus, colorimetrically by UV-Vis. Spectrophotometer using $SnCl_2$ indicator and by flame photometer, respectively [28,29]. Protein percentage in grains was calculated as N% × 6.25. Iron Fe, Zn, and Si were measured for both grains and straw by Inductively Coupled Plasma Spectrometry (ICP-*Ultima 2 JY* Plasma).

### 2.2. Statistical Analysis

The one-way analysis of variance (ANOVA) was carried out using the Co-State software (Ver. 6.311) to determine the statistical significance of the treatment effects at a significance level LSD of 0.05 [30].

## 3. RESULTS AND DISCUSSION

### *3.1. Soil available nutrients*

Table 3 shows the variation in the soil available NPK resulted from the different treatments as mean of the two seasons. Foliar application of Fe and/or Zn significantly increased the soil available N by 235.3% and 34.8%, P by 355% and 235%, and K by 140%, respectively. Combination between the K-humate and Fe or Zn significantly increased the available P (by 115 and 65%) and K (by 103.1 and 42.5%), respectively, but no effect on the available N. Combination between the SA and Fe or Zn significantly increased the K only but affected the available N and P in an opposite direction. The K-silicate application significantly decreased the available N and almost no effect on the P or K when combined with the Fe. When combined with the Zn, it significantly increased the NPK. Foliar application of the GA3 combined with the Fe and/or Zn significantly increased the soil available N (by 34.8 and 51.2%), P (by 65 and 85%) and K (by 59.5 and 91.1%), respectively.

**Table 3 Available NPK nutrients in soil after harvest (mg/kg)**

| Treatment | Available NPK (mg/kg) | | |
|---|---|---|---|
| | N | P | K |
| Cont. | 20.1 | 2.1 | 65.0 |
| Fe (Cont.) | 23.1 | 2.5 | 70.0 |
| Fe + H | 21.2 | 2.7 | 78.0 |
| Fe + SA | 20.2 | 2.6 | 75.1 |
| Fe + Si | 19.1 | 2.5 | 78.5 |
| Fe + GA3 | 23.1 | 2.3 | 80.7 |

**Comment [F27]:** Indicate the study site, season , crop and the year of study.

UNDER PEER REVIEW

| | | | |
|---|---|---|---|
| **Zn (Cont.)** | 22.1 | 2.4 | 75.0 |
| **Zn + H** | 21.2 | 2.5 | 80.6 |
| **Zn + SA** | 23.0 | 2.6 | 70.0 |
| **Zn + Si** | 22.2 | 2.7 | 90.0 |
| **Zn + GA3** | 24.0 | 2.7 | 90.0 |
| **L. S. D. $_{5\%}$** | 2.26 | 0.68 | 8.35 |

### *3.2. Yield and yield components*

Table 4 shows some of the lentil yield components as affected by the studied treatments as mean of the two seasons. According to the LSD values at the *P* = .05 significance level and comparing with the control, the lentil plant height significantly increased by different treatments except for the foliar application by SA along with both Fe and Zn which was non-significant. Same treatment gave the lowest significance increase by 12.5% for the number of branches/plant. The Fe + Si treatment gave the most significant increase in the plant height and the number of branches/plant by 28.57% and 58.33%, respectively. The Fe and Fe + SA treatments were non-significant and the lowest significant increase effect on the number of pods/plant. The most significant increase by 42.86% was shown by the foliar application of silicate along with Fe or Zn (Fe + Si and Zn + Si treatments) and the GA3 with Zn. The variation in the 1000-seed weight was non-significant for all treatments.

**Table 4 Yield and yield components**

**Comment [F28]:** Indicate the study site, season ,crop and the year of study.

| Treatment | Plant Height (cm) | No. branches/plant | No. Pods/plant | 1000 seed wt. (g) | Seed yield (kg/ha) | Straw yield (kg/ha) |
|---|---|---|---|---|---|---|
| **Cont.** | 28 | 2.4 | 14 | 27.5 | 1250 | 3750 |
| **Fe (Cont.)** | 32 | 3.2 | 16 | 29.3 | 1510 | 4228 |
| **Fe + H** | 35 | 3.6 | 19 | 29.5 | 1650 | 4455 |
| **Fe + SA** | 29 | 2.7 | 15 | 28.0 | 1320 | 3960 |
| **Fe + Si** | 36 | 3.8 | 20 | 28.1 | 1630 | 4564 |
| **Fe + GA3** | 35 | 3.5 | 19 | 28.2 | 1700 | 4760 |
| **Zn (Cont.)** | 31 | 3.1 | 17 | 28.2 | 1610 | 4508 |
| **Zn + H** | 34 | 3.6 | 19 | 29.1 | 1720 | 4730 |
| **Zn + SA** | 29 | 2.7 | 17 | 27.5 | 1310 | 3930 |
| **Zn + Si** | 35 | 3.7 | 20 | 29.3 | 1800 | 5040 |
| **Zn + GA3** | 35 | 3.6 | 20 | 29.3 | 1850 | 5180 |
| **L.S.D. $_{5\%}$** | 2.1 | 0.3 | 2.1 | 3.6 | 99.2 | 280.5 |

Again, the foliar spray by SA along with both Fe and Zn was non-significant for both the seeds and straw yield (kg/ha) as yield parameters. The highest significant increase in the seeds and straw yield (kg/ha) was by 36, 44 and 48% and by 26.93, 34.4 and 38.13% resulting from the treatments Fe + GA3, Zn + Si and Zn + GA3, respectively.

Comparison between Fe and Zn different treatments indicated that the effect of Zn treatments is more pronounced than Fe for the estimated yield components except for the plant height and the number of branches/plant. The Fe and Zn foliar application significantly increased the lentil seed yield by 20.8% and 28.8%, respectively.

The significant effect of Zn on the lentil yield may be due to its role as a critical micronutrient for health plant function during essential stages of growth and development. Alloway (2008) reported that high Zn concentration have improved vigour seed compared to those with lower Zn [23]. The data obtained confirm with data obtained [20,31].

**Comment [F29]:** Decide on one method of writing citations, i.e. numerical numbers or names?

Application of humate combined with Fe and/or Zn increased the seed yield by 32% and 37.6%, respectively, compared to control. The mechanism of humate activity in plant growth simulation have been proposed as increasing cell membrane permeability, nutrient and oxygen uptake, respiration into cell of plant tissue, enhanced root, leaf and shoot growth and simulated

UNDER PEER REVIEW

germination. Plant shows improved physiological and metabolic processes and increased respiratory activity attributed to the inner venation of humic acid.

Combined application of Si with Fe and Zn significantly increased the lentil height, branches number and seed yield kg/ha by 30.4 and 44% for the Fe + Si and Zn + Si, respectively. Silicon affects the plant growth mechanisms, i.e. alleviate many abiotic stress, like lodging, drought, radiation, high temperature, salt, nutrient imbalance, increasing photosynthetic activity. Si is applied in several countries for increasing productivity and sustainable production. It was beneficial for barely, wheat, corn, sugar cane, cucumber, tomato and other crops. Silicon foliar application increase pea dry shoot and NPK contents, increased maize yield and its components at 100 and 200 ppm Si [32].

The treatments Fe + GA3 and Zn + GA3 showed the highest significant increase in the seeds yield (kg/ha) by 36 and 48%, respectively. The obtained data are in agreement with those obtained previously. Application of GA3 significantly affected the growth, yield, and yield components of maize grain yield [32].

### *3.3. Nutrient content in the lentil seeds and straw*

#### ***3.3.1. The NPK (g/kg) total content***

Data in Table 5 present the mean of the two seasons for the variation in the total NPK content in both lentil seeds and straw in addition to the seeds' protein. All treatments significantly increased the protein (%) and N (g/kg) in seeds compared with the control at a significance level *P = .05.* Individual application of Fe or Zn increased the protein (%) by 23.53 and 17.03%, and the N (g/kg) by 19.06 and 17.06%, respectively. Combination of SA with either Fe or Zn resulted in the most significant increase by 38.7 and 84.21% in the protein (%) and by 38.74 and 84.26%, respectively. The combined Zn different applications increased the protein (%) more significantly than the combined Fe applications.

**Table 5 Total NPK in lentil seeds and straw, (g/kg)**

**Comment [F30]:** Insert study site, season, crop and year of study.

| Treatment | Seeds, (g/kg) | | | | Straw, (g/kg) | | |
|---|---|---|---|---|---|---|---|
| | Protein % | N | P | K | N | P | K |
| **Cont.** | 19.38 | 31.00 | 2.10 | 5.10 | 8.10 | 1.20 | 11.2 |
| **Fe (Cont.)** | 23.94 | 38.30 | 3.95 | 6.14 | 8.06 | 1.20 | 12.42 |
| **Fe + H** | 26.04 | 41.66 | 4.45 | 6.14 | 9.41 | 1.05 | 7.33 |
| **Fe + SA** | 26.88 | 43.01 | 4.05 | 6.14 | 8.06 | 1.05 | 11.31 |
| **Fe + Si** | 24.57 | 39.31 | 4.05 | 7.49 | 6.72 | 1.05 | 13.19 |
| **Fe + GA3** | 24.36 | 38.98 | 4.00 | 7.33 | 9.41 | 1.13 | 7.80 |
| **Zn (Cont.)** | 22.68 | 36.29 | 3.63 | 6.01 | 10.08 | 0.85 | 5.04 |
| **Zn + H** | 29.40 | 47.04 | 4.15 | 5.30 | 9.41 | 1.13 | 5.6 |
| **Zn + SA** | 35.70 | 57.12 | 4.05 | 7.33 | 10.75 | 1.3 | 5.87 |
| **Zn + Si** | 30.24 | 48.38 | 3.55 | 5.87 | 8.74 | 0.80 | 7.33 |
| **Zn + GA3** | 28.14 | 45.02 | 4.25 | 6.14 | 11.42 | 1.10 | 5.73 |
| **L.S.D. $_{5\%}$** | 1.58 | 1.38 | n.s. | 1.03 | 1.38 | 0.41 | 1.72 |

The variation in the seeds' content of P (g/kg) was non-significant compared to the control. Foliar application of either Fe or Zn increased the P (g/kg) by 88.1 and 72.86%, respectively. The Fe + H and the Zn + GA3 treatments showed the maximum increase by 111.9 and 102.38%, respectively.

All the Fe treatments while only two Zn treatments (Zn + SA and Zn + GA3) that significantly increased the K (g/kg) in seeds. The Fe + Si treatment showed the most significant increase by 46.86% while the Zn + H treatment exhibited the minimum non-significant increase by 3.92% compared to the control.

For the straw, only three Zn treatments that significantly increased the N (g/kg) by 24.44%, 32.72% and 40.99% for the Zn, Zn + SA and Zn + GA3 treatments, respectively. Again, the variation



UNDER PEER REVIEW

in the P (g/kg) was non-significant and all treatments decreased it in the straw compared with the control. The Fe + Si was the sole treatment significantly increased the K (g/kg) in the straw by 17.77%. The Fe and Fe + SA treatments increased it by 0.97 and 0.98%, respectively. Other treatments significantly decreased the K (g/kg) in the straw and the least significant decrease by 55% due to the Zn treatment.

***The Fe, Si, and Zn (mg/kg) content***

Table 6 exhibits the content of the Fe, Si, and Zn (mg/kg) in the lentil seeds and straw as mean of the two seasons. Foliar application of the humate combined with the Fe or Zn (Fe + H and Zn + H) showed the most significant increase in the Fe in the seeds (by 54.89 and 100.16%, respectively) and the maximum non-significant increase in the straw (57.86 and 38.23%, respectively). The second most effective in enhancing the Fe content treatments were the Zn + GA3 for the seeds and the Fe + SA for the straw [33].

**Table 6 The effect of different treatments on the content of Fe, Si and Zn in the seeds and straw of lentil**

Comment [F31]: Insert study site, season and year of study.

| Treatment | Seeds (mg/kg) | | | Straw (mg/kg) | | |
|---|---|---|---|---|---|---|
| | Fe | Si | Zn | Fe | Si | Zn |
| **Cont.** | 57.0 | 46.0 | 33.0 | 66.0 | 48.0 | 15.0 |
| **Fe (Cont.)** | 78.1 | 49.7 | 38.0 | 138.9 | 75.2 | 20.4 |
| **Fe + H** | 117.7 | 113.0 | 44.4 | 72.8 | 56.4 | 29.2 |
| **Fe + SA** | 85.4 | 68.8 | 45.6 | 111.0 | 75.4 | 44.8 |
| **Fe + Si** | 98.6 | 47.9 | 80.4 | 95.6 | 55.6 | 22.0 |
| **Fe + GA3** | 76.3 | 51.2 | 43.2 | 79.2 | 34.6 | 10.8 |
| **Zn (Cont.)** | 82.4 | 49.3 | 35.2 | 96.9 | 56.8 | 24.8 |
| **Zn + H** | 152.1 | 53.7 | 42.8 | 121.6 | 75.3 | 30.8 |
| **Zn + SA** | 91.4 | 41.2 | 50.4 | 94.4 | 43.1 | 12.8 |
| **Zn + Si** | 90.8 | 42.8 | 34.8 | 97.9 | 50.2 | 14.0 |
| **Zn + GA3** | 107.6 | 44.3 | 47.2 | 107.8 | 113.8 | 36.8 |
| **L.S.D. $_{5\%}$** | 3.39 | 2.73 | 3.67 | 3.75 | 2.76 | 4.38 |

Foliar application of Fe or Zn significantly increased the Si (mg/kg) in the seeds and straw by 7.96, 7.15% and 56.56, 18.33%, respectively. Combination between the humate with either the Fe or Zn significantly increased the Si (mg/kg) by 145.74% and 16.65%, respectively, in the seeds' and by 17.44% and 56.79%, respectively, in the straw. The Si increased by the combined Fe treatments in the order Si < GA3 < SA for seeds and GA3 < Si < SA for the straw. However, it decreased by the combined Zn treatments in the order SA < Si < GA3 for seeds and by the SA for the straw.

Combination between the silicate and Fe (Fe + Si) increased the seed content of Zn more than the silicate and Zn (Zn + Si).

The foliar application of GA3, Si, and SA mediated the accumulation levels of some heavy metals in maize leaves. The chemical structure of the GA3, SA and silicates affects the absorption of cations (or anions) by plant tissues [20,32,34].

**4. Conclusion**

The variation in the yield and quality of lintel (*Lens culinaris Medikus*) resulting from the foliar spray of gibberellic acid (GA3), salicylic acid (SA), K-humate and K-silicate accompanying the fertilization by Fe and Zn was studied. The data as mean of the two seasons were recorded and statistically analysed. The sole application of either Fe or Zn treatment significantly increased the soil available NPK followed by the GA3 application with either Zn or Fe then the Zn + Si treatment compared to the control. Other treatments exhibited divers effect on the soil available NPK. Generally, almost all treatments significantly increased the available K. The available N decreased by the Fe treatments (+ H, + SA, + Si) while increased by the Zn treatments (+ SA, + Si).

The lentil seed yield (kg/ha) was enhanced by the Zn treatments more than the Fe treatments. Foliar application of SA with either Fe or Zn increased the seed yield but non-significantly. Other treatments significantly increased the lentil seed yield compared to the control in the following

UNDER PEER REVIEW

order: Zn + GA3 > Zn + Si > Zn + H > Fe + GA3 > Fe + H > Fe + Si > Zn (Cont.) > Fe (Cont.) > Fe + SA > Zn + SA.

Application of the humate combined with the Fe or Zn (Fe + H and Zn + H) showed the most significant increase in the Fe in the seeds by 54.89 and 100.16%, respectively. Foliar application of Fe or Zn significantly increased the Si (mg/kg) in the seeds and straw by 7.96, 7.15% and 56.56, 18.33%, respectively. The Si increased by the combined Fe treatments in the order Si < GA3 < SA for seeds. However, it decreased by the combined Zn treatments in the order SA < Si < GA3 for seeds and by the SA for the straw. Combination between the silicate and Fe (Fe + Si) increased the seed content of Zn more than the silicate and Zn (Zn + Si).

The foliar application of GA3, Si, and SA mediated the accumulation levels of some heavy metals in maize leaves. The chemical structure of the GA3, SA, and silicates affects the absorption of cations (or anions) by plant tissues.

## REFERENCES


[1] Abdel-Daiem M, Hassan MF, Hamdi YA, Abdel-Ghaffar AS. Nitrogen fixation and yield of faba bean, lentil and chickpea in response to selected agricultural practices in Egypt. In: Summer field R.J. (eds.) *World crops: Cool season food legumes.* Current Plant Science and Biotechnology in Agriculture, 5. Springer, Dordrecht, 1988.

[2] Biçer BT, Şakar D. Stability parameters in lentil. J. Cent. Eur. Agri. 2006;7(3), Prosinac 439–444.

[3] Pourazar K, Mirshekari B. Invigoration of lentil (*Lens culinaris* L .) seeds by hormonal priming with kinetin and gibberellic acid, ARPN. J. Agric. Biol. Sci. 2015;10, 324–329.

[4] Singh D, Singh CK, Taunk J, Tomar RSS, Chaturvedi AK, Gaikwad K, Pal M. Transcriptome analysis of lentil (*Lens culinaris Medikus*) in response to seedling drought stress. BMC Genomics. 2017;18, 206.

[5] Abdel CG, Al-Rawi IMT. Influence of varying gibberellic acid on growth and yield of three lentil (*lens culinaris Medik* sub species *culinaris*) cultivars grown under supplementary irrigation. Int. J. Environ. Water. 2012;1,137–153.

[6] Tuna ALC, Kaya M, Dikilitas D, Higgs D. The combined effects of gibberellic acid and salinity on some antioxidant enzyme activities, plant growth parameters and nutritional status in maize plants. Environ. Exp. Bot. 2008;62(1), 1–9.

[7] Hadi F, Bano A, Fuller MP. The improved phyto extraction of lead (Pb) and the growth of maize (*Zea mays* L.): the role of plant growth regulators (GA3 and IAA) and EDTA alone and in combinations. Chemosphere. 2010;80,457–462.

[8] Mehrabi A, Khavarinegad R, Safari Z, Kian S. The effect of gibberellic acid (GA3) on seed germination and early growth of lentil seedlings under salinity stress. Middle-East J. Sci. Res. 2014;19,995–1000.


**Comment [F32]:** Itallize all the journal names in the references.

**Comment [F33]:** Itallize.

UNDER PEER REVIEW

[9] Kawano T, Furuichi T. Salicylic Acid as a Defense-Related Plant Hormone: Roles of oxidative and calcium signaling paths in salicylic acid biology. (Hayat, S. and Ahmad, A. (eds.), Salicylic Acid - A Plant Hormon. Springer. 2007; 277–321.

[10] Bose SK, Yadav RK, Mishra S, Sangwan RS, Singh AK, Mishra B, Srivastava AK, Sangwan NS. Effect of gibberellic acid and calliterpenone on plant growth attributes, trichomes, essential oil biosynthesis and pathway gene expression in differential manner in *Mentha arvensis* L. Plant Physiol. Biochem. 2013;66,150–158.

[11] Du J, Yao Y, Ni Z, Sun Q. Cloning and characterization of an up-regulated GA 20-oxidase gene in hybrid maize. Prog. Nat. Sci. 2009;19,161–166.

[12] Janmohammadi M, Nasiri Y, Zandi H, Kor-Abdali M, Sabaghnia N. Effect of manure and foliar application of growth regulators on lentil (*Lens culinaris*) performance in semi-arid highland environment. Bot. Lith. 2014;20,99–108.

[13] Misra N, Saxena P. Effect of salicylic acid on proline metabolism in lentil grown under salinity stress. Plant Sci. 2009;177,181–189.

[14] Prabha D, Negi YK. Seed treatment with salicylic acid enhance drought tolerance in Capsicum. World J. Agric. Res. 2014;2(2),42–46.

[15] Echer FR, Rosolem CA. Plant growth regulator losses in cotton as affected by adjuvant and rain. Ciência Rural. 2012;42,2138–2144.

[16] El-Zohiri SS. Performance comparison of three alternatives for GA3 on growth, earliness, and total yield of Globe Artichoke. Middle East J. Appl. Sci. 2015;3,636–644.

[17] Blanchard JR, Farrell L. Guide to Sources for Agricultural and Biological Research, National Agricultural Library (U.S.), University of California Press. 1981.

[18] Tehranifar A, Atefe A. Effect of humic acid on nutrient uptake and physiological characteristics of fragaria ananassa'camarosa. VII. International Strawberry Symposium 1049. 2012.

[19] Caixian T, Robson AD, Dilworth MJ. A split-root experiment shows that iron is required for nodule initiation in *Lupinus angustifolius* L. New Physiologist. 1990;115(1),61-67.

[20] Zeidan MS, Hozayn M, Abd El-Salam MEE. Yield and Quality of Lentil as Affected by Micronutrient Deficiencies in Sandy Soils. J. Appl. Sci. Res. 2006;2(12),1342-1345.

[21] Ghasemian V, Ghalavand A. The effect of iron, zinc and manganese on quality and quantity of soybean seed. J. Phytol. 2011;2(11),73-79.

[22] Heidarian AR, Kord H, Mostafavi K, Lak AP, Mashhadi FA. Investigating Fe and Zn foliar application on yield and its components of soybean (*Glycine max* (L) Merr.) at different growth stages. J. Agric. Biotech. Sust. Devel. 2011;3(9),189-197.

[23] Alloway BJ. Zinc in soils and crop nutrition. IZA Publications. International Zinc Association, Brussels. 2004;1–116.

[24] Marschner H. Marschner's Mineral Nutrition of Higher Plants, Academic Press. 1982.

[25] Pandey N, Pathak GC, Sharma CP. Zinc is critically required for pollen function and fertilisation in lentil." J. Trace Elements in Med. Bio. 2006;20(2),89-96.

UNDER PEER REVIEW

[26] FAO. World Reference Base for Soil Resources. A framework for international classification, correlation and communication. Soil Taxonomy, Food and Agriculture Organization of the United Nations, Rome, Italy. 2014.

[27] Chapman HD, Pratt PF. *Methods of analysis for soils, plants and waters.* Division of Agricultural Sciences, University of California, Riverside. 1961.

[28] Black CA. Methods of soil analysis. Part I, American Society of Agronomy. Madison, Wisconsin, USA. 1965.

[29] Jackson ML. Soil Chemical Analysis. Prentice-Hall, Inc., Englewood Cliffs. NJ, USA. 1973.

[30] Gomez KA, Gomez AA. Statistical Procedures for Agriculture Research. A Wiley – Inter Science Publication, John Wiley and Sons, Inc. New York, USA. 1984.

[31] Okaz AMA, El–Gareib EA, Kadry Ay Negm W, Zahran FAF. Micronutrient application to lentil plants grown on newly reclaimed sandy soils. Proc. 6th Conf. Agron., Al–Azhar Univ., Cairo, Egypt, Sept. Vol. II, 1994;737-752.

[32] El-Kamar FA, Shaban KhA, Rashad RT. The efficiency of using salicylic acid, gibberellic acid and silicon, on the productivity of the maize plant (*Zea maize L.*) under salt- affected soil conditions. J. Soil Sci. Agric. Eng., Mansoura Univ. 2013;4 (10),1021-1035.

[33] Dogan Y, Togay Y, Togay N, Kulaz H. Effect of humic acid and phosphorus applications on the yield and yield components in lentil (*Lens culinaris medic.*). Legume Res., An International J. 2014;37(3),316-320.

[34] Rashad RT, Hussien RA. A comparison study on the effect of some growth regulators on the nutrients content of maize plant under salinity conditions. Ann. Agric. Sci. 2014;59(1),89–94.

UNDER PEER REVIEW