

GROWTH AND YIELD RESPONSES OF CABBAGE CULTIVARS AS INFLUENCED BY ORGANIC AND INORGANIC FERTILIZERS

ABSTRACT

An experiment was conducted at the Horticulture Farm of Sher-e-Bangla Agricultural University, Dhaka from November 2015 to February 2016 to evaluate the growth and yield responses of cabbage cultivars as influenced by organic and inorganic fertilizers. The experiment comprised of two different factors such as (1) three varieties viz. V_1 (Atlas 70), V_2 (Keifu 65) and V_3 (Autumn 60) and (2) Four different fertilizers viz. F_0 (Control), F_1 (Cow dung), F_2 (Poultry manure), and F_3 (Inorganic fertilizer). The experiment was set up in Randomized Complete Block Design with three replications. The experimental plot was fertilized as per treatment with organic and inorganic fertilizers. Among the varieties, Atlas 70 (V_1) achieved the highest results of Plant height (31.94 cm), Leaf length with petiole (32.00 cm), Stem length (4.194 cm), Diameter of head (20.24 cm), Weight of whole plant (2.23 kg/plant), gross yield (46.67 t/ha), marketable yield (45.29 t/ha) and Economic production (1.576 kg/plant) at the time of harvest. With the interaction effect of variety and fertilizer; V_1F_2 (Atlas 70 $\times$ Poultry manure) represented the highest Weight of whole plant (2.56 kg/plant), gross yield (62.14 t/ha), marketable yield (61.52 t/ha) and Economic production (1.85 kg/plant). Therefore, Atlas 70 coupled with poultry manure can be the most suitable for enhanced yield and economic production of cabbage.

Keywords: Growth, Yield, Cabbage, Cultivars, Organic, Inorganic, Fertilizers.

1. INTRODUCTION

Cabbage (*Brassica oleracea* L.) is ~~a one of the~~ cole crops, member of the Brassicaceae family. ~~Cabbage is an important fresh and processing vegetable crop in most of the countries of the world.~~ Cabbage ~~is a vegetable crop and~~ generally is grown in during Rabi season in Bangladesh. ~~Cabbage is an important fresh and processing vegetable crop in most of the countries of the world.~~ Cabbage is an important and nutritious winter leafy vegetable in Bangladesh. It contains a range of essential vitamins and minerals as well as small amount of protein and good caloric value (Haque, 2006). The productivity of cabbage per unit area is quite low as compared to the developed countries of the world ~~(provide some data)~~. Various factors such soil nutrient management, irrigation, variety, plant population per unit area, are involved for better growth of cabbage. Among the factors, suitable variety and nutrient supply is the important inputs for realizing higher cabbage yield and its nutrient content.

Variety is an important factor for successful crop yield. An improved variety represents a higher yield than a wild one. Generally nutrient requirement is determined by the variety of crops. High yielding variety requires more nutrients than the local or wild variety. Generally it depends on its vegetative and reproductive characters. And it was also mentioned that vegetable variety and history of fertilizer use are important factors to be considered in the development of a soil nutrient management program (Huang, 2006).

The cultivation of cabbage is required proper supply of plant nutrients. The requirement of these plants nutrients can be provided by applying inorganic fertilizer or organic manure or both. However, farmers are now showing interest in organic farming because of, they are more aware about the residual effect of chemical substances used in the crops field and environmental degradation. Besides, the excess application of inorganic fertilizer causes hazard to public health and to the environment. But the application of both organic and inorganic fertilizer combined, can increase the yield as well as keep the environment sound (Hsieh *et al.*, 1996). Considering the above factors, the present experiment was undertaken to identify the best variety that could be suggestive for growth of cabbage for the farmers of Bangladesh. In addition attempt was undertaken to determine the best organic fertilizer option for better growth of cabbage and to determine the combination of variety and fertilizer management of cabbage.

2. MATERIAL AND METHODS

2.1 Experimental Site

The experiment was conducted at the Horticultural farm of Sher-e-Bangla Agricultural University, Dhaka, Bangladesh from November 2015 to February 2016. The location of the experimental site was 23°74'N latitude and 90°35'E longitude and at an elevation of 8.2 m from sea level. The climate of experimental site was under the subtropical climate, characterized by three distinct seasons, the winter season from November to February and the pre-monsoon or hot season from March to April and the monsoon period from May to October. The soil of the experimental area belongs to the Modhupur Tract (AEZ No 28). It had shallow red-brown terrace soil. The selected plot was medium high land and the soil series was Tejgaon.

2.2 Planting Material

Three varieties were used as planting materials viz. (i) Atlas – 70, (ii) Keifu – 65 and (iii) Autumn – 60. Seeds of cabbage cultivars were used in the experiment and the seeds were collected from a commercial seed trader.

2.3 Organic and Inorganic Materials

Fertilizers (4 levels): F₀: Control, F₁: Cowdung at the rate of 15 t/ha, F₂: Poultry manure at the rate of 15 t/ha and F₃: Inorganic fertilizer- Urea at the rate of 330 kg/ha, TSP (Triple Super Phosphate) at the rate of 200 kg/ha, and MP (Murate of Potash) at the rate of 250 kg/ha.

2.4 Experimental Design and Treatments

The experiment was laid out in Randomized Complete Block Design with three replications. There were 12 treatment combinations such as F₀V₁, F₀V₂, F₀V₃, F₁V₁, F₁V₂, F₁V₃, F₂V₁, F₂V₂, F₂V₃, F₃V₁, F₃V₂ and F₃V₃. Total number of plots was 36 and the size of the each unit plot was 2.4 m × 1.6 m. The distance maintained between two blocks and two plots were 1.0 m and 0.5 m, respectively. Plant spacing 60 cm × 40 cm was maintained in this experiment.

2.5 Growth condition of Cabbage & Measurements of Parameters

Seedlings were grown following proper methods and all of the cultural practices were done properly. Application of manure and fertilizers were applied as per treatment. Healthy and uniform sized seedlings were transplanted in the main field. Intercultural practices were done as per requirements. For controlling leaf caterpillars Nogos @ 1 ml/L water were applied two times at an interval of 10 days starting soon after the appearance of infestation. All cabbage head were not matured at a same time, harvesting was done at 15 February to 02 March

76 .Different yield contributing data have been recorded from the mean of five harvested plants
77 which was selected at random of each unit plot of every harvesting stage.

78 2.6 Data Collection and Analysis


79 Five plants were randomly selected from each unit plot for the collection of data. The plants
80 in the outer rows and the extreme end of the middle rows were excluded from the random
81 selection to avoid the border effect. The height of the plants was measured from the ground
82 level to the tip of the highest leaves. To record the diameter, the cabbage heads were
83 sectioned vertically at the middle position and the horizontal distance from one side to the
84 other side of the widest part of the sectioned head was measured. The thickness of head
85 was measured as the vertical distance from the lower to the upper most leaves of head. The
86 data obtained for different parameters were statistically analyzed to find out the significance
87 difference of variety and different fertilizer application on yield and yield contributing
88 characters of cabbage. The mean values of all the characters were calculated and analysis
89 of variance was performing-performed by the 'F' (variance ratio) test. The significance of the
90 difference among the treatment combinations means was estimated by the Duncan's
91 Multiple Range Test (DMRT) at 5% level of probability (Gomez and Gomez 1984).

92 3. RESULTS AND DISCUSSION

93 3.1 Growth parameters

94 3.1.1 Plant height

96 Variety is an important factor considering plant height. Under the present study, plant height
97 was significantly influenced by different varieties of cabbage cultivar at different days after
98 transplanting (DAT) (Figure 1). Results showed that the cabbage cultivar Atlas 70 (V₁) was
99 evident for highest plant height at all growth stages. The tallest plant at 15, 30, 45 DAT and
100 at harvest were 26.81, 29.29, 30.88 and 31.94 cm respectively was obtained with Atlas 70
101 (V₁).




111 Fig.1: Plant height at different growth stages of three Cabbage Cultivars-cultivars

112 The competition in accordance with plant height among the cultivars the smallest plant was
113 demonstrated with Autumn 60 (V3) and the lowest plant height at 15, 30, 45 DAT and at

114 harvest were 23.83, 27.68, 29.58 and 31.04 cm respectively which was statistically identical
115 with Keifu 65 (V2) at 30, 45 DAT and at harvest respectively. This might be due to the
116 genetic variations among the varieties used under the present study. The varietal effect on
117 plant height was supported by Haque (2005).

118 Plant height was significantly affected by different manures and fertilizers under the present
119 study (Figure 2). It is evident that plant height was the highest with inorganic fertilizer (F_3) at
120 different growth stages of different varieties of cabbage cultivars. The highest plant height
121 was 26.00, 29.39, 31.03 and 32.55 cm at 15, 30, 45 DAT and at harvest respectively. On the
122 other hand, the lowest plant height (24.48, 26.98, 29.12 and 30.52 cm at 15, 30, 45 DAT and
123 at harvest, respectively) was with control treatment (F_0) which was statistically identical with
124 cow dung (F_1) treated crop at harvest. This result might be due to cause of rapid
125 performance on growth characters and rapid release of nutrients of inorganic fertilizer for
126 plant height where organic fertilizer has a slow nutrient release capacity that caused lower
127 plant height. Results under the present experiment on plant height were supported by Souza
128 *et al.* (2008).



140 **Fig 2: Effect of different fertilizer on plant height of different cabbage cultivar at**
141 **different growth stages**

142 Interaction effect of variety and different type manure and fertilizer affected plant height
143 significantly under the present study (Table 1). Different treatment combination viewed
144 different plant height at different days after transplanting (DAT). It was observed that highest
145 plant height was achieved with V_1F_3 and that was 28.30, 30.37, 32.00 and 33.11 cm at 15,
146 30, 45 DAT and at harvest respectively which was closely followed by V_1F_2 at 15, 30 DAT
147 and V_3F_3 at harvest. On the other hand the lowest plant height; 22.46, 25.60, 27.80 and
148 29.93 cm at 15, 30, 45 DAT and at harvest respectively was obtained with V_3F_0 which was
149 statistically identical with V_2F_1 and V_3F_1 at harvest.

151

152


153 **Table 1: Interaction effect of fertilizer management and three different cabbage**
 154 **cultivars on plant height at different growth stages**

Treatments	Plant height (cm)			
	15 DAT	30 DAT	45 DAT	At harvest
Interaction effect of variety and fertilizer				
V ₁ F ₀	25.97 bc	28.40 de	30.10 cd	31.12 de
V ₁ F ₁	26.03 bc	28.40 de	30.32 bc	31.51 cd
V ₁ F ₂	26.92 ab	30.00 ab	31.10 b	32.03 bc
V ₁ F ₃	28.30 a	30.37 a	32.00 a	33.11 a
V ₂ F ₀	25.02 bc	26.93 f	29.47 de	30.51 ef
V ₂ F ₁	24.18 cd	26.37 f	29.10 e	30.13 f
V ₂ F ₂	25.63 bc	28.97 cd	30.10 cd	31.52 cd
V ₂ F ₃	24.95 bc	28.47 d	30.00 cd	32.03 bc
V ₃ F ₀	22.46 d	25.60 g	27.80 f	29.93 f
V ₃ F ₁	23.91 cd	27.73 e	29.20 e	30.31 f
V ₃ F ₂	24.20 cd	29.03 cd	30.20 cd	31.40 cd
V ₃ F ₃	24.77 bc	29.35 bc	31.10 b	32.52 ab
LSD _{0.05}	1.976	0.6709	0.7497	0.7184
CV (%)	5.62	7.19	6.44	8.24

155

156 Means in a same column followed by different letter (s) are significantly different at $P < 0.05$; (V₁-Atlas
 157 70, V₂-Keifu 65; V₃-Autumn 60 & F₀-Control, F₁-Cow dung, F₂-Poultry manure, F₃-Inorganic fertilizer)

158 3.1.2 Number of leaves/plant



168 **Fig 3: Number of leaves/plant at different growth stages of three different cabbage**
 169 **cultivars**

170 Number of leaves/plant is an important parameter considering the highest performance of
 171 cabbage yield (Figure 3). Autumn – 60 (V₃) gave an idea about highest number of

leaves/plant at all growth stages of cabbage cultivar. The highest number of leaves/plant at 15, 30, 45 DAT and at harvest (17.58, 20.50, 22.00 and 22.75 respectively) was with autumn – 60 which was statistically identical with Keifu – 65 (V_2) at all growth stages. On the contrary the lowest number of leaves/plant at 15, 30, 45 DAT and at harvest (16.50, 17.42, 18.77 and 18.75 respectively) was obtained with Atlas – 70 (V_1). These results might be due to cause of genetical characters of cultivars that caused a higher and lower number of leaves/plant.

Significant variation was observed in the case of number of leaves/plant at different days after transplanting (DAT) (Figure 4). It was measured that the highest number of leaves/plant was obtained with inorganic fertilizer (F_3) and the highest number of leaves/plant was 18.00, 20.44, 21.35 and 22.02 at 15, 30, 45 DAT and at harvest respectively which was statistically identical with treatment of Poultry manure (F_2) at all growth stages of cabbage cultivars. The lowest number of leaves/plant was found to be at 15, 30, 45 DAT and at harvest 15.78, 18.11, 19.01 and 19.67 respectively with control treatment (F_0). The results obtained from the experiment on number of leaves/plant were conformity with Vimala (2006), Pankaj (2006) and Muhammad and Javed (2001) who reported an increased increase in number of leaves per plant with the application of different manures and fertilizers.

The Interaction effect of variety and different types manure and fertilizer affected number of leaves/plant significantly under the present study (Table 2). Different treatment combination viewed different number of leaves/plant according to the treatment at different days after transplanting (DAT). It was observed that highest number of leaves/plant was achieved with V_3F_3 and that was 18.33, 22.33, 24.00 and 25.00 at 15, 30, 45 DAT and at harvest respectively which was closely followed by V_2F_1 , V_2F_2 , V_3F_1 and V_3F_2 at harvest. On the other hand the lowest number of leaves/plant; 15.33, 16.67, 18.03 and 18.00 at 15, 30, 45 DAT and at harvest respectively was obtained with V_1F_0 which was statistically identical with V_1F_1 at all growth stages of cabbage cultivars.




Fig 4: Effect of different fertilizer on number of leaves/plant of different cabbage cultivar at different growth stages

Table 2: Interaction effect of fertilizer management and three different cabbage cultivars on number of leaves/plant at different growth stages

Treatments	Number of leaves/plant			
	15 DAT	30 DAT	45 DAT	At harvest
Interaction effect of variety and fertilizer				
V ₁ F ₀	15.33 d	16.67 f	18.03 e	18.00 d
V ₁ F ₁	15.35 d	15.67 f	17.03 e	18.04 d
V ₁ F ₂	18.33 a	18.67 e	20.00 d	20.00 cd
V ₁ F ₃	17.00 a-d	18.67 e	20.00 d	19.00 d
V ₂ F ₀	15.67 cd	19.33 c-e	21.00 cd	22.00 bc
V ₂ F ₁	17.68 ab	19.00 de	22.00 bc	23.00 ab
V ₂ F ₂	18.00 ab	21.00 b	22.00 bc	23.00 ab
V ₂ F ₃	18.36 a	20.33 b-d	20.04 d	22.00 bc
V ₃ F ₀	16.33 b-d	18.33 e	18.00 e	19.00 d
V ₃ F ₁	18.33 a	20.67 bc	23.00 ab	24.00 ab
V ₃ F ₂	17.33 a-c	20.67 bc	23.00 ab	23.00 ab
V ₃ F ₃	18.69 a	22.33 a	24.00 a	25.00 a
LSD _{0.05}	1.704	1.313	1.607	2.161
CV (%)	5.78	4.24	7.11	8.42

Means in a same column followed by different letter (s) are significantly different at $P < 0.05$; (V₁-Atlas 70, V₂-Keifu 65; V₃-Autumn 60 & F₀-Control, F₁-Cow dung, F₂-Poultry manure, F₃-Inorganic fertilizer)

3.1.3 Stem length

Under the present study, stem length was significantly influenced by different cabbage cultivars (Table 3). Different varieties showed different stem length and it was deliberate at the time of harvest. It was defined that Atlas – 70 (V₁) verified the highest stem length at harvest (4.194 cm) which was closely followed by Keifu – 65 (V₂) and the lowest stem length (3.678 cm) among the cultivars was obtained autumn – 60 (V₃) at harvest. Varietal effect was observed on shoot/stem length due to its phenotypical characters (Haque, 2005) and this result on stem length is supported by Haque, 2005.

Manure and fertilizer effect on stem length was significant under the present study. It is evident that different types of manure and fertilizer showed different stem length (Table 3). The highest stem length (4.203 cm) was indicated with the treatment of Cow dung (F₁) which was statistically identical with control (F₀) treatment and Inorganic fertilizer (F₃) treated plot. On the other hand, the lowest root length was measured with Poultry manure (F₂) treatment. Data of stem length under the present study was in agreement with Souza *et al.* (2008).

Interaction effect of different variety and manures and fertilizer had significant effect on stem length. Different treatment combination showed different root length (Table 3). The highest root length (4.557 cm) was observed with V₁F₁ which was statistically identical with V₂F₁ and statistically similar with V₃F₀. On the other hand, the lowest stem length (3 cm) was observed with V₃F₂ which was closely followed by V₂F₀.

Table 3: Interaction effect of fertilizer management and three different cabbage cultivars on growth parameters; root length and stem length at harvest

Treatments	Root length at harvest (cm)	Stem length at harvest (cm)
Effect of variety		
V ₁	18.03 b	4.194 a
V ₂	22.31 a	3.938 ab
V ₃	22.62 a	3.678 b
LSD _{0.05}	1.319	0.4300
Effect of fertilizer		
F ₀	21.22 a	4.019 a
F ₁	20.26 b	4.203 a
F ₂	21.67 a	3.481 b
F ₃	20.79 ab	4.043 a
LSD _{0.05}	1.167	0.4966
Interaction effect of variety and fertilizer		
V ₁ F ₀	18.45 c	4.333 ab
V ₁ F ₁	16.11 d	4.557 a
V ₁ F ₂	18.89 c	3.777 c-e
V ₁ F ₃	18.67 c	4.110 a-c
V ₂ F ₀	22.94 a	3.333 fg
V ₂ F ₁	22.39 a	4.546 a
V ₂ F ₂	22.89 a	3.667 d-f
V ₂ F ₃	21.00 b	4.193 a-c
V ₃ F ₀	22.27 a	4.390 ab
V ₃ F ₁	22.27 a	3.943 b-d
V ₃ F ₂	23.22 a	3.000 g
V ₃ F ₃	22.71 a	3.380 e-g
LSD _{0.05}	1.105	0.4078
CV (%)	5.33	7.58

Means in a same column followed by different letter-(s) are significantly different at $P < 0.05$; (V₁-Atlas 70, V₂-Keifu 65; V₃-Autumn 60 & F₀-Control, F₁-Cow dung, F₂-Poultry manure, F₃-Inorganic fertilizer)

3.2 Yield Parameters

3.2.1 Thickness of head

Generally thickness of the head of cabbage cultivar control yield and quality of the crop and it is greatly influenced by different varietal characters. The Results showed that there was no significant effect among the three cabbage cultivar considering thickness of head (Table 4). In spite of non-significant variation, the highest (13.44 cm) and lowest (12.76 cm) thickness of the head was achieved by autumn – 60 (V₃) and Keifu – 65 (V₂) respectively. The data obtained on thickness of the head was conformity with Haque (2005).

Thickness of the head was significantly influenced by manure and fertilizer under the present study (Table 4). It is evident that the highest thickness of head (13.78 cm) was obtained with by the treatment of Inorganic fertilizer (F₃) which was significantly different from all other treatment. On the other hand, the lowest thickness of head (12.55 cm) was measured with Control (F₀) treatment which was significantly same with Cow dung (F₁). Souza *et al.* (2008), Bimova (2008) showed the similar results which supported the data on the thickness of the head under the present study.

The interaction effect of different variety and manures and fertilizer had a significant effect on thickness of head. Different treatment combination showed different thickness of head (Table 4). The highest thickness of head (14.44 cm) was observed with V_3F_3 and the lowest thickness of head (11.83 cm) was with V_1F_0 . The results obtained from all other treatments were significantly different from highest and the lowest thickness of the head.

3.2.2 Diameter of head

Diameter of head is a measurement of the size of actual cabbage shape which indicates the yield amount and/or market value. Significant variation was observed in case of diameter of head among the cabbage cultivar (Table 4). Result revealed that the highest diameter of head (20.24 cm) was achieved with by Atlas – 70 (V_1) where the lowest (18.03 cm) was with Autumn – 60 (V_3). Similar results were obtained by Haque (2005) and Muhammad and Javed (2001) with their experiments.

Table 4: Interaction Effect of fertilizer management and three different cabbage cultivars on yield contributing parameters Thickness of head and Diameter of head at harvest

Treatments	Thickness of head (cm) at harvest	Diameter of head (cm) at harvest
Effect of variety		
V_1	13.03	20.24 a
V_2	12.76	18.85 b
V_3	13.44	18.03 c
LSD _{0.05}	NS	0.8160
Effect of fertilizer		
F_0	12.55 c	18.06 d
F_1	12.63 c	18.46 c
F_2	13.36 b	19.37 b
F_3	13.78 a	20.25 a
LSD _{0.05}	0.2375	0.3511
Interaction effect of variety and fertilizer		
V_1F_0	11.83 f	19.45 bc
V_1F_1	12.48 e	19.17 c
V_1F_2	13.48 bc	20.89 a
V_1F_3	13.89 b	21.44 a
V_2F_0	13.09 cd	18.18 d
V_2F_1	12.28 e	18.11 d
V_2F_2	13.11 cd	19.11 c
V_2F_3	13.01 d	19.99 b
V_3F_0	12.52 e	16.56 e
V_3F_1	13.33 cd	18.11 d
V_3F_2	13.48 bc	18.11 d
V_3F_3	14.44 a	19.33 c
LSD _{0.05}	0.4113	0.6082
CV (%)	4.48	5.66

Means in a same column followed by different letter (s) are significantly different at $P < 0.05$; (V_1 -Atlas 70, V_2 -Keifu 65; V_3 -Autumn 60 & F_0 -Control, F_1 -Cow dung, F_2 -Poultry manure, F_3 -Inorganic fertilizer)

270 Diameter of head was significantly influenced by manure and fertilizer under the present
271 study (Table 4). It is evident that the highest diameter of head (20.25 cm) was obtained with
272 the treatment of Inorganic fertilizer (F_3) which was significantly different from all other
273 treatment. On the other hand, the lowest diameter of head (18.06 cm) was measured with
274 Control (F_0) treatment. Data measurement on head diameter was in agreement with Souza
275 *et al.* (2008) and Vimala (2006).

276 Interaction effect of different variety and manures and fertilizer had significant effect on
277 diameter of head. Different treatment combination showed different diameter of head (Table
278 4). The highest diameter of head (21.44 cm) was observed with V_1F_3 which was statistically
279 same with V_1F_2 and the lowest diameter of head (16.56 cm). The results obtained from all
280 other treatment were significantly different from highest and lowest diameter of head.

281

282 3.2.3 Weight of whole plant

283 Weight of whole plant was significantly influenced by different cabbage cultivar (Table 5).
284 Results showed that the highest whole plant weight (2.23 kg/plant) was with Atlas – 70 (V_1)
285 where the lowest (1.96 kg/plant) was with Autumn – 60 (V_3). Supported results with the
286 present study were achieved by Haque (2005), Muhammad and Javed (2001) who reported
287 the higher weight of plant found in Atlas- 70.

288 Weight of whole plant was significantly influenced by manure and fertilizer under the present
289 study (Table 5). It is evident that the highest whole plant weight (2.41 kg/plant) was obtained
290 with the treatment of Inorganic fertilizer (F_3) which was significantly same with Poultry
291 manure (F_2) treated plot. On the other hand, the lowest whole plant weight (1.80 kg/plant)
292 was measured with Control (F_0) treatment which was significantly same with Cow dung (F_1)
293 treated plot that is suggested by Pankaj (2006) and Hsieh (2004).

294 | Interaction effect of different variety and manures and fertilizer had a significant effect on
295 whole plant weight. Different treatment combination showed different whole plant weight
296 (Table 5). The highest whole plant weight (2.56 kg/plant) was observed with the treatment
297 combination of V_1F_2 . The lowest whole plant weight (1.57 kg/plant) was obtained with V_3F_0
298 which was statistically identical with V_3F_1 . The results obtained from all other treatment were
299 significantly different from highest and lowest whole plant weight.

300 3.2.4 Marketable yield

301 Marketable yield was significantly affected by different variety used in the experiment (Table
302 5). The highest marketable yield (27.42 kg/plot and 45.29 t/ha) was obtained from Atlas – 70
303 (V_1) but the variety, Autumn – 60 (V_3) viewed lowest marketable yield (21.57 kg/plot and
304 35.95 t/ha) which was statistically different from others varieties. The results achieved by
305 Haque (2005), Muhammad and Javed (2001) was similar to the present study as they found
306 higher marketable yield with the similar variety.

307

308

309

Table 5: Interaction effect of fertilizer management and three different cabbage cultivars on yield parameters; weight of whole plant at harvest, gross yield, marketable yield and economic production (kg/plant) at harvest

Treatments	Weight of whole plant at harvest (kg/plant)	Gross yield (kg/plot)	Marketable yield (kg/plot)	Gross yield (t/ha)	Marketable yield (t/ha)	Economic production (kg/plant) at harvest
Effect of variety						
V ₁	2.23 a	28.00 a	27.42 a	46.67 a	45.29 a	1.576 a
V ₂	2.10 b	23.57 b	22.51 b	39.27 b	37.51 b	1.311 b
V ₃	1.96 c	22.75 b	21.57 c	37.91 c	35.95 c	1.226 b
LSD _{0.05}	0.093	0.9466	0.8169	1.100	1.199	0.2159
Effect of fertilizer						
F ₀	1.80 b	15.65 c	14.34 c	26.08 c	23.90 c	1.178 b
F ₁	1.86 b	24.38 b	23.32 b	40.64 b	38.87 b	1.198 b
F ₂	2.30 a	29.61 a	28.92 a	49.34 a	47.92 a	1.523 a
F ₃	2.41 a	29.45 a	28.75 a	49.09 a	47.64 a	1.584 a
LSD _{0.05}	0.339	1.093	0.9433	1.270	1.384	0.2492
Interaction effect of variety and fertilizer						
V ₁ F ₀	2.09 de	16.57 e	15.31 g	27.61 f	25.52 f	1.450 cd
V ₁ F ₁	1.88 f	26.93 c	26.17 d	44.89 c	43.62 c	1.257 de
V ₁ F ₂	2.56 a	37.28 a	36.91 a	62.14 a	61.52 a	1.853 a
V ₁ F ₃	2.39 a-c	31.24 b	31.31 b	52.06 b	50.51 b	1.743 ab
V ₂ F ₀	1.87 f	15.77 e	14.43 gh	26.28 fg	24.05 fg	1.107 ef
V ₂ F ₁	1.97 ef	23.86 d	22.79 e	39.76 d	37.98 d	1.250 de
V ₂ F ₂	2.23 cd	27.53 c	26.50 d	45.88 c	44.17 c	1.463 cd
V ₂ F ₃	2.34 bc	27.10 c	26.30 d	45.16 c	43.84 c	1.423 cd
V ₃ F ₀	1.57 g	14.60 e	13.28 h	24.34 g	22.13 g	0.977 f
V ₃ F ₁	1.63 g	22.36 d	21.01 f	37.26 e	35.02 e	1.087 ef
V ₃ F ₂	2.11 de	24.01 d	22.85 e	40.01 d	38.08 d	1.253 de
V ₃ F ₃	2.51 ab	30.02 b	29.15 c	50.04 b	48.58 b	1.587 bc
LSD _{0.05}	0.186	1.893	1.634	2.199	2.398	0.2074
CV (%)	6.14	4.51	5.83	7.18	4.62	7.46

Means in a same column followed by different letter (s) are significantly different at $P < 0.05$; (V₁-Atlas 70, V₂-Keifu 65; V₃-Autumn 60 & F₀-Control, F₁-Cow dung, F₂-Poultry manure, F₃-Inorganic fertilizer)

Marketable yield was significantly influenced by application of different manure and fertilizer according to the treatment under the present study (Table 5). It is evident that the highest marketable yield (28.92 kg/plot and 47.92 t/ha) was obtained with the treatment of poultry manure (F₂) which was statistically identical with inorganic fertilizer treated plot (F₃). On the other hand, the lowest marketable yield (14.34 kg/plot 23.90 t/ha) was measured with Control (F₀) treatment. The results obtained from the experiment were conformity with Hsieh (2004) and Chan *et al.* (2008) who reported better marketable yield with the application of different manures and fertilizers.

Interaction effect of different variety and manures and fertilizer had significant effect on marketable yield of cabbage cultivars. Different treatment combination showed different yield (Table 5). The highest marketable yield (36.91 kg/plot and 61.52 t/ha) was observed with the treatment combination of V₁F₂. The treatment combination, V₁F₃ and V₃F₃ also showed higher yield but significantly lower than V₁F₂. The lowest marketable yield (13.28 kg/plot and

328 22.13 t/ha) was obtained with V_3F_0 which was closely related to V_2F_0 . The results obtained
329 from all other treatments were significantly different from highest and lowest yield. Similar
330 findings are observed with Yau (2006).

331 4. CONCLUSION

332
333 Judicial application of organic and inorganic fertilizers can minimize the application inorganic
334 fertilizer to reduce the hazardous effect on public health and environment. The result showed
335 that V_1F_2 (Atlas – 70 × Poultry manure) performed best in producing higher yield than other
336 treatments comprised with other variety and fertilizer application under the present study. On
337 the other hand interactions of variety (Atlas – 70) and organic fertilizer (Poultry manure)
338 showed its superiority in producing higher cabbage yield and economic production.
339 Therefore, it may be concluded that Atlas – 70 can be used along with poultry manure can
340 be used for higher yield and economic production of cabbage.

341 REFERENCES

- 342
343 Altieri, M. and Nicholls, F. N. Tolerance and effect of leaf fertilization treatments on
344 vegetables. *Gartebauwissenschaft*. 2003;51 (2): 58-62.
- 345 Bimova Impact of organic fertilizers on total antioxidant capacity in head cabbage. Faculty of
346 Horticulture, Mendel University of Agriculture and Forestry in Brno, Lednice, Czech Republic.
347 2000.
- 348 Edris, K. M., Islam, A. T. M. T., Chowdhury, M. S. and Haque, A. K. M. Detailed Soil Survey
349 of Bangladesh Agricultural University Farm, Mymensingh. Dept. Soil Survey, Govt. People's
350 Republic of Bangladesh. 1979;118 p.
- 351 FAO. Food and Agricultural Organization of the United Nations, Soil Survey Project of
352 Bangladesh. Soil Resources Tech. Rep. 1988; pp. 101-159.
- 353 FAO. Production Year Book. Food and Agricultural of the United Nations. Rome, Italy. 1988;
354 42: 190-193.
- 355 Gomez, K. A. and A. A. Gomez. Statistical Procedure for Agricultural Research (2nd ed.), A
356 Wiley Inter Science Publication, John Wiley and Sons, New York. 1984; 680p.
- 357 Haque, K.M.F. Influence of variety and time of planting on the yield performance and nutrient
358 contents of cabbage (*Brassica oleracea* L.). Dhaka, Bangladesh: Bangladesh Council of
359 Scientific and Industrial Research. Bangladesh-Journal-of-Scientific-and-Industrial-
360 Research. 2005; 40(3/4): 211-218.
- 361 Hsieh, C. F., H. C. Fang, K. Nan and K. N. Hsu. Effect of continuous use of organic manures
362 on the growth and yield of vegetable soybean' and cabbage. Bulletin of Taichung District.
363 Agric. Improvement Sta., Japan, 1995; 46: 1-10.
- 364 Huang, F. Varietal influence in organic farming on autumn cabbage. Hangzhou, China:
365 Zhejiang Academy of Agricultural Sciences. Acta-Agriculturae-Zhejiangensis. 2006; 18(1):
366 58-61.

- 367 Letourneau, A. F. The characteristics of mineral nutrition and yield formation in cabbage
368 cultivar in relation to fertilizers and mineral uptake by cabbage plants. *Agrokhimiya*, 1996; 9:
369 61-67.
- 370 Muhammad, I. and Javed, U. Response of Chinese cabbage cultivars to different levels of
371 nitrogen with constant doses of phosphorus and potassium under the agro-climatic
372 conditions of Chilas (Diamer). *Sarhad-Journal-of-Agriculture*. 2001; 17(1): 81-85.
- 373 Pankaj, S. Integrated effect of bio-inoculants, organic and inorganic fertilizer on growth and
374 yield of cabbage. Hisar, India: Agricultural Research Information Centre. *Crop-Research-*
375 *Hisar*. 2006; 32(2): 188-191.
- 376 Souza, P. A., Souza, G. L. F. M., Menezes, J. B. and Bezerra, N. F. Evaluations of cabbage
377 cultivar grown under organic compost and mixed mineral fertilizers. *Horticultura Brasileira*.
378 2008; 26(1): 143-145.
- 379 Tindall, M. Mineral and organic fertilizing in cabbage. Residual effect for commercial
380 cultivation on yield and quality performance with organic farming. *Hort. Bras.*, 2000; 6(1) :
381 15-20.
- 382 UNDP. Land Resource Appraisal of Bangladesh for Agricultural Development Report 2: Agro-
383 ecological Regions of Bangladesh, FAO, Rome, Italy, 1988. p. 577.
- 384 P. Vimala, M.K. Illias, H. Salbiah. Effect of rates of organic fertilizer on growth, yield and
385 nutrient content of cabbage (*Brassica oleracea* var. *capitata*) grown under shelter. *Acta-*
386 *Horticulturae*. 2006; (710): 391-397. <https://doi.org/10.17660/ActaHortic.2006.710.47>
- 387 P.Y. Yau, S. Ahmad, M.K. Illias, K. Ganisan. Evaluation of cabbage cultivars (*Brassica*
388 *oleracea* var. *capitata*) under plastic rain-shelter on mineral soils in the lowlands. *Acta Hortic*,
389 2006; 710: 343-346. <https://doi.org/10.17660/ActaHortic.2006.710.39>