

# Original Research Article

## Effect of Different Levels of Nitrogen, Phosphorus and Potassium on Insect and Pests in T. Aman Rice (*Oryza sativa* L.)

### ABSTRACT

The study was conducted in the experimental area of Sher-e-Bangla Agricultural University (SAU), Sher-e-Bangla Nagar, Dhaka-1207, Bangladesh during the period from July to October 2016 to find out the effect of different levels of NPK on insect pests in transplant aman rice. BRRI dhan33 were used as the test crop in this experiment. The experiment comprised the different NPK (Nitrogen, Phosphorus and Potassium) fertilizers dose as treatment where,  $T_0$  = Untreated control, no NPK fertilizers,  $T_1$  = NPK @ 45, 50, 40 kg/ha,  $T_2$  = NPK @ 70, 25, 40 kg/ha,  $T_3$  = NPK @ 70, 50, 20 kg/ha,  $T_4$  = NPK @ 70, 50, 40 kg/ha,  $T_5$  = NPK @ 70, 50, 60 kg/ha,  $T_6$  = NPK @ 70, 75, 40 kg/ha and  $T_7$  = NPK @ 95, 50, 40 kg/ha. The experiment was laid out in a Randomized complete block design (RCBD) with three replications. Data were recorded on different types of insects and pests that were identified for the entire growing period with their number and incidence on rice plants. During the entire growing period 5 selected hills/plot were monitored with clean observation and yellow stem borer, leaf folder, rice hispa, grasshopper, brown plant hopper, green leaf hopper and rice bug insects and pests were observed. The lowest number of infestation of insects and pest was observed from  $T_5$ , whereas the highest number was found from  $T_7$  treatment. In case of incidence percent of dead heart, data was observed at 25, 45 and 65 days after transplanting (DAT), respectively. Data recorded from each plot revealed that the lowest incidence of dead heart was observed from  $T_5$  (3.64%, 4.23% and 4.47%, respectively), while the highest incidence of dead heart was found from  $T_7$  (10.37%, 13.56% and 14.73%, respectively) treatment. In terms of white head incidence, at 60, 70 and 80 DAT, data recorded from each plot revealed that the lowest incidence of white head was found from  $T_5$  (2.35%, 2.66% and 3.12%, respectively), whereas the highest incidence of white head was observed from  $T_7$  (7.57%, 8.26% and 8.64%, respectively).

**Keywords:** *Oryza sativa*, nitrogen, phosphorus, potassium, insects, pests

### 1. INTRODUCTION

Rice (*Oryza sativa*) is the most important food crop around the world and the staple food for approximately more than two billion people in the Asia (Hien *et al.*, 2006). Rice is more nutritious than any other cereal crops and is an ideal host for over 800 species of insect (Barr and Smith, 1975). In tropical Asia, more than 100 species of insects are persistent to rice. In Bangladesh, about 175 insect pest species have been reported, which cause damage to the rice plants (Mustafi *et al.*, 2007), of these 20-30 species are economically important (Miah and Karim, 1984). The estimated loss of rice in Bangladesh due to insect pests and diseases amounts to 1.5 to 2.0 million tons (Siddique, 1992). At high population density, crop loss may be 100% (Rahman *et al.*, 2004). Major pests cause damage about 28% to Aman crops and the estimated annual loss of rice in Bangladesh due to insect pests and diseases amounts 1.5 to 2.0 million tons (BRRI, 1989). Nutrient management is the most important practice in rice production system, but it may affects response of rice to insect pests. If there is positive interaction between nutrients and pest can be identified, can provide guidelines for optimizing total agro-ecosystem function (Magdoff *et al.* 2000). Some aman rice pests are green leaf hopper, rice hispa, green stink bug, rice leafroller, yellow stem borer and rice bug. The beneficial insects are categories as predator and parasites, collectively known as natural enemies which are able to interact with their prey and consequently regulate them at reasonably lower level. 99 species of parasites and 88 species of predator of rice insect pests have been recorded in Bangladesh (Wahiduzzaman, 1993). The application of nitrogen fertilizer in plants can normally increase herbivore's feeding preference, food consumption, survival, growth, reproduction, and population density, except few examples that nitrogen fertilizer reduces the herbivore performances. Higher nitrogen doses cause for higher sucking pests incidence. Phosphorus has not been considered as important or limiting as nitrogen for phytophagous insects. However, phosphorus is determinant of growth rate and population density. Phosphorus is an important component for the population growth of phytophagous insects as it is required for RNA synthesis. Potassium induced change in rice plant

at profound effect on insect–host interactions. Increase of K in rice plant causes reduction in the feeding rate of BPH *Nilaparvata lugens* (Vaithilingan et al., 1975) and *Nephotettix* sp. (Subramanian and Balasubramnian, 1976). To make the green revolution successful & to mitigate the adverse effects of fertilizers to the crop productive environment. Judicious use of fertilizers is considered as one of the important aspect of cultural practices in IPM which influence the activity of insect pests and ultimate effect on growth, development and yield of crop plant. Keeping in view, the present study was undertaken to assess the effect of different levels of fertilizers on incidence of arthropod fauna on rice.

Present study was aimed to identify optimum doses of NPK for lower incidence of insects and pests and higher grain yields and better quality of rice, to ensure the minimum use of chemical nutrients and find out the proper dose of nutrient components (NPK) against these insects and pests

## 2. MATERIALS AND METHODS

### 2.1 Experimental site

The experiment was conducted during the period from July 2016 to October 2016.

The present piece of research work was conducted in the experimental area of Sher-e-Bangla Agricultural University (SAU), Sher-e-Bangla Nagar, Dhaka-1207, Bangladesh. The geographical location of the experimental site was under the subtropical climate and its climatic conditions is characterized by heavy rainfall during the month of April to September and scanty rainfall during the rest period of the year.

### 2.2 Experimental material:

BRRRI dhan33, high yielding rice variety developed by Bangladesh Rice Research Institute (BRRRI), Gazipur, Bangladesh was used as experimental material

### 2.3 Treatments:

The experiment comprised of the following NPK dose as treatment.

T<sub>0</sub> = Untreated control, no NPK fertilizers,

T<sub>1</sub> = NPK @ 45, 50, 40 kg/ha

T<sub>2</sub> = NPK @ 70, 25, 40 kg/ha

T<sub>3</sub> = NPK @ 70, 50, 20 kg/ha

T<sub>4</sub> = NPK @ 70, 50, 40 kg/ha

T<sub>5</sub> = NPK @ 70, 50, 60 kg/ha

T<sub>6</sub> = NPK @ 70, 75, 40 kg/ha

T<sub>7</sub> = NPK @ 95, 50, 40 kg/ha

In each of the treatment Nitrogenous fertilizer was applied in three splits at equal amount where phosphorus and potash fertilizer were applied at single dose as basal.

### 2.4 Experimental design and layout:

The experiment was laid out in a randomized complete block design (RCBD) with three replications, where the experimental area was divided into three blocks representing the replications to reduce soil hetero-genetic effects. Each block was divided into 8 unit plots as treatments demarked with raised bunds. Thus, the total numbers of plots were 24. The unit plot size was 4.0 m × 2.5 m. The distance maintained between two blocks and two plots were 0.5 m and 0.5 m, respectively.

### 2.5 Intercultural Operations:

Fertilizers other than NPK such as zinc, boron, Sulphur were applied as per recommended for BRRRI dhan33 by Bangladesh Rice Research Institute. Other intercultural operations such as raising of seedlings, land preparation, manuring, irrigation and drainage, weeding were done as per necessity. Assessing Infestation level

### 2.6 Observations:

Five hills were selected at random per replicate for each treatment. The dead heart tiller and white head infested tiller were counted. In case of dead heart, it was counted in vegetative growth stage and white head infested tillers was counted at reproductive stage converted into per plant. Hopper burn was counted in tillering, panicle initiation, before ripening and after ripening stage. The observation was recorded at the first observation of symptom and was continued up to maturity at 7 days interval.

Equation for percent Dead Heart, White Head and Hopper Burn infestation

Number of Dead Heart/white head/hopper burn infested tiller per five hills were counted and divided by total number of tillers in five hills and then multiplied by 100 to assess the percentage of infestation. For Example,

No. of dead heart infested tiller

% dead heart/ infested tillers = ----- x 100

Total no. of tiller per five hills

## 2.7 Collection of Data and Statistical analysis

The data obtained for different characters were statistically analyzed to observe the significant difference among the treatments. The mean values of all the characters were calculated and analysis of variance was performed by using MSTAT-C software. The significance of the difference among the treatments means was estimated by the Least Significant Difference (LSD) test at 5% level of probability (Gomez and Gomez, 1984).

## 3. RESULTS AND DISCUSSION

### 3.1 Number of different insect population

During the entire growing period 5 selected hills/plot were monitored with clean observation and yellow stem borer, leaf folder, rice hispa, brown plant hopper, green leaf hopper and rice bug insect pests was observed in different growth stage as insect populations of rice plants. The number of these identified insects were recorded and presented in Table 1. For different levels of NPK fertilizers number of different insect pests showed statistically significant differences under the present trial.

### 3.2 Yellow stem borer

The data presented in table 1 revealed that the number of yellow stem borer varied from 1.67 to 9.33 for different levels of NPK fertilizers. The lowest number (1.67) of yellow stem borer was observed from T5 (NPK @ 70, 50, 60 kg/ha) treatment whereas the highest number (9.33) was found from T7 (NPK @ 95, 50, 40 kg/ha) treatment. Optimum doses of NPK fertilizers was more effective in controlling of yellow stem borer, whereas in excessive application of NPK fertilizers increased the incidence of yellow stem borer. Rangini *et al.* (2005) reported that yellow stem borer (YSB) infestation was extensively occurred early tillering to maximum tillering stage.

### 3.3 Leaf folder

In the present study, the number of leaf folder varied from 1.13 to 12.67 upon described treatments. Data revealed that the lowest number (1.13) of leaf folder was recorded from T5 (NPK @ 70, 50, 60 kg/ha) treatment, while the highest number (12.67) was observed from T7 (NPK @ 95, 50, 40 kg/ha) treatment. de Kraker *et al* found that the average density of leaf folder larvae at the highest nitrogen level was eight times more than that at the zero-nitrogen level.

### 3.4 Rice hispa

The range of rice hispa was 1.13 to 3.27 for different levels of NPK fertilizers under the study. The lowest number (1.13) of rice hispa was recorded from T5 treatment (NPK @ 70, 50, 60 kg/ha) while the highest number (3.67) was observed from T7 treatment (NPK @ 95, 50, 40 kg/ha). Singh *et al.* (1990) in Punjab indicated that the NPK at 120:60:60 kg/ha increased the susceptibility of rice to infestation by rice hispa.

### 3.6 Brown plant hopper

The lowest number (1.93) of brown plant hopper was found from T5 (NPK @ 70, 50, 60 kg/ha) treatment, whereas the highest number (9.47) was recorded from T7 treatment (NPK @ 95, 50, 40 kg/ha). Madhuri (2016) reported that the lowest BPH population (2.87/hill) was found in control treatments which was devoid of all types of nutrients. Whereas, without micronutrients other treatments with only nitrogen produced higher incidence or in combination with boron, zinc etc caused lower incidence.

### 3.7 Green leaf hopper

In consideration of green leaf hopper, data revealed that the number of green leaf hopper in 5 selected hills vary from 2.47 to 11.67 for different levels of NPK fertilizers. The lowest number (2.47) of green leaf hopper was recorded from T5 treatment (NPK @ 70, 50, 60 kg/ha), while the highest number was observed from T7 (11.67) treatment (NPK @ 95, 50, 40 kg/ha). Nath and Bhagabati (2005) reported that the green leaf hopper population was first appeared in the rice seedbed during June-July, reaching the peak in October -November in the main rice field and disappeared from field from December to May.

### 3.8 Rice Bug

In case of rice bug, the numbers of rice bug differ from 1.73 to 8.60 per 5 selected hills due to different levels of NPK fertilizers. The lowest number (1.73) of rice bug was observed from T5 treatment (NPK @ 70, 50, 60 kg/ha), whereas the highest number (8.60) was found from T7 treatment (NPK @ 95, 50, 40 kg/ha). Tsueda *et al.* (2002) studied on the occurrence of rice bugs, a total of 22 species, in rice fields. They also observed that *Stenotus rubro vittatus* was the important species and the peak occurrence of it coincided with the date of heading of early-ripening rice.

178 **Table 1. Number of major insect pests in BRRI dhan33 during the growing period for different**  
 179 **levels of NPK fertilizers**

Treatment	Number of different insect pests in 5 selected hills						
	Yellow stem borer	Leaf folder	Rice hispa	Grasshopper	Brown plant hopper	Green leaf hopper	Rice bug
T0	7.67 b	10.27 b	3.27 b	11.13 b	7.27 b	9.27 b	6.33 b
T1	3.13 g	2.73 f	1.60 f	4.27 f	2.47 g	2.87 fg	2.60 f
T2	6.87 c	8.13 c	2.87 c	8.33 c	5.87 c	7.87 c	5.53 c
T3	6.33 d	7.33 d	2.47 d	6.33 d	5.33 d	6.13 d	4.80 d
T4	5.80 e	5.33 e	2.13 e	5.47 de	4.87 e	4.87 e	4.13 e
T5	1.67 h	1.13 g	1.13 g	3.33 g	1.93 h	2.47 g	1.73 g
T6	5.27 f	4.93 e	1.67 f	4.93 ef	4.27 f	3.07 f	3.80 e
T7	9.33 a	12.67 a	3.67 a	12.73 a	9.47 a	11.67 a	8.60 a
LSD(0.05)	0.470	0.751	0.293	0.899	0.467	0.522	0.495
CV(%)	4.65	6.52	7.07	7.26	5.13	4.95	6.04

180

T0 = Untreated control, no NPK fertilizers

T1 = NPK @ 45, 50, 40 kg/ha

T2 = NPK @ 70, 25, 40 kg/ha

T3 = NPK @ 70, 50, 20 kg/ha

T4 = NPK @ 70, 50, 40 kg/ha

T5 = NPK @ 70, 50, 60 kg/ha

T6 = NPK @ 70, 75, 40 kg/ha

T7 = NPK @ 95, 50, 40 kg/ha

181

182

### 183 3.9 Infestation of rice by different insects and pests in different stages

184 Dead heart incidence by yellow stem borer at 25, 45 and 60 Days after transplanting

185 In case of incidence of dead heart, at 25 DAT, data recorded from each plot revealed that the lowest  
 186 incidence of dead heart was observed from T5 (3.64%), while the highest incidence of dead heart was  
 187 found from T7 (10.37%) treatment. At 45 DAT, from each plot it was revealed that the lowest  
 188 incidence of dead heart was observed from T5 (4.23%), while the highest incidence of dead heart  
 189 from T7 (13.56%) treatment. At 65 DAT, data recorded from each plot revealed that the lowest  
 190 incidence of dead heart was observed from T5 (4.47%), while the highest incidence of dead heart was  
 191 found from T7 (14.73%) treatment. In case of incidence of dead heart decrease/increase over control,  
 192 the highest decrease was observed in T5 at 25, 45 and 60 DAT as -61.36, -59.17 and -59.58  
 193 respectively. On the other hand, the increase of dead heart incidence was found from T7 at 25, 45 and  
 194 60 DAT as +10.08, +30.89 and +33.18 respectively. Ramzan *et al.* (1992) reported that high  
 195 infestation of yellow stem borer is correlated with the high use of nitrogenous fertilizers in rice field.

196

197 **Table 2. Incidence of rice yellow stem borer (dead heart) infestation in BRRI dhan33 for**  
 198 **different levels of NPK fertilizers**

Treatment s	Incidence of rice yellow stem borer infestation (dead heart/plot) at different days after transplanting (DAT)					
	1 <sup>st</sup> observation (25 DAT)		2 <sup>nd</sup> observation (45 DAT)		3 <sup>rd</sup> observation (65 DAT)	
	Dead heart (%)	Decrease/increase over control (%)	Dead heart (%)	Decrease/increase over control (%)	Dead heart (%)	Decrease/increase over control (%)
T0	9.42 b	--	10.36 b	--	11.06 b	--
T1	4.24 e	-54.99	5.16 e	-50.19	5.66 d	-48.82

T2	7.27 c	-22.82	8.47 c	-18.24	9.08 c	-17.90
T3	6.08 d	-35.46	6.24 d	-39.77	6.57 d	-40.60
T4	5.77 d	-38.75	5.96 de	-42.47	6.41 d	-42.04
T5	3.64 f	-61.36	4.23 f	-59.17	4.47 e	-59.58
T6	4.66 e	-50.53	5.48 de	-47.10	6.10 d	-44.85
T7	10.37 a	+10.08	13.56 a	+30.89	14.73 a	+33.18
LSD(0.05)	0.517	--	0.810	--	1.124	--
CV(%)	4.59	--	6.22	--	8.01	--

199

200 In a column, numeric data represents the mean value of 3 replications; each replication is derived  
201 from each plot of each treatment

202 In a column means having similar letter(s) are statistically identical and those having dissimilar  
203 letter(s) differ significantly as per 0.05 level of probability

204

T0= Untreated control, no NPK fertilizers

T2= NPK @ 70, 25, 40 kg/ha

T4 = NPK @ 70, 50, 40 kg/ha

T6 = NPK @ 70, 75, 40 kg/ha

T1 = NPK @ 45, 50, 40 kg/ha

T3 = NPK @ 70, 50, 20 kg/ha

T5 = NPK @ 70, 50, 60 kg/ha

T7 = NPK @ 95, 50, 40 kg/ha

205

206

207

208

209

### 210 3.10 White head incidence by yellow stem borer at 60, 70 and 80 DAT

211 In terms of white head incidence, at 60 DAT, 70 DAT and 80 DAT, data recorded from each plot  
212 revealed that the lowest incidence of white head was found from T5 (2.35%, 2.66% and 3.12%  
213 respectively), while the highest incidence of white head was observed from T7 (7.57%, 8.26%  
214 and 8.64% respectively) treatment. In case of incidence of white head decrease/increase over control,  
215 the highest decrease was observed in T5 at 60, 70 and 80 DAT as -54.19, -54.45 and -47.56  
216 respectively. Whereas, the increase of white head incidence was found from T7 treatment at 60, 70  
217 and 80 DAT as +47.56, +41.44 and +45.21 respectively. Chakraborty (2011) observed that incidence  
218 of white head (WH) was 206.72% higher than the control field when the field was fertilized by 140 kg  
219 N/ha.

220 **Table 4. Incidence of rice yellow stem borer (white head) infestation in BRRI dhan33 for**  
221 **different levels of NPK fertilizers**

Treatment s	Incidence of rice yellow stem borer infestation (white head/plot) at different days after transplanting (DAT)					
	1 <sup>st</sup> observation (60 DAT)		2 <sup>nd</sup> observation (70 DAT)		3 <sup>rd</sup> observation (80 DAT)	
	White head (%)	Reduction over control (%)	White head (%)	Reduction over control (%)	White head (%)	Reduction over control (%)
T0	5.13 b	--	5.84 b	--	5.95 b	--
T1	3.08 d	-39.96	2.97 e	-49.14	3.45 de	-42.02
T2	4.69 b	-8.58	5.16 b	-11.64	5.67 b	-4.71

T3	3.96 c	-22.81	4.06 c	-30.48	4.38 c	-26.39
T4	3.66 c	-28.65	3.83 cd	-34.42	4.08 c	-31.43
T5	2.35 e	-54.19	2.66 e	-54.45	3.12 e	-47.56
T6	3.19 d	-37.82	3.24 de	-44.52	3.99 cd	-32.94
T7	7.57 a	+47.56	8.26 a	+41.44	8.64 a	+45.21
LSD(0.05)	0.436	--	0.731	--	0.559	--
CV(%)	5.92	--	9.25	--	6.49	--

222

223 In a column, numeric data represents the mean value of 3 replications; each replication is derived  
224 from each plot of each treatment

225 In a column means having similar letter(s) are statistically identical and those having dissimilar  
226 letter(s) differ significantly as per 0.05 level of probability

227

T0= Untreated control, no NPK fertilizers

T2= NPK @ 70, 25, 40 kg/ha

T4 = NPK @ 70, 50, 40 kg/ha

T6 = NPK @ 70, 75, 40 kg/ha

T1 = NPK @ 45, 50, 40 kg/ha

T3 = NPK @ 70, 50, 20 kg/ha

T5 = NPK @ 70, 50, 60 kg/ha

T7 = NPK @ 95, 50, 40 kg/ha

228

229

230

### 231 3.11 Leaf folder incidence in leaf at 30, 45 and 60 DAT

232 In consideration of leaf folder incidence, at 30 DAT, 45 DAT and 60 DAT data recorded from each plot  
233 revealed that the lowest incidence of leaf folder was found from T5 (3.47%, 3.94% and 4.04%  
234 respectively), while the highest incidence of leaf folder was observed from T7 (8.05%, 8.56% and  
235 9.18% respectively) treatment. In case of incidence of leaf folder decrease/increase over control, the  
236 highest decrease was observed in T5 at 30, 45 and 60 DAT as -43.30, -40.66 and -41.79 respectively.  
237 Whereas, the highest increase of leaf folder incidence was found from T7 treatment at 30, 45 and 60  
238 DAT as +31.54, +28.92 and +32.28 respectively. Mahadev *et al.* (1995) reported that the crop applied  
239 with N either alone or coupled with P exhibited a lowered the incidence of leaf folder in rice field at  
240 vegetative stage.

241 **Table 5. Incidence of leaf folder infestation in BRRI dhan33 for different levels of NPK**  
242 **fertilizers**

Treatment s	Incidence of leaf folder at different days after transplanting (DAT)					
	1 <sup>st</sup> observation (30 DAT)		2 <sup>nd</sup> observation (45 DAT)		3 <sup>rd</sup> observation (60 DAT)	
	Leaf infestation (%)	Reduction over control (%)	Leaf infestation (%)	Reduction over control (%)	Leaf infestation (%)	Reduction over control (%)
T0	6.12 b	--	6.64 b	--	6.94 b	--
T1	3.84 ef	-37.25	4.56 e	-31.33	4.72 e	-31.99
T2	5.47 c	-10.62	5.93 c	-10.69	6.12 c	-11.82
T3	4.92 d	-19.61	5.34 d	-19.58	5.64 cd	-18.73



T4	4.57 d	-25.33	4.91 de	-26.05	5.23 de	-24.64
T5	3.47 f	-43.30	3.94 f	-40.66	4.04 f	-41.79
T6	4.05 e	-33.82	4.67 e	-29.67	4.91 e	-29.25
T7	8.05 a	+31.54	8.56 a	+28.92	9.18 a	+32.28
LSD(0.05)	0.403	--	0.502	--	0.643	--
CV(%)	4.57	--	5.15	--	6.29	--

243

244 In a column, numeric data represents the mean value of 3 replications; each replication is derived  
245 from each plot of each treatment

246 In a column means having similar letter(s) are statistically identical and those having dissimilar  
247 letter(s) differ significantly as per 0.05 level of probability

248

T0= Untreated control, no NPK fertilizers

T2= NPK @ 70, 25, 40 kg/ha

T4 = NPK @ 70, 50, 40 kg/ha

T6 = NPK @ 70, 75, 40 kg/ha

T1 = NPK @ 45, 50, 40 kg/ha

T3 = NPK @ 70, 50, 20 kg/ha

T5 = NPK @ 70, 50, 60 kg/ha

T7 = NPK @ 95, 50, 40 kg/ha

249

250

### 251 3.12 Brown plant hopper incidence at 35, 50 and 65 DAT

252 In consideration of brown plant hopper incidence, at 35 DAT, 50 DAT and 65 DAT data recorded from  
253 each plot revealed that the lowest incidence of brown plant hopper was found from T5 (4.23%, 4.55%  
254 and 4.78% respectively), while the highest incidence of brown plant hopper was observed from T7  
255 (7.88%, 7.12% and 7.95% respectively) treatment. In case of incidence of BPH decrease/increase  
256 over control, the highest decrease was observed in T5 at 35, 50 and 65 DAT as -33.28, -26.14 and -  
257 29.50 respectively. Whereas, the highest increase of BPH incidence was found from T7 treatment at  
258 35, 50 and 65 DAT as +24.29, +15.58 and +17.26 respectively. Similar findings also reported by  
259 Sarwar (2012) earlier.

260 **Table 6. Incidence of brown plant hopper infestation in BRRI dhan33 for different levels of NPK**  
261 **fertilizers**

Treatment s	Incidence of brown plant hopper at different days after transplanting (DAT)					
	1 <sup>st</sup> observation (35 DAT)		2 <sup>nd</sup> observation (50 DAT)		3 <sup>rd</sup> observation (65 DAT)	
	Tillers infestation (%)	Reduction over control (%)	Tillers infestation (%)	Reduction over control (%)	Tillers infestation (%)	Reduction over control (%)
T0	6.34 b	--	6.16 b	--	6.78 b	--
T1	4.78 cd	-24.61	4.94 de	-19.81	5.03 de	-25.81
T2	6.12 b	-3.47	5.78 bc	-6.17	6.14 bc	-9.44
T3	5.86 b	-7.57	5.56 bcd	-9.74	5.94 c	-12.39
T4	5.22 c	-17.67	5.34 cd	-13.31	5.78 c	-14.75
T5	4.23 d	-33.28	4.55 e	-26.14	4.78 e	-29.50

T6	4.93 c	-22.24	5.23 cde	-15.10	5.55 cd	-18.14
T7	7.88 a	+24.29	7.12 a	+15.58	7.95 a	+17.26
LSD(0.05)	0.578	--	0.726	--	0.667	--
CV(%)	5.69	--	7.26	--	6.22	--

262 In a column, numeric data represents the mean value of 3 replications; each replication is derived  
 263 from each plot of each treatment

264 In a column means having similar letter(s) are statistically identical and those having dissimilar  
 265 letter(s) differ significantly as per 0.05 level of probability

266

T0= Untreated control, no NPK fertilizers

T1 = NPK @ 45, 50, 40 kg/ha

T2= NPK @ 70, 25, 40 kg/ha

T3 = NPK @ 70, 50, 20 kg/ha

T4 = NPK @ 70, 50, 40 kg/ha

T5 = NPK @ 70, 50, 60 kg/ha

T6 = NPK @ 70, 75, 40 kg/ha

T7 = NPK @ 95, 50, 40 kg/ha

267

268

### 269 3.13 Green leaf hopper incidence at 35, 50 and 65 DAT

270 For green leaf hopper incidence, at 35 DAT, 50 DAT and 65 DAT, data recorded from each plot  
 271 revealed that the lowest incidence of green leaf hopper was found from T5 (1.12%, 1.34% and 1.56%  
 272 respectively), whereas the highest incidence of green leaf hopper was observed from T7 (3.96%,  
 273 4.18% and 4.34% respectively) treatment. In case of incidence of Green leaf hopper  
 274 decrease/increase over control, the highest decrease was observed in T5 at 35, 50 and 65 DAT as -  
 275 59.71, -54.73 and -50.16 respectively. Whereas, the highest increase of Green leaf hopper incidence  
 276 was found from T7 treatment at 35, 50 and 65 DAT as +42.45, +41.22 and +38.66 respectively.  
 277 Mahadev *et al.* (1995) reported that the crop applied with N either alone or coupled with P exhibited a  
 278 higher degree of incidence of green leaf hopper.

279

280

281

282 **Table 7. Incidence of green leaf hopper infestation in BRRI dhan33 for different levels of NPK**  
 283 **fertilizers**

Treatment s	Incidence of green leaf hopper at different days after transplanting (DAT)					
	1 <sup>st</sup> observation (35 DAT)		2 <sup>nd</sup> observation (50 DAT)		3 <sup>rd</sup> observation (65 DAT)	
	Leaf infestation (%)	Reduction over control (%)	Leaf infestation (%)	Reduction over control (%)	Leaf infestation (%)	Reduction over control (%)
T0	2.78 b	--	2.96 b	--	3.13 b	--
T1	1.58 f	-43.17	1.88 e	-36.49	1.94 d	-38.02
T2	2.59 bc	-6.83	2.78 bc	-6.08	2.97 b	-5.11
T3	2.24 cd	-19.42	2.46 cd	-16.89	2.81 b	-10.22
T4	2.03 de	-26.98	2.14 de	-27.70	2.45 c	-21.73
T5	1.12 g	-59.71	1.34 f	-54.73	1.56 e	-50.16



T6	1.84 ef	-33.81	1.94 e	-34.46	2.12 cd	-32.27
T7	3.96 a	+42.45	4.18 a	+41.22	4.34 a	+38.66
LSD(0.05)	0.380	--	0.367	--	0.332	--
CV(%)	9.51	--	8.53	--	6.94	--

284 In a column, numeric data represents the mean value of 3 replications; each replication is derived  
 285 from each plot of each treatment

286 In a column means having similar letter(s) are statistically identical and those having dissimilar  
 287 letter(s) differ significantly as per 0.05 level of probability

288

T0= Untreated control, no NPK fertilizers

T2= NPK @ 70, 25, 40 kg/ha

T4 = NPK @ 70, 50, 40 kg/ha

T6 = NPK @ 70, 75, 40 kg/ha

T1 = NPK @ 45, 50, 40 kg/ha

T3 = NPK @ 70, 50, 20 kg/ha

T5 = NPK @ 70, 50, 60 kg/ha

T7 = NPK @ 95, 50, 40 kg/ha

289

290

### 291 3.14 Rice hispa incidence in leaf at 25, 40 and 55 DAT

292 For rice hispa incidence, at 25 DAT, 40 DAT and 55 DAT data recorded from each plot revealed that  
 293 the lowest incidence of rice hispa was found from T5 (1.15%, 1.46% and 1.66% respectively), whereas  
 294 the highest incidence of rice hispa was observed from T7 (4.86%, 5.14% and 5.83% respectively)  
 295 treatment. In case of incidence of Rice hispa decrease/increase over control, the highest decrease  
 296 was observed in T5 at 25, 40 and 55 DAT as -63.72, -58.87 and -56.08 respectively. Whereas, the  
 297 highest increase of Rice hispa incidence was found from T7 treatment at 25, 40 and 55 DAT as  
 298 +53.31, +44.79 and +54.23 respectively. Pathak *et al.* (1999) reported that minimum rice hispa  
 299 incidence in control plots and enhanced doses of N resulted in significant increase in rice hispa  
 300 incidence in rice field.

301

302 **Table 8. Incidence of rice hispa infestation in BRRl dhan33 for different levels of NPK fertilizers**

Treatment s	Incidence of rice hispa at different days after transplanting (DAT)					
	1 <sup>st</sup> observation (25 DAT)		2 <sup>nd</sup> observation (40 DAT)		3 <sup>rd</sup> observation (55 DAT)	
	Leaf infestation (%)	Reduction over control (%)	Leaf infestation (%)	Reduction over control (%)	Leaf infestation (%)	Reduction over control (%)
T0	3.17 b	--	3.55 b	--	3.78 b	--
T1	1.90 e	-40.06	1.67 g	-52.96	1.86 f	-50.79
T2	2.89 bc	-8.83	3.03 c	-14.65	3.25 c	-14.02
T3	2.56 cd	-19.24	2.74 d	-22.82	2.85 d	-24.60
T4	2.34 d	-26.18	2.43 e	-31.55	2.67 d	-29.37
T5	1.15 f	-63.72	1.46 g	-58.87	1.66 f	-56.08
T6	1.85 e	-41.64	2.05 f	-42.25	2.24 e	-40.74
T7	4.86 a	+53.31	5.14 a	+44.79	5.83 a	+54.23

LSD(0.05)	0.443	--	0.254	--	0.355	--
CV(%)	9.80	--	5.26	--	6.69	--

In a column, numeric data represents the mean value of 3 replications; each replication is derived from each plot of each treatment

In a column means having similar letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.05 level of probability

T0= Untreated control, no NPK fertilizers

T2= NPK @ 70, 25, 40 kg/ha

T4 = NPK @ 70, 50, 40 kg/ha

T6 = NPK @ 70, 75, 40 kg/ha

T1 = NPK @ 45, 50, 40 kg/ha

T3 = NPK @ 70, 50, 20 kg/ha

T5 = NPK @ 70, 50, 60 kg/ha

T7 = NPK @ 95, 50, 40 kg/ha

### 3.15 Rice bug incidence at 45, 55 and 65 DAT

In case of rice bug incidence, at 45 DAT, 55 DAT and 65 DAT data recorded from each plot revealed that the lowest incidence of rice bug was found from T5 (1.78%, 1.96% and 2.04% respectively), while the highest incidence of rice bug was observed from T7 (4.22%, 4.75% and 5.08% respectively) treatment. In case of incidence of Rice bug decrease/increase over control, the highest decrease was observed in T5 at 45, 55 and 65 DAT as -48.85, -46.30 and -46.60 respectively. Whereas, the highest increase of Rice bug incidence was found from T7 treatment at 45, 55 and 65 DAT as +21.26, +30.14 and +32.98 respectively. Mahadev *et al.* (1995) reported that the crop applied with N either alone or coupled with P exhibited a lower incidence of rice bug.

**Table 9. Incidence of rice bug infestation in BRR dhan33 for different levels of NPK fertilizers**

Treatment s	Incidence of rice bug at different days after transplanting (DAT)					
	1 <sup>st</sup> observation (45 DAT)		2 <sup>nd</sup> observation (55 DAT)		3 <sup>rd</sup> observation (65 DAT)	
	Panicle infestation (%)	Reduction over control (%)	Panicle infestation (%)	Reduction over control (%)	Panicle infestation (%)	Reduction over control (%)
T0	3.48 b	--	3.65 b	--	3.82 b	--
T1	2.15 e	-38.22	2.22 ef	-39.18	2.45 e	-35.86
T2	3.14 bc	-9.77	3.42 bc	-6.30	3.68 bc	-3.66
T3	2.95 cd	-15.23	3.13 bcd	-14.25	3.34 cd	-12.57
T4	2.84 cd	-18.39	2.97 cd	-18.63	3.12 d	-18.32
T5	1.78 e	-48.85	1.96 f	-46.30	2.04 f	-46.60
T6	2.56 d	-26.44	2.64 de	-27.67	2.67 e	-30.10
T7	4.22 a	+21.26	4.75 a	+30.14	5.08 a	+32.98
LSD(0.05)	0.403	--	0.514	--	0.367	--
CV(%)	7.66	--	9.14	--	6.19	--

In a column, numeric data represents the mean value of 3 replications; each replication is derived from each plot of each treatment  
In a column means having similar letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.05 level of probability

T0= Untreated control, no NPK fertilizers	T1 = NPK @ 45, 50, 40 kg/ha
T2= NPK @ 70, 25, 40 kg/ha	T3 = NPK @ 70, 50, 20 kg/ha
T4 = NPK @ 70, 50, 40 kg/ha	T5 = NPK @ 70, 50, 60 kg/ha
T6 = NPK @ 70, 75, 40 kg/ha	T7 = NPK @ 95, 50, 40 kg/ha

#### 4. CONCLUSION

Considering the above finding it is revealed that if the nitrogen dose increases, the incidence of insect pest also increased. In the term of percent of infestation, NPK @ 70, 50, 60 kg/ha (T5) was the better for the incidence of insect pest. However, imbalanced nutrition like 100% N, 100% P, 100% K etc. incited more insect incidence, whereas balanced nutrients resulted in lower incidence of insect pests. Among the different levels of NPK fertilizers, NPK @ 70, 50, 60 kg/ha was the better for the rice cultivation. Considering the situation of the present experiment, further studies in the following areas may be suggested: Different pest management practices and rice variety may be use in future study. This experiment should be carried out in different agro-ecological zones AEZ) of Bangladesh for confirmation of the results.

#### 5. REFERENCES

- Hien, N.L., Yoshihashi, T. and Sarhadi, W.A. (2006). Evaluation of aroma in rice using KOH method, molecular markers and measurement of 2-acetyl-1pyrroline concentration. *Japanese J. Trop. Agric.* 50: 190-198.
- Barr, B.A.S. and Smith, R.F. (1975). Crop loss to insects, diseases and other pests. USAID, Pest management and related environmental protection project. University of California, Berkley. p. 64.
- Mustafi, B.A.A., Salam, M.A., Mondal, M. and Kalam, M.A. (2007). Modern cultivation of rice (In Bangla). Bangladesh Rice Research Institute, Gazipur, Bangladesh. p. 66.
- Miah, S.A. and Karim. (1984). Rice pest management technology, BRRI, Gajipur. pp. 43-49.
- Siddique, A.K.M.T. (1992). "An overview of the ETL of rice pest and 1PM scope in the content of M.V. Rice production" In Proc. of the Workshop on Experiences with Modern Rice Cultivation in Bangladesh. BRRI, Gazipur. pp. 33-45.
- Rahman, M.T., Khalequzzaman, M. and Khan, M.A.R. (2004). Assessment of infestation and yield loss by stem borers on variety of rice. *J. Asia Pacific Entomol.* 7(1): 89-95.
- BRRI (Bangladesh Rice Research Institute) (1985). Dhan Chaser Samashy. BRRI, 1985. Annual Report for 1984. Bangladesh Rice Research Institute, Gazipur. p. 207.
- Magdoff, F. and Havan, E. (2000). Building soils for better crops. SARE, Washington DC.
- Wahiduzzaman, M. (1993). Studies of the insect predatory behavior of the black drongo in rice ecosystem. M.S. thesis, Institute of Post Graduate, Delhi, India.
- Vaithilingan, C. (1975). Impact of potassium on pests of rice. *Entomol. views.* p. 126.
- Subramanian and Balasubramanian. (1976). Influence of Phosphorus fertilizer on rice pests. *Madraj journal.* pp. 561-565.
- Gomez, K.A. and Gomez, A.A. (1984). Statistical procedure for agricultural research. International Rice Research Institute. John Wiley and Sons, New York. pp. 139-240.
- Rangini, J.C., Thangaraju, D. and David, P.M.M. (2005). Relative abundance of rice stem borer species in Tamil Nadu. *Madras Agril. J. Pub.* 87(4-6): 228-234.
- de Kraker J. The potential of natural enemies to suppress rice leaffolder populations. [PhD thesis]. Wageningen: Wageningen Agricultural University, 1996.
- Singh, G., Kathapal, T.S. and Gupta, S.P. (1990). Leaching of phorate and carbofuran in soils as affected by zinc. *J. of Ecology.* 17(2): 132-135.
- Madhuri G. (2016). Effect of different levels of fertilizers on incidence of arthropod fauna on rice. MS thesis. Orissa University of Agriculture and Technology, India.
- Nath, P. and Bhagabati, K.N. (2005). Population dynamics of leaf hopper vectors of Rice tungro virus in Assam. *Indian Phytopatho.* 55(1): 92-94.
- Tsueda, H., Yajima, M., Taguchi, Y. and Suzuki, T. (2002). Occur rent tendency of rice sting bugs in South flat area of Gifu Prefecture. *Proc.The Kansai Plant Protection Soc.* 4: 13-20.

- 377 19.Ramzan, M., Hussain, S. and Akhter, M. (1992). Incidence of insect pests on rice crop under  
378 various nitrogen doses. *J. of Animal Plant Science*. 17(3-4): 20.
- 379 20.Chakraborty K. (2011). Influence of inorganic N fertilizer on plant characters, yield generation and  
380 the incidence of yellow stem borer in the field of local scented paddy cultivar Tulaipanji.  
381 *International J. of Applied Biology and Pharmaceutical Technology*. 2(4): 305-309.
- 382 21.Mahadev, P., Das, M., Ghose, M.R. and Mukherjee, N. (1995). Effect of fertilizer treatments on  
383 growth, productivity, insect pests and diseases incidence on rice. *Madras Agricultural J*.  
384 82(9/10): 525-529.
- 385 22.Sarwar,M. (2012). Effects of potassium fertilization on population buildup of rice stem borers  
386 (lepidopteron pests) and rice (*Oryza sativa* L.) yield. *J. of Cereals and Oil seeds*. 3(1): 6-9.
- 387 23.Pathak, K.A. and Ram, S. (1999). Effect of fertilizers (NPK) on the incidence of pest complex of  
388 paddy in Manipur. *Indian J. of Entomology*. 4(2): 409-411.