

Control of Rhizome Rot Disease of Ginger through Application of Fungicides

Md. Kamal Hosain¹, Sk. Md. Mobarak Hossain¹, Zoynal Abedin², Md. Mukul Islam^{1*},
Md. Atiar Rahman³, Md. Shazadur Rahman⁴ and Md. Abu Zafor⁵

¹Department of Plant Pathology, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh.

²Kazi and Kazi Tea Estate LTD., Panchagarh, Bangladesh.

³Department of Agroforestry and Environment, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh.

⁴Department of Agricultural Chemistry, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh.

⁵Department of Soil Science, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh.

Authors' contributions

This work was carried out in collaboration between all authors. Authors MKH, SMMH and MMI designed the study, wrote the protocol and wrote the first draft of the manuscript. Authors ZA and MAR managed the literature searches and analysis of the study. Author MSR and MAZ managed the experimental process. All authors read and approved the final manuscript.

Corresponding author E-mail: mukul.hstu11@gmail.com

ABSTRACT

Aims: To determine the effect of seed treatment and foliar spray of fungicide on rhizome rot of ginger.

Study Design: The study was laid out in a randomized complete block design with three replications.

Place and Duration of Study: The experiment was conducted at the plant pathology Field Laboratory Of Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh during 2016-2017.

Methodology : Fungicide namely Contuf 5 EC @ 1ml/L water, Bavistin DF @ 1 g/L water, Companion @ 2 g/L water, Fiasta Z-78 @ 2 g/L water, Matco 72 WP @ 2 g/L water, Blitox 50 WP @ 2 g/L water, Cabriotop @ 3 g/L water were applied as seed treatment and foliar spray with one control plot (without fungicide). The total number of unit plots was 24 and the size of unit plot was 15 m × 7 m.

Results: Fungicide Cabriotop was found as the most effective in increasing plant growth and by reducing disease severity of rhizome rot of ginger. Among the treatments, capriotop exhibited least disease severity (45.08%) of rhizome and maximum yield (4.13 t/ha) of ginger as compared to control plot (69.04% disease severity and 0.33 t/ha rhizome).

Conclusion: Application of the fungicide Cabriotop may be recommended for better performance in controlling of rhizome rot disease of ginger.

Keywords: Ginger, rhizome rot, fungicides, disease severity

1. INTRODUCTION

Ginger (*Zingiber officinale* Rosc.) under the family zingiberaceae is one of the most important spices crops in world as well as in Bangladesh. It is an herbaceous tropical and sub-tropical perennial plant and seems to have originated in South-East Asia, probably in India [1,2]. Ginger is an important commercial

crop in tropical and sub-tropical countries. It has medicinal value, particularly in traditional medicines of India [3]. Fresh ginger, ginger powder from dry ginger and oil are all used for this purpose. Fresh ginger is unique for its flowery flavor and spicy taste. It is also used in jams and marmalades. The syrup in which ginger is preserved is valued for pickle and sauce making. It is also used in the production of ginger bread [4]. In addition, ginger is used popularly for chewing purposes [5]. Ginger is cultivated in several parts of the world, and the most important countries viz. India, China, Nigeria, Sierra Leone, Indonesia, Bangladesh, Australia, Fiji, Jamaica and Nepal. Among them, India and China are the dominant suppliers to the world market [6]. In Bangladesh, the yield of ginger is not enough to fulfill the national demand of the country. Thus, a huge amount of ginger has to be imported from abroad to fulfill the national demand. Disease is a major constraint for the production of healthy rhizomes and can cause even total failure of crop [7]. Ginger is affected by various diseases, such as, rhizome rot, bacterial wilt, soft rot, leaf blight etc. Among all of these, rhizome rot is the most damaging [8]. Rhizome rot or soft rot is a highly destructive disease in ginger; in some areas of the world. Soft rot is known to destroy 80 to 90% yield of the annual crop [9]. In Bangladesh, rhizome rot caused by *Pythium* spp. which can survive in soil for long periods of time results in loss of total production if infection initiates at early stage of plant growth. Yield of ginger is drastically reduced by water logging of the soil. The infected rhizomes become rotten and the crop is completely destroyed [10]. The disease is important because it causes economic losses to growers resulting in decreased prices of products to the consumers. It is very important to identify the factors affecting disease severity and to develop control measures for rhizome rot. Control measures such as seed treatment, soil treatment, soil amendment, sanitation, drainage, intercropping etc. have some effect in controlling the disease [11]. Rhizome rot of ginger can be controlled by the application of fungicides viz. Contuf 5 EC, Bavistin DF, Companion, Fiasta Z-78, Matco 72 WP, Blitox 50 WP, Cabriotop, etc. Many researchers have investigated the chemical control of the disease with promising results [12,13]. Systemic and contact fungicides like Bavistin 50WP, Ridomil Gold MZ-72, Captan, Dithane M-45, Copper Oxychloride and Bordeaux mixture etc. were reported effective against the disease [6]. Therefore, the present study was undertaken to identify the most effective fungicide available in Bangladesh for controlling rhizome rot disease of ginger.

2. MATERIALS AND METHODS

2.1 Experimental site: The experiment was conducted at the Plant Pathology Research Field of Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh.

2.2 Duration of the Experiment: The experiment was conducted during April, 2016 to January, 2017.

2.3 Soil type: Soil of the experimental plot was sandy loam with good drainage capacity. The experimental plot was medium high land with the p^H range from 5.5 to 6.0.

2.4 Variety and preparation of propagating unit: Local variety rhizomes of diseased free having 1-2 buds whose average weight 55-60 g. were selected. The collected rhizomes were preserved under soil about one month for pre-sprouting of seed.

2.5 Preparation of the main field: The selected land was harrowed, ploughed and cross-ploughed several times followed by laddering to get a fine tilth. Weeds, stubbles and rubbishes were removed, proper drainage channels were made around the experimental plots to create a desirable tilth of soil for planting of ginger rhizomes.

2.6 Application of manure and fertilizers: Well decomposed cow-dung @ 7 t/ha was applied during land opening. TSP @ 4 t/ha and MOP @ 3 t/ha, Gypsum @ 2 t/ha were applied at the time of final land preparation. The entire amount of TSP, MOP and Gypsum were applied during the final land preparation of land. Urea was applied after first weeding @2 t/ha.

2.7 Design of the experiment: Single factor RCBD (Randomized Complete Block Design)

2.8 Layout of the experiment: The experiment was conducted with 3 replications having 8 treatments.

2.9 Treatment of the experiment: The treatments were as follows:

- T₀ = Untreated (Control),
- T₁ = Treatment of ginger with Contaf 5 EC@1 ml /L,
- T₂ = Treatment of ginger with Bavistin DF@1 g /L,
- T₃ = Treatment of ginger with Companion@2 g /L,
- T₄ = Treatment of ginger with Fiasta Z-78@2 g /L,
- T₅ = Treatment of ginger with Matco 72 WP@2 g /L,
- T₆ = Treatment of ginger with Blitox 50 WP@2 g /L,
- T₇ = Treatment of ginger with Cabriotop@3 g /L.

The treatments were applied at two times. Firstly, at the time of sowing of rhizomes by dipping them in the treatment solutions for 15 minutes and finally sprayed on the plants at 60 DAS (days after sowing).

2.10 Time of planting of rhizomes: Pre-sprouted rhizomes were planted at 2nd week of April, 2016 at the rate of 1.6 t/ha.

2.11 Sowing of rhizome: Pieces of seed rhizome were sown at the rate of 65g (1 rhizome) per hole. The seeds (rhizome) were placed individually in the furrows and furrow was covered with soil. The plots were earthen up 20 cm high from the level of drain. Finally, the plot was covered with straw of dry rice.

2.12 Data collecting parameters: Data were collected on no. of tillers per plant, plant height (cm), no. of leaves per plant, healthy plant per plot, infected plant per plot and disease severity of plant (%) before harvesting. Data were recorded on 60, 90, 120, 150 and 180 days after sowing (DAS). Disease severity of plant (%) was calculated by using the formula:

$$\text{Disease severity of plants} = \frac{\text{Surface area of plants infected by disease}}{\text{Total surface area of plants}} \times 100$$

Healthy rhizome, diseased rhizome, wt. of healthy rhizome, wt. of diseased rhizome, percent disease severity and yield (t/ha) per plot were recorded at the time of harvest.

2.13 Harvesting: Date of harvesting of ginger was 15th January, 2017. Rhizomes from each plot were harvested separately. The weight of rhizomes was recorded in each plot in kg and it was converted into hectare.

2.14 Data analysis: Collected data were analyzed statistically by using the MSTAT-C computer package program [12].

3. RESULTS

3.1 Effect of fungicides on tiller number per plant

Tiller number per plant of ginger was recorded at 60, 90, 120, 150 and 180 DAS and the results are presented in Table 1. When tiller number per plant of ginger was recorded at 60 DAS, the highest (5.297) tiller number per plant was found in T₇, where ginger was planted in Cabriotop treated plots which was statistically similar with T₂, T₆, with the lowest (3.607) was in T₀. The highest (10.28) tiller number per plant at 90 DAS was found in T₇, where ginger was planted in Cabriotop treated plots which was statistically similar with T₂, T₄, T₅ and T₆ and the lowest (6.140) tiller number per plant was found in T₀. At 120 DAS tiller number per plant maximum was (18.32) in T₇, which was statistically similar with T₁, T₂, T₃, T₄, T₅ and T₆ and the minimum 13.85) tiller number per plant was observed in T₀. At 150 DAS, tiller number per plant ranged from 17.10 to 20.34 where the highest (20.34) was found in T₅, which was statistically similar with T₁, T₂, T₃, T₄, T₆ & T₇ and the lowest (17.10) again was in control. Tiller number per plant at 180 DAS, was

recorded the highest (20.91) in T₇ was statistically similar with T₁, T₂, T₃, T₄, T₅ and T₆ and the lowest (17.55) tiller number per plant was found in T₀.

Table 1. Effect of fungicides on tiller number per plant at different dates of observation

Treatments	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
Control (T ₀)	3.61 c	6.14 d	13.85 b	17.10 b	17.55 b
Contaf 5 EC (T ₁)	3.80 bc	7.80 c	16.76 a	18.69 ab	19.25 ab
Bavistin DF (T ₂)	5.19 a	9.55 ab	17.88 a	20.20 a	20.54 ab
Companion (T ₃)	4.25 bc	8.72 bc	17.46 a	19.61 ab	20.03 ab
Fiasta Z-78 (T ₄)	4.42 b	10.1 a	16.68 a	19.46 ab	20.04 ab
Matco 72 WP (T ₅)	4.29 bc	9.66 ab	17.22 a	20.34 a	20.86 a
Blitox 50 WP (T ₆)	5.25 a	9.40 ab	17.20 a	18.08 ab	18.71 ab
Cabriotop (T ₇)	5.30 a	10.29 a	18.32 a	20.24 a	20.91 a
LSD	0.71	1.25	1.86	2.58	2.86
CV %	8.99	8.01	6.30	7.68	8.28

Figures in a column having common letter(s) do not differ significantly at 5% level of significance. CV = Coefficient of variation;

3.2 Effect of fungicides on plant height (cm)

Plant height (cm) of ginger was recorded at 60, 90, 120, 150 and 180 DAS and the results are shown in Table 2. Plant height was recorded maximum (61.77 cm) in T₆ at 60 DAS, where ginger was planted with Blitox 50 WP treated plot which was statistically similar with T₁, T₄, T₅ and T₇ and the minimum (56.44) plant height was observed in T₀, where ginger was planted with control plot. Plant height at 90 DAS was maximum (71.63) in T₇ statistically similar with T₁ and T₄. The minimum (66.15 cm) plant height was found in T₃ was also statistically similar with T₀, T₂, T₅, and T₆. At 120 DAS, the maximum (81.05cm) plant height was found in T₇ was statistically similar with T₁, T₂, T₃, T₄, T₅ and T₆ while the minimum (76.33cm) plant height in T₀. However at 150 180 DAS, no significant differences were observed among the treatments.

Table 2. Effect of fungicides on plant height (cm) at different dates of observation

Treatments	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
Control (T ₀)	56.44 c	68.37 b	76.33 b	85.92 a	60.75 a
Contaf 5 EC (T ₁)	60.22 abc	69.84 ab	77.82 ab	81.07 a	79.86 a
Bavistin DF (T ₂)	59.56 abc	68.57 b	78.93 ab	81.66 a	82.19 a
Companion (T ₃)	56.78 bc	66.15 c	77.56 ab	79.45 a	80.18 a
Fiasta Z-78 (T ₄)	58.57 abc	69.95 ab	79.43 ab	82.27 a	82.69 a
Matco 72 WP (T ₅)	59.57 abc	68.62 b	77.89 ab	80.12 a	80.55 a
Blitox 50 WP (T ₆)	61.77 a	68.85 b	78.96 ab	81.32 a	81.81 a
Cabriotop (T ₇)	61.43 ab	71.63 a	81.05 a	82.82 a	83.25 a
LSD	4.31	2.04	3.16	7.42	22.22
CV %	4.15	1.69	2.30	5.18	16.08

Figures in a column having common letter(s) do not differ significantly at 5% level of significance. CV = Coefficient of variation;

3.3 Effect of fungicides on leaves/plant

Number of leaves/plant of ginger was recorded at 60, 90, 120, 150 and 180 DAS and the results are shown in Table 3. At 60 DAS, the highest (16.55) leaves/plant was found in T₁, statistically similar with T₂, T₃, T₄, T₆ and T₇ while the lowest (14.43) number of leaves/plant was recorded in T₀. Leaves/plant was recorded maximum (18.04) in T₇ at 90 DAS, which was statistically similar with other treatments except T₀ where the minimum (16.10) number of leaves/plant was observed. However at 120 DAS, no significant differences were observed among the treatments. The highest (30.68) leaves/plant were found in T₁ at 150 DAS, which was statistically similar with T₂, T₃, T₅ and T₇. On the other hand, the lowest (28.28)

leaves/plant were found in T_0 , was statistically similar to T_6 (28.41). At 180 DAS, all the treatments showed statistically significant results except T_0 where the lowest (26.31) leaves/plant were recorded.

Table 3. Effect of fungicides on leaves/plant at different dates of observation

Treatments	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
Control (T_0)	14.43 b	16.10 c	26.89 a	28.28 c	26.31 b
Contaf 5 EC (T_1)	16.55 a	17.81 ab	27.52 a	30.68 a	31.38 a
Bavistin DF (T_2)	15.88 ab	17.74 ab	27.12 a	30.16 abc	30.56 a
Companion (T_3)	15.35 ab	16.36 bc	26.85 a	29.47 abc	29.95 ab
Fiasta Z-78 (T_4)	15.79 ab	17.05 abc	26.80 a	28.43 bc	28.80 ab
Matco 72 WP (T_5)	15.93 ab	16.88 abc	27.02 a	30.22 ab	30.79 a
Blitox 50 WP (T_6)	15.55 ab	17.37 abc	26.64 a	28.41 bc	29.10 ab
Cabriotop (T_7)	15.92 ab	18.04 a	27.10 a	30.42 a	30.80 a
LSD	1.56	1.40	1.78	1.72	3.57
CV %	5.71	4.66	3.78	3.33	6.87

Figures in a column having common letter(s) do not differ significantly at 5% level of significance. CV = Coefficient of variation;

3.4 Effect of fungicides on healthy plants/plot at different

Healthy plants/plot of ginger was recorded at 60, 90, 120, 150 and 180 DAS and the results are shown in Table 4. In case of healthy plants per plot was found significant only at 60 DAS while at 90, 120, 150 and 180 DAS no significant differences were observed among the treatments. The highest (12.67) number of healthy plants/plot was found in T_7 at 60 DAS was statistically similar with T_1 , T_2 , T_3 , T_5 and T_6 while the lowest (10.67) healthy plant/plot was found in T_0 .

Table 4. Effect of fungicides on healthy plant/plot at different dates of observation

Treatments	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
Control (T_0)	10.67 c	9.33 a	8.00 a	6.67 a	5.00 a
Contaf 5 EC (T_1)	12.33 ab	10.33 a	9.33 a	8.33 a	6.00 a
Bavistin DF (T_2)	12.33 ab	10.33 a	9.00 a	6.66 a	5.00 a
Companion (T_3)	12.00 ab	10.00 a	8.33 a	6.33 a	5.33 a
Fiasta Z-78 (T_4)	11.33 bc	9.33 a	7.66 a	6.33 a	5.33 a
Matco 72 WP (T_5)	12.33 ab	10.33 a	8.66 a	7.33 a	6.00 a
Blitox WP 50 (T_6)	11.67 abc	10.00 a	9.00 a	7.00 a	5.33 a
Cabriotop (T_7)	12.67 a	11.00 a	9.00 a	7.66 a	6.00 a
LSD	0.97	1.68	2.33	2.49	2.55
CV %	4.67	9.56	15.47	20.20	26.50

Figures in a column having common letter(s) do not differ significantly at 5% level of significance. CV = Coefficient of variation;

3.5 Effect of fungicides on infected plants/plot

Infected plants/plot of ginger was also recorded at 60, 90, 120, 150 and 180 DAS and the results are shown in Table 5. Only at 60 DAS infected plants/plot was found significant while at 90, 120, 150 and 180 DAS no significant differences were observed among the treatments. At 60 DAS, infected plant per plot was highest (3.33) in control plots, which was similar with T_4 (2.67) and T_6 (2.33). However the lowest (1.33) number of infected plant per plot was recorded in T_7 .

Table 5. Effect of fungicides on infected plant/plot at different dates of observation

Treatments	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS
Control(T_0)	3.33 a	4.67 a	6.00 a	7.33 a	9.00 a

Contaf 5 EC(T ₁)	1.67 bc	3.67 a	4.67 a	5.67 a	8.00 a
Bavistin DF(T ₂)	1.67 bc	3.67 a	5.00 a	7.33 a	9.00 a
Companion(T ₃)	2.00 bc	4.00 a	5.67 a	7.67 a	8.67 a
Fiasta Z-78(T ₄)	2.67 ab	4.67 a	6.33 a	7.67 a	8.67 a
Matco 72 WP(T ₅)	1.67 bc	3.67 a	5.33 a	6.67 a	8.00 a
Blitox 50 WP(T ₆)	2.33 abc	4.00 a	5.00 a	7.00 a	8.67 a
Cabriotop (T ₇)	1.33 c	3.00 a	5.00 a	6.33 a	8.00 a
LSD	0.97	1.69	2.33	2.49	2.55
CV %	26.70	24.60	24.82	20.44	17.15

Figures in a column having common letter(s) do not differ significantly at 5% level of significance. CV = Coefficient of variation;

3.6 Effect of fungicides on disease severity of plant

Percent disease severity of ginger was recorded at 60, 90, 120, 150 and 180 DAS and the results are presented in Figure 1. At 60 DAS, disease severity varied significantly among the treatments and recorded minimum in T₆ while the maximum was in T₀. However, at the later stages of plant growth at 90, 120, 150 and 180 DAS disease severity were recorded maximum in T₀ while the minimum was found in T₇.




Fig.1. Effect of fungicides on disease severity of plant at different dates of observation

3.7 Effect of fungicide on healthy rhizome, diseased rhizome, percent disease severity and yield (t/ha) of rhizome

Effect of healthy rhizome, disease rhizome and yield (t/ha) of rhizome were recorded after **harvests and are** presented in the Table 6. Number and weight of healthy rhizome was recorded the highest in **the** Cabriotop treated plot followed by Contuf and Bavistin treated **plots**. On the other hand, the lowest number and weight of healthy rhizome was recorded in the Control treatment. Number and weight of disease rhizome was found maximum in the control treatment and minimum in Cabriotop treatment. Disease severity of rhizome was recorded maximum in control plot and the minimum was in Cabriotop treatment treated plot. Yield of the rhizome recorded the highest in Cabriotop treatment and minimum in Control treatment.

Table 6. Effect of fungicide on healthy rhizome, diseased rhizome, percent disease severity and yield (t/ha) of rhizome.

Treatments	Healthy rhizome		Diseased rhizome		Disease severity of rhizome	Yield (t/ha)
	Number /plot	Weight /plot	Number /plot	Weight /plot		
Control (T ₀)	5.00 e	0.15 c	28.33 a	0.73 a	71.03 a	0.33 c
Contaf 5 EC (T ₁)	17.67 b	0.84 b	14.67 bc	0.62 ab	59.67 a	1.84 b
Bavistin DF (T ₂)	12.33 c	0.36 c	23.67 abc	0.60 ab	65.88 a	0.76 c
Companion (T ₃)	10.33 cd	0.27 c	17.67 abc	0.37 bc	63.01 a	0.58 c
Fiasta Z-78 (T ₄)	8.33 cde	0.33 c	16.33 abc	0.49 abc	65.25 a	0.70 c
Matco 72 WP (T ₅)	7.00 de	0.35 c	15.33 abc	0.51 abc	69.04 a	0.74 c
Blitox 50 WP (T ₆)	6.33 de	0.34 c	27.00 ab	0.43 abc	70.32 a	0.72 c
Cabriotop (T ₇)	32.67 a	1.92 a	12.33 c	0.28 c	45.08 b	4.13 a
LSD	4.18	0.29	11.95	0.28	11.83	0.63
CV %	19.18	35.16	29.60	31.76	10.61	29.42

Figures in a column having common letter(s) do not differ significantly at 5% level of significance. CV = Coefficient of variation;

4. DISCUSSION

Rhizome rot is ranked one of the most destructive disease of ginger which may results in total loss of an affected crop. The effect of fungicides was investigated to minimize the losses caused by the disease. The results of the present study revealed that the fungicides Capriotop and Contaf 5 EC have an impact on early plant growth, rhizome yield with reduced disease incidence and severity. Sasikumar et al. [13] reported that the treatment of seed rhizomes with mancozeb 0.3 percent for 30 minutes before storage and once again before planting reduces the incidence of disease. In the present study Contaf 5 EC, Bavistin DF, Companion, Fiasta Z-78, Matco 72 WP, Blitox 50 WP and Cabriotop proved their effectiveness to increase healthy rhizome/plant compared to control. The results of the present experiment initially showed significant variation among the treatments on tiller number per plant and plant height. Significantly higher tiller number per plant and plant height were recorded when seed treatment and foliar spray were applied with Capriotop compared to control at 180 DAS. This might have happened due to reduction of primary and secondary inocula of pathogen present in seed and soil. The application of fungicide may also inhibit the growth of fungi. Similar result was reported by Rahman [11] who found that number of tiller per plant was the highest in Ridomil and Secure treated plot and the plant in control plot is shorter than the fungicidal treated plot. Under the present investigation, the fungicide Cabriotop was found as the most effective for yielding maximum number of leaves per plant compared to control treatment. Other fungicides tested also had some influence on number of leaves per plant over control. Maximum number of leaves per plant was the indication of minimum amount of disease. Initially, all the fungicide treated plots produced lower percent of infected plants compared to control treated. However, at 60 DAS the higher percent of healthy plants were found in fungicide treated plots compared to the control treated. The present findings are envisaged with the findings of Rahman [11], BARI [14] and BARI [15] all mentioned that Ridomil resulted in the lowest number of dead plants per plot whereas the highest was found in control. Ara [16] reported that the use of Secure and Antracol as seed treatment decreased the number of rhizome rot infected plants. Ramachandran et al. [17] also reported that five systemic fungicides namely Fosetyl aluminium, Metalaxyl, Oxadiazyl 25 WP, Propamocarb and Ethazole gave the best result in controlling rhizome rot of ginger. Although in this experiment, different fungicides were used as a treatment but Cabriotop treated plots resulted as the best by reducing disease severity and number of rhizome rot infected plants with highest yield. The present findings are in line with Hossain et. al. [18] who reported that fungicide Antracol, Follicur and Secure were effective in reducing the incidence and severity of rhizome rot and improved plant growth. Ichitani [19] also reported that rhizome rot incidence was reduced by seed treatment with fungicides namely Echlomezol and Methyl bromide. In case of yield of ginger, the plots treated with Cabriotop resulted in significantly highest yield of ginger (4.13 t/ha). The result of the present study was also supported by many other scientists who conducted different

experiments by using various fungicides. Ghorpade and Ajri [20], Dohroo and Sharma [21], Rathaiah [22] and Jayasekhar et al. [23] also reported the highest seed germination, lowest disease incidence and more yield of ginger through application of fungicide Ridomil.

5. CONCLUSION

From the present study, it can be concluded that the application of different fungicides have significant importance on plant height, number of tillers, number of leaves, infected plants, disease severity of plants and yield of ginger. But among all the treatments, Capriotop was found as the best with highest yield (4.13 t/ha) of ginger by increasing other yield contributing parameters and reducing disease severity.

REFERENCES

1. Burkill IH, Birtwistle W, Foxworthy FW, Scrivenor JB, Watson JG. A Dictionary of the Economic products of the Malaysia, Ministry of Agriculture and Co-operatives, Peninsula, Kuala Lumpur. 1966.
2. Purselove JW, Brown EG, Green CL, Robbins SRJ. Spices. Longman Inc. New York. USA. 1981;2(2).
3. Selvan MT, Thomas KG, Manojkumar K. Ginger (*Zingiber officinale* Rose). In: Singh HP, Sivaraman K, Selvan MT (editors). Indian Spices- Production and Utilization. Coconut Development Board, India. 2002;110-131.
4. Pruthi JS. Major Spices of India-Crop Management Post Harvest Technology. Indian Council of Agril. Res. New Delhi. 1993;12.
5. Purselove JW, Brown EG, Green CL Robbins SRJ.. Spices. Co-published in the United States with John Wiley & Sons. Inc. New York. 1988;2 (8): 447-462 & 2(9):533-540.
6. Sagar SD. Investigations on the etiology, epidemiology and integrated management of rhizome rot complex of ginger and turmeric. Ph.D. Thesis, Department of Plant Pathology, University of Agricultural Sciences, Dharwad. 2006.
7. Fageria MS, Choudhary BR, Dhaka RS. Vegetable crops production technology. Kalyani Publisher, New Delhi. 2006;11:223-227.
8. Chattopadhyay SB. Disease of Plants yielding drugs, dyes and spices. Indian council of Agric. Res. New Delhi. 1997;2(5): 66-67.
9. Dake GN. Diseases of ginger (*Zingiber officinale* Rose) and their management. J. Spices and Aromatic Crops. 1995;4:40-48.
10. Baruah HK, Baruah P, Baruah A. Text Book of Plant Pathology. Published by Mohan Primalani, Oxford and IBH publishing Co.66 Janpath, New Delhi. 1998;304-308.
11. Rahman MM. Integrated management of rhizome rot of ginger. M. S. Thesis, Dept. Plant Pathol. BAU, Mymensingh. 2001;65-69.
12. Gomez KA, Gomez AA. Duncan's Multiple Range Test. Statistical Procedure for Agril. Res. 2nd ed. A Wiley Inter-Science Publication, John and Sons, New York. 1984;202-215.
13. Sasikumar B, Thankamani CK, Srinivasan V, Devasahayam S, Santhosh J, Eapen KA, Zachariah JT. Ginger (Extension Pamphlet). 2009.
14. Bangladesh Agricultural Research Institute (BARI). Annual Report 2006-2007. Bangladesh Agril. Res. Inst. Joydebpur, Gazipur. 2007;159-160.
15. Bangladesh Agricultural Research Institute (BARI). Annual Report 2009-2010. Bangladesh Agril. Res. Inst. Joydebpur, Gazipur. 2010;172.
16. Ara A. Intercropping of ginger with indigenous plants and seed treatment with fungicides to control rhizome rot of ginger. M. S. Thesis, Dept. of Plant Pathol. HSTU, Dinajpur. 2013;51-57.
17. Ramachandran N, Dake GN, Sharma YR. Evaluation of systemic fungicides for efficacy against rhizome rot of ginger. Indian Phytopath. 1989;42(4):530-533.
18. Hossain SMM, Hasan MA, Alam MM. Study on chemical control of rhizome rot disease of ginger (*Zingiber officinale* Rose.). J. Sci. and Tech. 2015;13:75-81.
19. Ichitani T. Control of rhizome rot of ginger cultivated successively and protectively for immature rhizome production in plastic house. Proceedings of the Kansai Plant Prot. Society. 1980;22:7-11.

20. Ghorpade SA, Ajiri DS. Effectiveness of oilseed cakes in control of rhizome rot malady of ginger. J. Maharashtra Agric. Univ. 1982; 272-273.
21. Dohroo NP, Sharma SL. Evaluation of fungicides for the control of rhizome rot of ginger in storage. Indian Phytopathol. 1983;36:691-93.
22. Rathaiah Y. Control of soft rot of ginger with Ridomil. Dep. pl. Path. Coll. Agric. Dhawad, India. 1987;68:112.
23. Jayasekhar M, Prem Joshua J, Pillai AAO. Management of rhizome rot of ginger caused by *Pythium aphanidermatum*. Madras Agricultural J. 2000;87:170-171.