

**SORGHUM /LEGUME INTERCROP ON STEM  
BORER DAMAGE  
AND YIELD OF SORGHUM IN THE SOUTH  
EASTERN DRY AREAS OF ZIMBABWE.**

**Abstract**

*Stem borer (*Chilo partellus*) is one of the major pest of economic importance which affects sorghum production in the South eastern region of Zimbabwe. The experiment to establish the relationship between stem borer insect suppression by intercropping and grain yield in sorghum and six legumes was conducted under field conditions at Chiredzi Research Station which is in the South Eastern Lowveld (21°01'S, 31°33'E) from 2013 to 2015 cropping seasons. Treatments laid in randomized complete block design and replicated three times consisted of sorghum combined with cowpea, groundnut, pigeon pea and bambara. Monocropped treatments of legumes were cowpea, groundnut, pigeon pea, bambara and two sole sorghum treatments as controls. On one of the controls, an insecticide was applied while the other one remain untreated. Data collected on stem tunnel length, yield (grain and stover), land equivalent ratio and other pests, indicated that in sorghum sole plots where no chemical was applied, yield was reduced by 28% compared to sole plots where a pesticide was applied. In intercropped combinations of sorghum/groundnut, sorghum/pigeon pea and sorghum/cowpea, an incremental benefit of 10-38% was observed than all other treatments. No benefit was observed in sorghum bambara combinations. Stem borer and aphid had lower thresholds in intercrops. Predators populations recorded in intercrops reduced insect pest density than in monocrops. Thus, these findings indicated that intercropping can form a component of an integrated pest management program*

**Key words**

*Sorghum, legume, intercropping, land equivalent ratio, stem borer, yield*

33

## 34 1.0 INTRODUCTION

35

36 Intercropping legumes and non-legumes is an agricultural practice of cultivating two or more  
37 crops in the same place of land at the same time which is commonly practiced in many parts  
38 of the world in order to increase the productivity per unit area of the land (1). The crops are  
39 not necessarily sown at the same time and their harvest time may be quite different, but they  
40 are simultaneously grown for significant growing periods (2). Moreover, intercropping allows  
41 efficient use of both space and time to optimize beneficial effects (3). According to (4)  
42 intercropping promotes diversification and allows greater flexibility in adjusting to short- and  
43 long-term changes in the production and marketing situations, and also intercropping  
44 provides better weed control and reduces pest and disease incidence (5). Furthermore  
45 intercropping is a popular cropping system among small scale farmers in the tropics (6).  
46 Cereal/legume intercropping increased dry-matter production and grain yield more than their  
47 monocultures. The nitrogen transfer from legume to cereal increased the cropping system's  
48 yield and efficiency of nitrate uses. The taller cereal reduces biological nitrogen fixation and  
49 yield of the associated legume (7). According to (8), the competitive relationships between  
50 the non-legume and the legume affected the growth and yield of the leguminous crops in  
51 close proximity.

52

53 Sorghum (*Sorghum bicolor* L. Moench) belongs to the family *Gramineae*, including both  
54 wild and cultivated sorghum. Sorghum is the fifth important crop among the cereals in the  
55 world following rice, wheat, maize, and barley in total area planted and production [9].  
56 Sorghum is a principal cereal that forms an important staple diet throughout the semi-arid  
57 Asian and African regions (10). It is grown in regions receiving 300–1200 mm rainfall and in  
58 soils of pH range 5.0–10.0 (11). Sorghum is grown for grain, forage, syrup, and sugar. In  
59 2004 the total production of sorghum in the world was 57924 thousand tones and in Sudan  
60 was 2600 thousand tones (12). The total consumption of sorghum closely follows the global  
61 patterns of output, since most of it is consumed in the countries where it is grown. This is  
62 characterized by what most of smallholder farmers living the South East Lowveld of  
63 Zimbabwe rely on. These farmers produce a number of crops ranging from vegetables, cash  
64 crops and mostly sorghum as a source of their livelihood for food security.

65

66 Production of this cereal crop is reined by stem borer damage problems which salvage the  
67 crop to a mere fraction of the potential yield. *Chilopartellus* (Swinhoe) remains one of the  
68 economically important sorghum and maize stem borers in the dry areas of the south eastern  
69 Lowveld and East Africa (13). Stem borer damage to sorghum plants results primarily from  
70 leaf-feeding and stem tunnelling activities of the larvae. Characteristic leaf lesions and  
71 scarification caused by first and second larval instars of *C. partellus* are the first indications  
72 of infestation under natural field conditions (14). Yield reduction occurs as a result of leaf-  
73 feeding, dead hearts, stem tunnelling, direct damage to grain and increased susceptibility of  
74 attacked plants to stalk rots and lodging. Control of stem borers includes the use of chemical  
75 insecticides, host-plant resistance, cultural practices and biological control. This problem is  
76 aggravated by poor rains which are experienced in the area.

78 These resource marginalized farmers lack capital to purchase modern chemicals that are sold  
79 in most retail outlets/shops, to effectively fight the problem. Escalating costs and  
80 unavailability of chemicals (at times) has forced many smallholder farmers to use cheaper  
81 means such as intercropping. Surveys carried out in some parts of South Eastern Lowveld  
82 districts have revealed that most smallholder farmers have resorted to the use of different  
83 intercropping systems to alleviate the stem borer attacks on their crops. Whilst some of the  
84 farmers say that some of these control measures are effective, others have reservations on  
85 their efficacy. They only resort to these indigenous systems because they cannot afford or  
86 access the recommended pesticides. Adoption of effective intercrop practices for natural  
87 regulation of insect pests including stem borers remains crucial (15) (16), especially by these  
88 resource-poor farmers that lack the capacity of input-intensive plant protection measures.  
89 Groundnut (*Arachis hypogea*) is a short-duration legume crop grown by farmers in the  
90 savannah regions of Africa, and is readily intercropped with other medium duration crops  
91 such as pearl millet (17).

92 The objective of this trial was to evaluate the efficacy of different cultural indigenous legume  
93 intercropping knowledge systems for the control of stem borer in sorghum which is  
94 commonly grown in the marginalized low lying areas.

## 95 1.2 Methods and materials

### 97 1.2.1 Study site

The study was carried out at Chiredzi Research Station ( $21^{\circ}01'S$ ,  $31^{\circ}33'E$  429 m above sea level) located in the South Eastern Lowveld (Agro-ecological Region V) of Zimbabwe. It experiences temperatures ranging from  $29 - 39^{\circ}C$  and can reach up to  $42^{\circ}C$  and receive rainfall totals of 450-650mm year round and are common during the summer months and these are the favourable temperature requirements for stem borer multiplication. Triangle PE1 series such as shallow sandy clay soils dominate. The low latitude of 365-457 m above sea level is an effective safeguard against frost in all but the extreme circumstances. Minimum temperatures tend to run low in winter and frost can occur in low lying areas.

### 1.2.2 Experimental procedure and treatments

The trial was implemented at Chiredzi Research Station in the rain fed area. Treatment combinations of sorghum/cowpea, sorghum/groundnut, sorghum/bambara, sorghum/pigeon pea and sole crops of groundnut, cowpea, bambara and pigeon pea, replicated three times were laid on furrows spaced 0.90m wide in randomized complete blocks. Treatment controls comprised of sole sorghum plots where chemicals are applied and where no chemical was applied. Plantings were done with the first effective rains received in the area for the three seasons. Plot sizes are 8m long and 7 rows/ridges spaced at 0.9 m. Three to five seeds of sorghum variety SV4 are sown per hill and later thinned to one, with  $20 \times 90$  cm spacing between plants and between rows, respectively. Commercially available phosphate fertilizer at a rate of 150 kg/ha (18%  $P_2O_5$ ) is applied at planting and nitrogen at the rate of 34.5 kg/ha as ammonium nitrate applied at three weeks after crop emergence. Ten sorghum plants are randomly selected weekly from day of nitrogen fertilizer application until harvesting. Data on plant height, leaf damage, stem tunneling length, borer density and grain yield at harvest were recorded. Foliar damage ratings used for analysis were assessed at 8 weeks after emergence. Leaf damage based on a nine point visual rating scale (1, slightly visible damage; 9 severe foliar damage) is used using standardized chilo leaf damage scoring system (18) and (19). Plants are dissected and length of tunnel resulting from larval feeding recorded. Stem tunnel are expressed as the total length of stem tunneled as a percentage of plant height. Number of pupae, larvae and pupal cases from dissected plants are also recorded. Data for the three seasons was combined and subjected to GENSTAT version 14. Means were separated at  $\alpha=0.05$  by least squared differences (LSD) (20). Where percentages are used, the data is transformed using arc-sine transformation after adding 0.5 to each value. The data pertaining to the numbers of stem borers is transformed after adding 0.5 to each value by square-root transformation.

### 1.3 RESULTS


#### 1.3.1. Effect of stem borer damage on different cropping system of sorghum

Cropping system	Stem tunnel Length (m/plant)	Sorghum yield	Leaf severity score	Stover yield (kg)/plot	Larvae/plant
s/groundnut	50.4 <sup>a</sup>	5.18 <sup>b</sup>	2	18.67 <sup>b</sup>	1.469 <sup>a</sup>
s/cowpea	46.9 <sup>a</sup>	2.52 <sup>a</sup>	1.67	9.670 <sup>a</sup>	1.344 <sup>a</sup>
s/bambara	52.6 <sup>a</sup>	4.96 <sup>b</sup>	1.67	18.00 <sup>b</sup>	1.500 <sup>a</sup>
s/pigeon pea	48.8 <sup>a</sup>	4.33 <sup>b</sup>	1.67	14.17 <sup>b</sup>	1.431 <sup>a</sup>
s/sole (no sprays)	59.6 <sup>a</sup>	4.46 <sup>b</sup>	2	17.33 <sup>b</sup>	1.703 <sup>a</sup>
s/sole (sprayed)	44.3 <sup>a</sup>	5.65 <sup>b</sup>	1.67	23.67 <sup>c</sup>	1.153 <sup>a</sup>
<b>Grand mean</b>	<b>50.43</b>	<b>4.52</b>	<b>1.78</b>	<b>16.92</b>	<b>1.43</b>
<b>Lsd</b>	<b>17.94</b>	<b>1.721</b>	<b>1.135</b>	<b>4.403</b>	<b>0.6015</b>
<b>Cv %</b>	<b>24.1</b>	<b>15</b>	<b>35.1</b>	<b>10.3</b>	<b>28.6</b>
<b>Se</b>	<b>12.57</b>	<b>0.692</b>	<b>0.624</b>	<b>1.75</b>	<b>0.4215</b>
<b>pValue</b>	<b>0.537</b>	<b>0.030</b>	<b>0.942</b>	<b>&lt;.001</b>	<b>0.801</b>

**Table 1:** Effect of stem borer damage on different cropping systems of sorghum

#### 1.3.2 Effect of stem borer damage on sorghum stem on different cropping systems


No significant results on stem tunnel length at  $p < 0.05$  was noted among combinations. The sprayed control and the sorghum/cowpea combinations recorded shorter tunnel lengths than all other treatments. The unsprayed control record a longer tunnel length of 0.59 m.



**Fig 1: Effect of stem borer damage on sorghum stem on different cropping systems**


#### 1.3.4 Effect of stem borer damage on sorghum stover and grain yield among cropping systems

Significant lower mean stover and grain yield at ( $p < 0.05$ ) was observed (Fig 2 and 3) in plots where sorghum was combined with cowpea.



**Fig 2: Effect of stem borer damage on sorghum stover yield among different cropping systems**

Sorghum intercropped with groundnut, bambara and pigeon pea showed recorded almost the same yield of around 3.36 tonnes ha<sup>-1</sup>. The sprayed sole sorghum control out yielded all other treatments on both stover and grain yield.



**Fig 3: Effect of stem borer damage on sorghum grain yield among cropping systems**

Significant lower mean yield of 0.5 tonnes ha<sup>-1</sup>(Fig 3) ( $p<0.05$ )was recorded in plots where sorghum was intercropped with cowpea. Other combinations recorded the same grain yield of around 0.956 tonnes ha<sup>-1</sup>. No significant yield difference was recorded on the two sorghum sole plots.

### 1.3.4 Effect of sorghum legume intercropping on incidences of other pests among cropping systems

#### 1.3.4.1 Effect of legume intercrops on pest incidences

A significant higher aphid score of 1.14 ( $p<0.05$ ) was recorded among treatments (Table 2). The high mean aphid score was recorded in plots where sorghum was intercropped with groundnut and in plots where there was a cowpea sole crop. A lower scores was recorded in the cowpea and bambara combinations. In these respective treatments, higher ant populations were also recorded among treatments. Ladybird populations were observed in treatments where sorghum was combined with cowpea, pigeon pea as well in cowpea sole crops

<b>Cropping system</b>	<b>Ant</b>	<b>Aphid</b>	<b>Ladybird</b>	<b>Grasshopper</b>
s/groundnut	1.194 <sup>b</sup>	1.1454 <sup>c</sup>	0	0 <sup>a</sup>
s/cowpea	1.213 <sup>b</sup>	0.2357 <sup>ab</sup>	0.71	0.236 <sup>ab</sup>
s/bambara	0.408 <sup>ab</sup>	0.3333 <sup>ab</sup>	0	0 <sup>a</sup>
s/pigeon pea	0.236 <sup>ab</sup>	0.9024 <sup>bc</sup>	0.71	0 <sup>a</sup>
Sorghum sole (no sprays)	0.742 <sup>ab</sup>	0.7416 <sup>abc</sup>	0.24	0 <sup>a</sup>
Sorghum sole (sprayed)	0.408 <sup>ab</sup>	0.5690 <sup>abc</sup>	0.24	0 <sup>a</sup>
Groundnut sole	0.333 <sup>ab</sup>	0 <sup>a</sup>	0.41	0 <sup>a</sup>
Cowpea (sole)	1.040 <sup>b</sup>	0 <sup>a</sup>	0.86	0.569 <sup>b</sup>
Bambara (sole)	0.236 <sup>ab</sup>	0 <sup>a</sup>	0	0.236 <sup>ab</sup>
Pigeon pea (sole)	0 <sup>a</sup>	0.2357 <sup>ab</sup>	0.47	0.236 <sup>ab</sup>
<b>Grand mean</b>	<b>0.581</b>	<b>0.413</b>	<b>0.36</b>	<b>0.128</b>
<b>Lsd</b>	<b>0.8653</b>	<b>0.6697</b>	<b>1.113</b>	<b>0.467</b>
<b>Cv %</b>	<b>86.8</b>	<b>94.5</b>	<b>178.9</b>	<b>213.4</b>
<b>Se</b>	<b>0.5044</b>	<b>0.3904</b>	<b>0.649</b>	<b>0.2723</b>
<b>pValue</b>	<b>0.070</b>	<b>0.018</b>	<b>0.675</b>	<b>0.231</b>

**Table 2:** Other pest incidences among cropping systems

#### 1.3.4.2 Incremental benefit of intercropping among cropping systems.

An incremental benefit (Table 3) of between 9 -10 % were recorded in plots where sorghum was intercropped with groundnut and cowpea and pigeon pea. Over the seasons, no benefits were recorded in sorghum bambara combinations.

<b>Cropping System</b>	<b>Sorghum yield</b>	<b>Cowpea yield</b>	<b>g/nut yield</b>	<b>p/pea yield</b>	<b>Cow pea LER</b>	<b>G/Nut LER</b>	<b>P/Pea LER</b>	<b>Total</b>
S/groundnut	0.83		0.87			1.0		1.0
S/cowpea	0.97	1.66			1.0			1.0
S/bambara	1.07							0
S/pigeon pea	1.25			0.65			1.0	1.0
Sole sorghum (no spray)	2.02							1.0
Sole sorghum (sprayed)	2.48							1
Groundnut			1.75			2.02		1
Cowpea		2.79			1.9			1.0
Bambara								0
Pigeon pea				1.33			1	1

**Table 3:** Incremental benefit of sorghum legume intercropping among cropping systems

## 1.4 DISCUSSION



The effect of sorghum legume and insecticide applications in reducing stem borer activity and the resultant stem tunnelling in the level of crop damages by this stem borer was reflected in an appreciable non-significant relationships observed on stem tunneling (Table 1). This is in line with the findings of (21) who found out that green parts of legumes have dense covers of glandular hairs which are thought to exude a very acidic liquid. This, with a ph. of approximately 1.3 and a high content of malic acid produce a rancid smell and thus is thought to be a factor that repel stem borer or limit its ability to continue with stem tunnelling activities. Significant sorghum lower yield ( $p=0.030$ ) was recorded on plots where sorghum was combined with cowpea. This was also corroborated by (22), (23) who concluded that cowpea outcompete sorghum for water, light and nutrients (at few weeks after thinning) may result in lower sorghum yields and higher cowpea yield. Higher stover yield were recorded on sorghum/groundnut and sorghum/bambara combinations. This results in sorghum accumulating a lot of biomass during the growing season. This is in line with the work by (24), and (25) who found out that sorghum out compete groundnut and bambara in intercropping systems and the reverse for cowpea.

No statistical significant ( $p < 0.05$ ) on pest incidences was recorded among treatments. This might be due to diversity in the crop field that might have a profound effect on colonisation by insects (26), (27). It is also in line with (28) and (29) who found intercropping, an important cultural practice in crop pest management, primarily involves increasing the plant diversity of a given agro-ecosystem to aid reducing insect pest populations, and consequently, their attack. Populations of aphid predators (Table 2) such as ladybird were recorded among treatments. This goes in line with the findings of Dobson and Russell (unknown sources) who carried out an analysis of natural enemy impact on *chilo partellus* populations in 2005. Studies in Kenya by (30) also indicated that *Anthocorid* bugs (mainly *Orius* spp.) and ants (*Pheidole* spp., *Myrmecaria* spp. and *Camponotus* spp.) play the most important natural regulatory role on *chilo partellus*. This was also further explained by (31) who found out that intercrops facilitated the natural proliferation of predators.

Pest reduction in intercropping systems could be due to the “natural enemy effect” (32). Some plant combinations, for instance, with non-hosts lower the spread of pests within crops (33), (19). Non-host plants in such mixtures may emit chemicals or odours that adversely affect the pests, thereby conferring some level of protection to the host plant (34), (35). This

might be due to green parts of legume (24) (27). Studies indicate that crop diversification through intercropping, such as cereals with legumes, is effective in reducing insect pest damage (36), (37). Even plant diseases are believed to be less in intercropped agroecosystems due to increased crop diversity than those in sole crops (38), (39). Also, the intercropping of groundnut with pearl millet (*Pennisetum glaucum* L.) has particularly been found to increase the population of *Goniozus* sp., a parasitoid species that effectively manages leaf miner pest populations in ground nut (*Arachis hypogaea* L.) (40).

Results that were recorded on leaf severity, stem tunnelling and the number of larvae per plant were not significant among treatments. These are yield components and thus yield reduction due to stem borers occur as a result of leaf feeding, stem tunnelling, direct damage to cereal grain (41), (42). However, depending on the season and nutritional status of plant, crop yield reduction by stem borer feeding and tunnelling activities in Africa can fall between 10– 100% (28), (29). The unsprayed control recorded the longest tunnel length of 0.59 m while in intercrops it ranged from 0.3-0.5m. This is in line with (29) who found out that three to eight times more stems tunnelling and one to three times more cob damage were also recorded in monocropped maize with high stem borer larval densities (21– 48%) and yield loss (1.8 - 3.0 times greater) than in the intercropped counterparts. In contrast however, (29), in West Africa, found a considerably reduced amount of noctuid eggs laid by *Sesamia calamistis* Hampson and *Busseola fusca* Fuller due to reduced host found by the ovipositing adult moths in maize intercropped with grain legumes or cassava than those in the monocrop.

## 1.5 CONCLUSION

Intercropping, particularly with ground nut, pigeon pea and cowpea seems to encourage less stem borer infestation and abundance in sorghum, whilst additionally support high stover and grain yield. As such, the cultural practice is greatly encouraged over mono-cropping for stem borer pest management in sorghum grown in the south eastern Lowveld. Being an uncomplicated method of control and not capital-intensive, the practice should be readily adopted especially by resource-poor-farmers.

## 1.6 RECOMMENDATION

Farmers in the South Eastern Lowveld should intercrop using legumes in sorghum since intercrops have an incremental benefit. Legumes also provide nutrition to the soils as well as a source of relish. Legumes repel or attract other pests such as predators of aphid which is a pest of economic importance in drought stricken crops due to its sap sucking feeding habit.

## REFERENCES

1. S. Bhupinder S, Kalidindi U, Singh B, and Usha K 2003), “Nodulation and symbiotic nitrogen fixation of cowpea genotypes as affected by fertilizer nitrogen,” *Journal of Plant Nutrition*, vol. 26, no. 2, pp. 463–473. View at Publisher. View at Google Scholar · View at Scopus
2. Willey, R.W, (1990) “Resource use in intercropping systems,” *Agricultural Water Management*, vol. 17, no. 1–3, pp. 215–231. View at Google Scholar · View at Scopus
3. Potts M.J, (1990), “Influence of intercropping in warm climates on pests and diseases of potato, with special reference to their control,” *Field Crops Research*, vol. 25, no. 1-2, pp. 133–144, 1990. View at Google Scholar · View at Scopus
4. Campbel C.W (1990), “Techniques for producing export—quality, tropical horticultural crops,” *Horticultural Science*, vol. 25, pp. 31–33, 1990. View at Google Scholar
5. Finney D.J (1990), “Intercropping experiments, statistical analysis, and agricultural practice,” *Experimental Agriculture*, vol. 26, no. 1, pp. 73–81. View at Google Scholar · View at Scopus
6. Vandermeer J, (1989) *The Ecology of Intercropping*, Cambridge University Press.
7. K. Fujita, K. G. Ofosu-Budu, and S. Ogata, “Biological nitrogen fixation in mixed legume-cereal cropping systems,” *Plant and Soil*, vol. 141, no. 1-2, pp. 155–175, 1992. View at Publisher · View at Google Scholar · View at Scopus
8. R. Sangakkara, “Growth, yield and nodule activity of mungbean intercropped with maize and cassava,” *Journal of the Science of Food and Agriculture*, vol. 66, no. 3, pp. 417–421, 1994. View at Google Scholar · View at Scopus
9. I. C. Onwueme and I. D. Sinha, *Field Crop Production in Tropical Africa*, CTA, The Netherlands, 1993.

- 287 10. M. M. Ahmed, J. H. Sanders, and W. T. Nell, "New sorghum and millet cultivar  
288 introduction in sub-saharan Africa: impacts and research agenda," *Agricultural*  
289 *Systems*, vol. 64, no. 1, pp. 55–65, 2000. View at Publisher · View at Google Scholar  
290 · View at Scopus
- 291 11. N. Seetharama, S. Singh, and B. V. S. Reddy, "Strategies for improving rabi sorghum  
292 productivity," *Proceedings of the Indian National Science Academy*, vol. 56, no. 5-6,  
293 pp. 455–467, 1990. View at Google Scholar
- 294 12. FAO/WFP, "Special report FAO/WFP crop and food supply assessment mission to  
295 Sudan," 2007.
- 296 13. Warui, C.M. & Kuria, J.N. 1983. Population incidence and the control of maize stalk  
297 borers *Chilopartellus* (Swinhoe), *C. orichalcociliellus* Strand, and  
298 *Sesamia calamistis* Hamp in Coast Province, Kenya. *Insect Science and its Application*  
299 4: 11–18.
- 300 14. Sithole, S.Z. 1990. Status and control of *Chilopartellus* (Swinhoe) (Lepidoptera:  
301 Pyralidae) in southern Africa. *Insect Science and its Application* 11: 481–488.
- 302 15. Rao, M.R. (1986) Cereals in Multiple Cropping. In: Francis, C.A., Ed., *Multiple*  
303 *Cropping Systems*, Macmillan Publishing Company, New York, 34-39.
- 304 16. Verma, A.N. and Singh, S.I. (1989) Cultural Control of Sorghum Stem Borers. In:  
305 ICRISAT, *International Workshop on Sorghum Stem Borers*, ICRISAT Centre,  
306 Patancheru, 81-87.
- 307 17. Steiner, K.G. (1984) *Intercropping in Tropical Small Holder Agriculture with Special*  
308 *Reference to West Africa*. GTZ, Eschborn.
- 309 18. Dissmond, A. & Weltzien, H.C. 1986. Influence of sorghum and cow pea  
310 intercropping on plant pests in a semi-arid area of Kenya.  
311 *Mededelingen Faculteit Landbouwwetenschappen Rijksuniversiteit Gent* 51:1147–1155
- 312 19. ICRISAT 1990. Annual Report. International Crops Research Institute for the Semi-  
313 Arid Tropics (ICRISAT), Patancheru, Andhra Pradesh, India.
- 314 20. SAS INSTITUTE (1987). *SAS/STAT guide for personal computers*, version 6th  
315 edition. SAS Institute Inc., Cary, NC
- 316 21. Rembold, H and Winters, E. (1982) The Chemist's role in host plant resistance  
317 studies. In: *Proceedings of International Workshop on Heliothis Management* 15-20  
318 Nov 1981, ICRISAT Center, Patancheru, API, India. Pp 241-250
- 319 22. *Nitrogen Fixation in Tropical Cropping Systems*, Ken E. Giller, CABI Publishing,  
320 2001

23. Handbook for Innovative Maize-Legume Intercropping. 20 pp. SACRED Africa, Bungoma, Kenya. May 2002
24. Sustainable agriculture and intercropping – [www.leisa.info](http://www.leisa.info)
25. ATTRA Intercropping Principles and Practices - [www.attra.org/attra-pub/intercrop.html](http://www.attra.org/attra-pub/intercrop.html)
26. Risch SJ. Intercropping as cultural pest control: Prospects and limitations. *Environmental Management*. 2005;7(1):9–14.
27. Risch SJ, Andow D, Altieri MA. Agro—ecosystem diversity and pest control: data, tentative conclusions and new research directions. *Environmental Entomology*. 1983; 12:625–629.
28. J Ndemah, R. and Schulthess, F. (2002) Yield of Maize in Relation to Natural Field Infestations and Damage by Lepidopterous Borers in the Forest and Forest/Savanna Transition Zones of Cameroon. *Insect Science and Its Application*, 22, 183-193.
29. Chabi-Olaye, A., Nolte, C., Schulthess, F. and Borgemeister, C. (2005) Effects of Grain Legumes and Cover Crops on Maize Yield and Plant Damage by *Busseolafusca*(Fuller) (Lepidoptera: Noctuidae) in the Humid Forest of Southern Cameroon. *Agriculture, Ecosystem and Environment*, 65, 73-83.
30. van den Berg, H., Cock, M.J.W., Oduor, G.I. & Onsongo, E.K. (1993) Incidence of *Helicoverpaarmigera* (Lepidoptera: Noctuidae) and its natural enemies on smallholder crops in Kenya. *Bulletin of Entomological Research*, 83, 321-328.
31. Srinivasa Rao M, Dharma Reddy K, Singh TVK. Impact of intercropping on the incidence of *Marucavitrata* Geyer and *Helicoverpaarmigera* Hubner and their predators on Pigeonpea during rainy and post rainy seasons. *Shashpa*. 2004;11:61–70.
32. Baliddawa, C.W. (1985) Plant species diversity and crop pest control — an analytical review. *Insect Science and Its Application*. 1985; 6(4):479–487.
33. Gautan, R.C. and Kaushik, S. (1980) Studies on Planting Dates and Intercropping of Pearl Millet with Grain Legumes. *Indian Journal of Agronomy*, 25, 441-446.
34. Singh, K.A., Ahlawat, I.P.S. and Mahiendrapal, H. (1986) Cropping Systems Research 1. Concept, Needs and Directions. Proceedings of the National Symposium on Cropping Systems, New Delhi, Indian Society of Agronomy, Indian Agricultural Research Institute.
35. Reddy, S.R. (2012) Agronomy of Field Crops. Kalyani Publishers, New Delhi, 245-264.

- 354 36. Nwanze, K.F. (1989) Insect Pests of Pearl Millet in West Africa. *Acigoniaignefusalis*  
355 (Pyralidae: Lepidoptera); Distribution, Population Dynamics and Assessment of Crop  
356 Damage. Tropical Pest Management, 35, 137-142.  
357 <http://dx.doi.org/10.1080/09670878909371342>
- 358 37. Mailafiya, D.M. and Degri, M.M. (2012) Stem Borers' Species Composition,  
359 Abundance and Infestation on Maize and Millet in Maiduguri, Nigeria. Archives of  
360 Phytopathology and Plant Protection, 45, 1286-1291.
- 361 38. Altieri, M.A. and Liebman, M.Z. (1986) Insect, Weed and Plant Disease Management  
362 in Multiple Cropping Systems. In: Francis, C.A., Ed., Multiple Cropping Systems,  
363 Macmillan, New York, 183-218.
- 364 39. Gomez, A.A. (1990) Farming System Research Approach to Identifying Farmers'  
365 Production Problems. Farming System Research Approach, 8, 63-70.
- 366 40. Dhaliwal, G.S. and Arora, R. (1996) Principles of Insect Pest Management. National  
367 Agricultural Technology Information Centre, Ludhana.
- 368 41. Bosque-Pérez, N.A. and Marek, J.H. (1991) Effect of the Stem Borer  
369 *Eldanasaccharina* Walker (Lepidoptera: Pyralidae) on the Yield of Maize. Bulletin of  
370 Entomological Research, 81, 243-247.
- 371 42. Sétamou, M., Schulthess, F., Poehling, H.M. and Brgemeister, C. (2000) Monitoring  
372 and Modeling of Field Infestation and Damage by the Maize Ear Borer  
373 *Mussidianigrivenella* (Lepidoptera: Pyralidae) in Benin, West Africa. Journal of  
374 Economic Entomology, 93, 650- 657.