

The effect of traditional shade-drying method on seed germination and vigour of two varieties of tomato

ABSTRACT

The effect of traditional shade-drying method on the seed germination and vigour of two varieties of tomato was investigated. The experiment was conducted at the seed testing laboratory of The National Centre for Genetic Resources and Biotechnology (NACGRAB), Ibadan, Nigeria in July 2015. The seed samples were dried for one week in the drying chambers and thereafter evaluated for germination and vigour tests. The experiment was carried out in a completely randomized design with three replications, in 2 x 2 factorial scheme. Two varieties of tomato (Ibadan local and Alausa) and two drying methods: traditional (shade-drying with electric fan at temperature between 23.5 to 32.3°C) and mechanical (seed dryer at 35°C) were evaluated. The germination percentage of Ibadan local variety was significantly higher (87.2%) in comparison to the Alausa (79.0%). Mechanical drying at 35°C gave the higher germination percentage (94.16%), while seeds dried traditionally gave the germination of 72.0%. Moreover, the effect of drying method was not significant on germination index of tomato seeds, suggesting that seeds dried using both methods may not exhibit differential performance when subjected to unfavorable environmental conditions either on the field or during storage. The study indicates that drying using seed dryer at controlled drying temperature would enhance germination of tomato seeds better than traditional shade drying method.

Keywords: Tomato; drying; germination percentage; germination index.

14 1. INTRODUCTION

15 Tomato (*Lycopersicon lycopersicum* Mill.) is one of the most important vegetables worldwide
16 belonging to the Solanaceae family. It is cultivated for its fruits which contribute to a healthy,
17 well-balanced diet in human nutrition. Tomato fruits are rich in minerals, vitamins, essential
18 amino acids, sugars and dietary fibres. The fruits can be consumed fresh in salads or
19 cooked in sauces, soup and meat or fish dishes. They can be processed into purées, juices
20 and ketchup. Canning and drying are also economically important means of processing
21 tomato fruits.

22 Tomato is propagated by seed and the use of high quality seeds is very important for
23 successful production of this crop as the establishment of an adequate plant population in
24 the field is necessary to achieve high productivity [1]. Drying operation is a critical step in the
25 post-harvest processing of tomato seeds however, seed quality can be reduced during
26 drying due to injury caused by unfavorable drying conditions, although the causes and
27 impairment mechanisms are poorly understood [2]. The air temperature and relative humidity
28 of the environments during drying process are the two major factors that influence the
29 germination characteristics of seeds hence methods of drying must be carefully selected in
30 order to avoid injury caused by unfavorable drying conditions. There are several factors to
31 be considered when choosing seed drying method to be used which include seed volume
32 effectively harvested, harvest speed, drying time, energy consumption and end purpose of
33 seeds. There are different methods of seed drying, such as shade and sun drying, vacuum
34 drying, freeze drying and refrigeration drying with low relative humidity [3]. Other
35 recommended methods for safe drying of seeds include seed drying chambers, seed dryers
36 and controlled conditions [4] however, in many developing countries, such drying facilities
37 are limited due to the high cost of establishing, running and maintaining such facilities.

38 The fundamental objective of seed testing is to establish the quality level of seed. Seed
39 vigour and germination tests aimed at differentiating low and high quality seeds from each
40 other. The Association of Official Seed Analysts [5] defined seed germination as 'the
41 emergence and development from the seed embryo of those essential structures which, for
42 the kind of seed in question, are indicative of the ability to produce a normal plant under
43 favourable conditions'. Germination capacity therefore forms a crucial aspect of seed quality
44 hence the germination tests are used worldwide to determine the maximum germination
45 potential of a seed batch under optimum conditions. The International Seed Testing
46 Association (ISTA) [6] defined seed vigour as "the sum total of those properties of the seed
47 that determine the level of activity and performance during germination and seedling
48 emergence". The aspect of performance associated with seed vigour include (i) rate and
49 uniformity of seed germination and seedling growth, (ii) field performance, including the
50 extent, rate and uniformity of seedling emergence, and (iii) performance after storage and
51 transport particularly the retention of germination capacity. The concept of seed vigour
52 implies that two seed lots having similar germination level may perform differently due to
53 differences in vigour potential when subjected to poor field conditions. Speed of emergence
54 of seedlings is one of the oldest seed vigour concepts and vigorous seeds have been shown
55 to germinate rapidly. Speed of germination has been measured by various techniques and
56 given many different names such as: emergence rate index, germination rate, germination
57 index and speed of germination. The tests have important advantages. They are
58 inexpensive, rapid, require no specialized equipment, and most importantly do not
59 necessitate additional technical training.

60 Tomato fruits harvested at physiological maturity usually contain high moisture of 60% to
61 70% hence sooner the seed is extracted, cleaned and dried better will be the quality. In
62 addition, the rate at which moisture evaporates from the seed surface also determines the
63 quality of tomato seeds. If moisture evaporates too rapidly, the resulting moisture stress can

damage the embryo. If the moisture evaporates too slowly, it may favour invasion of pathogen therefore seeds should be dried carefully to arrest stress damage.

Investigations have been conducted on effective methods of drying seeds of some solanaceous crops such as chilli [7, 8] and tomato [9]. However, most of the studies laid much emphasis on traditional methods of drying probably due to the high cost of establishing, running and maintaining mechanical methods of drying such as drying chambers or seed dryers. In order to develop an efficient drying method without compromising seed health and quality, the National Centre for Genetic Resources and Biotechnology (NACGRAB) located in Ibadan, Nigeria who has institutional mandate for genetic resources conservation and utilization in Nigeria recently procured seed dryer for drying of seeds especially for small seeded crops like tomato, however, information on comparative performance between traditional method (shade-drying with electric fan at temperature between 23.5 to 32.3°C) which was in practice at NACGRAB and mechanical method (seed dryer at 35°C) on seed quality of tomato is not available. The aim of this study therefore was to compare these two drying methods with a view to identify the best method suitable for a successful and cost effective production of biologically viable tomato seeds.

2. MATERIAL AND METHODS

2.1 Plant Materials and Seed Production

The seeds of two varieties of tomato: Ibadan local and Alausa, which are popular among the farmers in the South West Nigeria were sourced from the seed gene bank of the National Centre for Genetic Resources and Biotechnology (NACGRAB), Ibadan. Seed production was carried out at the experimental field of the NACGRAB during the growing seasons of 2015.

2.2 Seed Processing

Fruits of the two varieties of tomato were harvested at physiological maturity stage and seeds were extracted directly after harvesting. The extraction was done by hand to minimize mechanical damage. The seeds of each variety were partitioned into two equal parts and samples from each variety were subjected to two drying methods: traditional (shade-drying with electric fan at temperature between 23.5 to 32.3°C) and Mechanical (seed dryer at 35°C). The seed samples were dried for one week in the drying chambers and thereafter evaluated for germination and vigour tests.

2.3 Experimental Design

The experiment was conducted at the seed testing laboratory of NACGRAB in July 2015. The experiment was carried out in a completely randomized design with three replications, in 2 x 2 factorial scheme. Two varieties of tomato and two drying methods: traditional (shade-drying with electric fan at temperature between 23.5 to 32.3°C) and mechanical (seed dryer at 35°C) were evaluated.

2.4 Standard Germination and Vigour Tests

One hundred seeds of each variety were drawn and evaluated for standard germination test in three replications. The test was assayed by placing the seeds in germination plastic containers lined with four layers of tissue paper moistened with 15 ml of distilled water. The containers were covered and placed in a germinating chamber at 25 ± 2°C. The seeds were kept moist every day for seven days. Germination percentages were calculated by expressing the number of seedlings in a replicate that emerged 7 days after planting as a percentage of the number of seeds planted according to ISTA rules [10]. Germination Index (GI) was calculated by taking the germination counts at 5, 7 and 9 days after planting using the following formula:

$$GI = \frac{\text{No of germinated seed}}{\text{Days of first count}} + \dots + \frac{\text{No of germinated seed}}{\text{Days of final count}}$$

2.5 Data Analysis

Data on germination percentage were subjected to analysis of variance (ANOVA) using Statistical Analysis Software, SAS Version 9.1 [11]. Data on percentages do not conform to normal distribution, the germination data were therefore log transformed before subjecting them to the ANOVA. However, since ANOVA did not detect any significant difference between transformed and untransformed values, untransformed values are hereby presented. Pertinent means were separated by the use of the least significant difference (LSD) at 0.05 level of probability.

3. RESULTS AND DISCUSSION

3.1 Germination Performance of Tomato Seeds as Influenced by Variety and Drying Methods

Analysis of variance (ANOVA) revealed that effect of tomato variety was significant ($P < 0.05$) while effect drying methods (DRY) was highly significant ($P < 0.01$) on germination of tomato seeds (Table 1). The germination percentage of Ibadan local was significantly higher (87.17%) when compared with the germination percentage of Alausa (79.0%) (Table 2). This result emphasises the fact that genetic constitution of any seedlot is a major determinant of its quality. The result agrees with findings of some authors who reported variations among genotypes different species of crops. Tame and Elam [12] reported significant variation among three soybean varieties for germination after 360 days in storage. Similarly, Omar et al. [13] also reported significant variation among three cultivars of wheat for germination, viability (Electrical conductivity and Acidity %) and seedling vigor (radical length, plumule length and seedling dry weight) after storage at different periods.

Table 1. Mean squares from the analysis of variance for the germination test and germination index on tomato seeds at NACGRAB, Ibadan

Source of variation	DF	Germination (%)	Germination index (days)
Rep	2	73.58ns	0.00ns
Variety (VAR)	1	200.08*	0.01ns
Drying methods (DRY)	1	1474.08**	0.00ns
VAR x DRY	1	102.08ns	0.02ns
Error	6	27.91	0.00
Total	11	190.08	0.01
CV		6.36	1.56
Mean		83.08	4.1

DF - Degrees of freedom; CV - coefficient of variation; *, ** - significant at probability level of 0.05 and 0.01, respectively; ns - not significant

Similarly, effect of drying methods was significant on germination of tomato seeds. Mechanical drying method using the seed dryer at 35°C gave the higher germination

percentage (94.16%) while seeds dried traditionally under shade with electric fan gave the germination percentage of 72.0% (Table 2). This significantly lower in seed germination for traditional shade-drying using electric fan might be as a result slow evaporation of moisture from the seed surface coupled with fluctuation in temperature of the room conditions which may favour invasion of pathogen. Although, Gowda et al. [9] concluded that combined sun and shade drying resulted in the highest seed germination of tomato (94%) however from this study, mechanical drying method using the seed dryer at 35°C gave better germination percentage compared to traditional shade-drying using electric fan. This result corroborated with the findings of Ali et al. [14] who concluded that sorghum seeds dried with seed dryer gave the highest germination percentage compared to those dried using silica gel, or under shade and sun. They further reported that drying with silica gel and shade are good alternative methods of drying. Nevertheless, our results show that traditional shade-drying using electric fan could also serve as alternative or low cost method of drying tomato seeds. This is in line with the report of [8] who investigated the effect of seed drying methods namely, sun-drying, shade-drying and freeze drying on physiological quality of chilli seeds and concluded that shade-drying was significantly better than other methods of drying. Furthermore, our results show the effect of drying method was not significant on germination index of tomato seeds suggesting that tomato seeds dried using both methods may not exhibit differential performance when subjected to unfavorable environmental conditions either on the field or during storage.

Table 2. Effect of varieties and drying methods on seed germination of tomato at NACGRAB, Ibadan

Factors	Seed germination (%)	Germination index (days)
Variety		
Ibadan local	87.17a	4.13a
Alausa	79.00b	4.01a
LSD	7.46	0.09
Drying Methods		
Traditional	72.00b	4.11a
Mechanical	94.16a	4.08a
LSD	7.46	0.09

CONCLUSION

This study led to the conclusion that the drying method highly affects the seed germination of tomato. From this study, mechanical drying at 35°C would give better germination compared

to traditional shade-drying using electric fan drying. Moreover, the effect of drying method was not significant on germination index of tomato seeds suggesting that tomato seeds dried using both methods may not exhibit differential performance when subjected to unfavorable environmental conditions either on the field or during storage. Furthermore, the study indicated that traditional shade-drying using electric fan could serve as alternative method of drying tomato seeds

REFERENCES

1. Owolade OF, Fawole B, Osikanlu YOK. Fungi associated with maize seed discolouration and abnormalities in south-west Nigeria. Trop. Agric. Res. Extension. 2001; 3 (2) 103-105.
2. Cordova-Tellez L, Burris JS. Alignment of lipid bodies along the plasma membrane during the acquisition of desiccation tolerance in maize seed. Crop Sci. 2002; 42(6): 1982-1988.
3. Ellis RH, Roberts EH. The potential of ultra-dry storage of seeds for genetic conservation. University of Reading, UK. 1991; 43.
4. Franke LB, Miguel AP, Torres R, Rodrigo RL. Performance of different drying methods and their effects on the physiological quality of grain sorghum seeds (*S. bicolor* (L.) moench). Revista Brasileira de Sementes. 2008; 30(3):177-184.
5. Association of Official Seed Analysts. (2009). Rules for testing seeds. 2004. AOSA, Ithaca, NY.
6. ISTA (International Seed Testing Association) (1993) International rules for seed testing. *Seed Science Technology* 21 (Suppl.).
7. Nagaraja A., Basavaraja PK, Yogeesh H. Effect of drying methods on fruit and seed quality of chilli. South Indian Horticulture, 1998; 46(3/6): 262-265.
8. Chakradhar P, Laxmipriya S, Ananda CS and Saroj KM. Studying the effect of different fruit and seed drying methods on seed quality of two different genotypes of chilli (*Capsicum annuum* L.). International Journal of Science, Environment and Technology, 2016; 5(5):3262-3270.
9. Gowda SJ, Talukdar KC, Ramaiah H. Effect of drying methods on seed quality in tomato (*Lycopersicon lycopersicum*). Seed Research. 1990; 18(2):126-129.
10. ISTA. International Rules for Seed Testing, 2003. Zurich, Switzerland, ISTA. 2003.
11. SAS. SAS/STAT User's Guide version 6, 4th edition. SAS Institute, Cary, North Carolina, USA. 1990.

- 212 12. Tame VT, Elam Y. Effect of storage materials and environmental conditions on
213 germination of percentage of soybean (*Glycine max* (L.) Merr) seeds in Yola, Nigeria.
214 International Journal of Agricultural Sciences and Natural Resources. 2015; 2(4): 90-94.
- 215 13. Omar AM, Sorour FA, El-Sayed SA, Naywa ES. Effect of storage periods, cultivators,
216 environments and package materials on germination, viability and seedling vigour of
217 wheat grains. Journal of plant production, Monsura University, 2012; 3(b):1075-1087.
- 218 14. Ali ZB, Mohammad ED, Mustafa MA, El Tahir I. Effects of low cost drying methods on
219 seed quality of Sorghum bicolor (L.) Monech. African Journal of Plant Science. 2010;
220 4(9): 339-345,

221

222

223