

1 **Biology and Dispersal of the Watermelon Bug *Coridius viduatus* (F.)**
2 **(Heteroptera: Dinidoridae) on Different Cucurbit Crops, in North Darfur**
3 **State, Sudan.**
4

5
6
7 **ABSTRACT**

8 The watermelon bug, *Coridius viduatus* (F.) is a real threat to watermelon *Citrullus*
9 *lanatus* (Thunb.) in western Sudan, where over 80% of the population relies
10 economically on agriculture. In order to overcome this constraint, a study was carried
11 out at University of Alfashir, North Darfur State, to investigate biology, food
12 preference and dispersal of watermelon bug. A survey was conducted on season
13 (2013/2014) to determine the movement and dispersal of the watermelon bug in the
14 area around Alfashir. Biology of the bug was studied under laboratory conditions,
15 preoviposition, oviposition, incubation and post oviposition periods were calculated.
16 Food preference and non-preference by the bug to four watermelon varieties and two
17 cucurbit ones were also evaluated; a field experiment was conducted, a randomized
18 complete block design was used. The field survey results indicated that there was a
19 regular movement from plant shelters, mountain crevices and soil cracks to the field
20 crop and back again to aestivation sites. Results showed that the bugs preferred
21 improved watermelon varieties (Crimson, Sugar baby and Congo) to the local
22 watermelon variety (Saphinga), the different life cycle stages of the bug were
23 determined. The bug aestivation shelters were determined, local watermelon
24 varieties could be cultivated however further work should be done to improve their
25 productivity.

26 **Key words:** *Citrullus lanatus*, *Coridius viduatus* (F.) watermelon- bug, Cucurbitaceae,
27 dispersal, preference.

28
29 **1. INTRODUCTION**

30 The watermelon, *Citrullus lanatus* (Thunb.) is originated in Africa and has been cultivated
31 for more than 4000 years in the drier parts of the continent and throughout India and
32 parts of Asia. It was grown in an area 3.2 million ha and the total production was 89.1 million
33 tons in 2010 [1]. It is used as desert fruit and thirst quencher and in the very dry parts of
34 Africa, it is consumed by both man and his animals as a source for water. The roasted
35 seeds are popular as food in some areas, especially in west Africa and southern China
36 and they contain a semi-drying oil [2]. Watermelon is a warm season crop and can be
37 cultivated year round in the tropics, the most producing countries are China (70,000,000
38 tons), Turkey (4,044,144 tons), Iran (3,800,000 tons), Brazil (2,079,547 tons) and Egypt
39 (1,874,710 tons), worldwide production of watermelon was 95,211,432 tons in season 2012,
40 watermelon contain about 92% water and 6% sugar by weight, it is a source of vitamin C [3].

41 There are several insects causing damage on watermelon such as watermelon beetles,
42 aphids, flies, ants, etc. However, the most serious one is the Black watermelon bug *Coridius*

viduatus (F.) (Heteroptera:Dinidoridae).The genus *Coridius* includes this species is currently replacing the old genus *Aspongopus* [4], in the Sudan watermelon is considered to be an important economic food crop both for man and his animals and is widely grown in Kordofan, Darfur, Kassala, Khartoum, Blue Nile, and white Nile States [5][6].There exist, in the far northern parts of the Darfur Region, wild local cucurbits varieties including watermelon. The environmental conditions were suitable for watermelon bug to develop, hide and rest along the study area. Therefore the grown populations have invaded the traditional water melon growing areas in the sandy clay soils and emigrated to attack the plant grown in valleys. Heavily infested plants wilt and finally die. In this regard melon bug is the most important insect pest in western Sudan [7] it is also found in almost all African, Arabian countries, Iran and Turkey. Nymphs and adults are gregarious, those from one egg cluster feed close together, and transmit bacterial and viral diseases. On the other side, in the remote territories of Sudan, oil from the bug *Coridius viduatus* is extracted and used as sweet oil without any poisons effect. This oil corresponds in its main content with the most of animal oils. It has been investigated that watermelon bug oil is used as food and medicine in dermatological diseases due to its antibacterial effect [9]. Biofuel production from this insect was reported in this country as well. Hundred grams of watermelon fruits contains 92.6g moisture, 0.5g protein, 0.2g fat and 4.6g carbohydrates; whole watermelon seed contains moisture (4.94), fat (25.87), protein (18.96), ash (2.31), fiber (39.84) and carbohydrate (8.38) g [10].

the eggs laid in clusters in lower surface of the leaves, the dark brown eggs are laid in groups about 1cm across on the lower surface of the leaf or in long rows along stems or veins, The female lays about 300-500 eggs in masses of dark colour [5] [13] [14] [15][16]. Young nymphs of the melon bug range in colour from, pale-brown to reddish black depending on their age. The newly hatched nymphs are found near the eggs cluster, Nymphs of the last instars reassemble the adult in colour and appearance but without fully developed wings. They are often seen feeding on the host in great numbers [6].The incubation period is (7-8) days, hatchability is (95-100%), eggs hatch to give Nymphs, there are five nymphal instars with a period of (42-45) days to give adult bugs, there are two generations per year[13][17].

Heavy infestation generally occur during the winter season, the bugs concentrate for aestivation, after massing in shady, low lying places [18],The bug seriously reduces the crop vigour and productivity. Some studies were carried in Sudan talked biology, ecology, bio control and integrated management of the watermelon bug in different ecological zones and even under different cropping systems [19] but meager or absent studies about the movement or dispersal of the bug from host plants to the aestivation sites. The objectives of this study are to investigate the biology of the bug under laboratory conditions, food preference and dispersal of watermelon bug.

2. MATERIAL AND METHODS

2.1 Field Survey

A survey was conducted in selected secured Localities viz. Alfashir, Dar Alsalam, Kelimendo, Wada and some other rural areas around Al-Fashir city; the capital of North Darfur State Latitude 13° 37' 40.55" N, Longitude 25° 20' 57.70"E.The survey took place on winter, summer and autumn (October-2013, March-2014 and August-2014).The objectives of this survey were to determine the sheltering sites and the nature of the watermelon bug movement. These areas are characterized by poor dry Savannah and sandy soils predominate as well as scattered shrubs and trees where water melon is grown at wide scales for both, fruits and seeds. The numbers of the pest were calculated during summer season to identify the sheltering sites and the pest density.

2.2 Watermelon Bug Food Preference and Non- Preference on Different Watermelon and Cucurbit Varieties in North Darfur State:

An experiment was conducted at the demonstration farm of Al-Fashir University; the objective of this experiment was to study food preference and non-preference of melon bug. An area of 740 m² was divided into 18 plots (4x5m) each contain three mustabas one meter width and 3meters length, a completely randomized block design was used, 6 treatments (varieties) were replicated 3 times. Seeds of four watermelon varieties, (three are genetically improved, i.e. Crimeson, Congo, Sugar baby and one local watermelon Safinga) and tow pumpkin varieties (genetically improved Ananas, and local Harshayia) were sown in a randomized complete block design and replicated three times, in the first week of July, 2014 with the onset of rainfall. Cultural practices were followed as recommended, Observations were recorded on weekly bases intervals and six counts were obtained. Analysis of variance was done and the least significant difference Lsd at 0.05 was calculated.

2.3 Biological studies

2.3.1Life cycle:

After survey that carried out in the study areas around Al-Fashir in season 2013 – 2014, insects' stock culture has initially been established from eggs deposited by female watermelon bugs.

Mature adult melon bugs were collected by hands early in the mornings and late evenings during late November and early December 2013, from Sarafayia, Magdoob and Umgidaibo areas around Al-Fashir. The bugs were put in transportable wooden cage and carried to the main laboratory of the Faculty of Environmental Sciences and Natural Resources then transferred into a mass rearing box measures (20x25x30cm), the bottom side was covered with a hard wood, the 4sides were covered with fine mesh, the top side was covered with a movable door to facilitate insect inspection. Adults were daily fed with fresh food, small leaves, stems, growing shoots of water melon; laboratory reared progeny of the overwintering adults were remain alive and continue egg laying on the wooden cages floor on leave under sides and cage corners[10].

Eggs lay in clusters, rows or singly were left to incubate at room temperature (37±2.6°C). After eggs hatching the first nymphs transferred into plastic containers, sorted out and uniform ages kept in groups, each group was composed of 5nymph instars as one unit of similar ages and kept in separate cages, Ibrahim metallic cage, the number of cages were 10. The total number of the first instars amounted (50) nymphs and were fed with foods (watermelon leaves, stems, young growing shoots, young fruits), and replaced every 24 hours by fresh succulent food. However, after hatching, the first, second , third, fourth and fifth instar stages up to adult emergency were summed up in days. Ibrahim cage was designed by making use of local materials as described below.

2.3.1.1 Ibrahim cage description

It was designed by making use of some local material to study the life cycle of melon bug. It composed of empty cylindrical perfume container (Mahlabia), its height was 8cm and the diameter was 8.4cm with 4windows (Holes) their length was 5.9cm, the width was 4.6cm to allow smooth ventilation, the inner walls were lined all through with fine sieve/mesh to hinder escaping of nymphs and permit close watch over, as well as enabling the nymphs to cling onto during resting and food sucking periods and molting process. However, nymphs remain in tight contact with food most of the time. The bottom of cage was mulched with filter paper where food was placed and the top of the cage covered with

137 a piece of cloth made of cotton and fixed with a rubber band. It services as protective
138 cover and allows ventilation. This cage was used for bug groups rearing at arrange of 5
139 nymphs per cage [10].

140 3. RESULTS AND DISCUSSION

141 3.1 Field Survey

142 The results of the filed survey that took place on winter, summer and autumn (October-2013,
143 March-2014 and August-2014) in the study areas. It had been found that the most important
144 sheltering plants for the black watermelon bug *Coridius viduatus* were Remta (*Eragrostis*
145 *aspera*), Argassy (*Chrozophora brocchiana*), Eyear (*Momordica balsamina*), Dura (*Sorghum*
146 *bicolor*), Mrahabaeb (*Cymbopogon giganteus*), Gdgad (*Gloriosa superba*), Tontob (*Capparis*
147 *sepiaria*), Hashab (*Acacia Senegal*), Laout (*Acacia nubica*), Gedaem (*Grewia villosa*),
148 Khbaesh (*Haemanthus multiflorus*), Sider (*Ziziphus spina-cristi*) and Osher (*Calotropis*
149 *procera*). The black watermelon bugs were found hiding between the leaves of these plants;
150 in addition to mountain crevices and soil cracks. Adult melon bugs undergo the dry season in
151 aestivation. These shelters differ from that reported in Northern State, Sudan where the black
152 watermelon bug shelter on Sodom apple *Calotropis procera*, Neem *Azadirachta indica* and
153 some *Acacia* spp. But the main sheltering plant was date palm *Phoenix dactylifera* [8]
154 Watermelon bug *C. viduatus* seriously attacks the watermelon crops, in Northern Kordofan
155 and Darfur states. This crop is considered as most important one, since it is used as a source
156 for drinking water because of water shortage in that area during the summer months.

157 It was also noted that melon bugs prefer cucurbit crops specially the watermelon. Adults and
158 nymphs suck the juice of the main stems, leaves and the developing fruits, heavily infested
159 plants shed their leaves and young fruits. The bug attacked other crops like Dura *Sorghum*
160 *bicolor* and pearl millet *Pennisetum gluacum*; these crops act as a source of infestation.

161 During this survey many observations were recorded; before sunset the bugs moved to the
162 upper parts of the host plant for a while then returned to the lower parts at night. Numbers of
163 bugs were huge in the sheltering plants around the watermelon fields and reached up to 3000
164 bugs per tree. They were found at the lower part of trees, shrubs, and weeds (mentioned
165 above) in clusters.

166 The results showed that the bug was found in valley flows under Hills and traditional stores in
167 the villages. It could be argued that from these observations the black watermelon bug
168 *Coridius viduatus* passed the off season hiding under these plants, trees, shrubs in summer
169 and start to migrate to attack field crops sorghum and pearl millet during autumn the same
170 observation reported later [20]. Its population increases toward the end of the rainy season
171 and attack watermelon and other cucurbit crops in winter (October to March), then back to
172 aestivation sites in late March. Generally we can mention that there was a regular movement
173 started during early August from plant shelters, mountain crevices and soil cracks to the field
174 crops especially watermelon and back again to aestivation sites during summer.

175

176 3.2 Biological Studies

177 Studies on the life cycle of the black watermelon bug *Coridius viduatus* (Fabricius) were
 178 conducted in the Entomology laboratory of the Faculty of Environmental Sciences and Natural
 179 Resources, University of Alfashir, under room temperature in season (2013-2014). Eggs were
 180 found laid on the leaves, stem and growing shoots of watermelon and arranged in clusters.
 181 The results showed that, the black watermelon bug endures incomplete metamorphosis and
 182 passed through five nymph instars. This result was in the same line with many investigators
 183 [8] [15] [16]. The different developmental stages during the course of the bug life span were
 184 described as follows:

185 **3.2.1Pre oviposition period**

186 The mean preoviposition period of the mated female was 5.70 ± 2.53 day. This
 187 period ranged from 2 to 9 days (Table 1) [19] [8]

188 **Table 1.The number of eggs laid by a female, preoviposition, oviposition, post**
 189 **oviposition periods and life span of the black watermelon bug *Coridius***
 190 ***viduatus* in days reared under laboratory conditions.**

No.	Total number of eggs Laid/Female	Pre oviposition period	Oviposition Period	Post oviposition period	Life span of female	Life span of male
1	160	9	45	5	59	30
2	210	3	45	7	55	29
3	265	5	29	7	41	27
4	419	6	43	6	55	30
5	519	2	39	4	39	25
6	315	6	55	3	53	23
7	220	9	50	5	64	29
8	251	7	41	6	54	27
9	361	8	52	3	63	28
10	215	2	57	5	64	30
Mean	293.50	5.70	45.6	5.1	54.70	27.80
Range	160-519	2-9	29-57	3-7	39-64	23-30
SE±	105.16	2.53	8.34	1.37	8.36	2.23

191

192 **3.2.2 Oviposition period and fecundity:**

193 It was found that the egg is small in size, cylindrical in shape when newly laid it was
 194 green in colour and gradually changed to reddish-brown just before hatching. The
 195 eggs either laid singly in rows along the running stems of watermelon shoot or eggs
 196 lay in clusters glued to each other along the running stem forming a straight bead
 197 like structure, frequently eggs were laid on the lower surface of the leaves of
 198 watermelon. However, during the course of this study, few eggs were laid on the
 199 cloth pieces that covered the oviposition cage.

200 The mean oviposition period of watermelon bug reared under laboratory conditions
 201 was 45.60 ± 8.43 days ranged from 29 to 57 days. The Maximum number of eggs laid
 202 by female was 519 and minimum was 160 eggs. The mean of eggs laid/female was
 203 293.50 ± 105.16 (Table 1), this results are in accordance with [8][18] [18] and it differ
 204 [19]. The maximum number of eggs laid by a female was during the third week from
 205 the beginning of egg laying.

206 **3.2.2 Postoviposition period**

207 Post oviposition period varied from one female to another, the mean post oviposition period
 208 was 5.10 ± 1.37 days ranged from 3 to 7 days this result in the same line with other investigator
 209 [8]. The duration of total life span for female and male were 54.78 ± 8.36 and 27.80 ± 2.23
 210 respectively (Table1).

211 **3.2.3 Hatchability%:**

212 The mean hatchability was 92.50 ± 7.16 ,

213
$$Hatchability\% = \frac{emergednymphs}{totaleggslaid} \times 100$$
 (Table 2). This result coincides with the same that
 214 mentioned by [11] also indicates the high fecundity of the pest.

215 **Table 2. Incubation period in days and hatchability % of watermelon bug**

216 ***Coridius viduatus* reared under laboratory conditions.**

Serial	Egg	Hatching	No. of	No. of	incubation	Hatchability%
No.	laying	date	egg laid	egg	period	
	Date			hatched		
1	27-11	7-12	20	18	10	90
2-	27-11	7-12	20	19	10	95
3-	27-11	7-12	20	20	10	100
4-	27-11	7-12	20	16	10	80

5-	27-11	10-12	20	20	13	100
6-	28-11	10-12	20	20	12	100
7-	29-11	9-12	20	16	10	80
8-	29-11	8-12	20	19	9	95
9-	30-11	7-12	20	19	7	95
10-	30-11	11-12	20	18	11	90
Range	-	-	-	16-20	7-13	80-100
Mean	-	-	20	18.5	10.20	92.50
SE±	-	-	0.00	1.43	1.54	7.16

217

218 **3.2.4 First nymph instar**

219 The first nymph instar emerged by pushing the eggshell, small in size, reddish
220 brown to black. Nymph's head, antennae and thorax changed to black after few
221 minutes from hatching, while the abdomen was orange in colour and appeared flat
222 in shape. It moves slowly searching for food and shelter. The mean duration of this
223 nymph instar was (7.40±0.49) days and ranged from 6 to 8 days this finding
224 coincides with other workers [8] [19]. The other four nymph instars each had unique
225 characters and showed different duration period, the mean total nymph instars is
226 51.6±2.42 (Table 3).

227 **3.2.5 Second nymph instar**

228 The body of this nymph stage is generally flat, with orange colour abdomen, creamy
229 thorax and dark brown thoracic appendages. The head and its appendages were
230 also dark brown. The mean duration of this stage was (11.3±1.10) days and the
231 result reported by [19]. This nymph moved quickly searching for food and shelter
232 (Table 3).

233 **3.2.6 Third nymph instar**

234 The body shape of this nymph instar is oval in shape, creamy in colour with two
235 creamy traverse stripes in the dorsal sides, the nymph head, antennae and thorax
236 were dark brown. The mean duration and range of this nymph instar were found to
237 be (6.1±0.7) days, 5 to 7 days respectively (Table 3), this results in agreement with
238 other workers [7][19].

239 **3.2.7 Fourth nymph instar**

The structure of this nymph was featured out, the abdomen become flattened with brown colour interrupted with creamy stripes along each of body sides. The mean duration period was (12.7±1.79) days and the range was 10 to 15 days (Table 3) this result slightly differ from [19] this may be due to differences in the climate conditions.

3.2.8 Fifth nymph instar

This stage is nearly similar to the fourth nymph instars in shape and colour. The main duration period took (14.1±0.83) days, ranged from 13 to 15 days (Table 3) these developmental stages came as reported by [19] and slightly differ from [8].

Table 3. The durations of nymph developmental stages of the melon bug *Coridius viduatus* (Fabricius) in days under laboratory conditions.

No.	First instar	Second instar	Third instar	Forth instar	Fifth instar	Total nymph days
1	7	10	6	10	13	46
2	7	12	6	10	13	50
3	8	11	6	11	15	51
4	7	11	7	12	13	50
5	8	12	7	13	13	53
6	8	13	5	13	14	53
7	8	13	6	14	14	55
8	7	10	6	14	15	52
9	7	10	7	15	15	54
10	7	11	5	15	14	52
Mean	7.4	11.3	6.1	12.7	14.1	51.6
Range	7.8	10-13	5-7	10-15	13-15	46-55
SE±	0.49	1.10	0.70	1.79	0.83	2.42

3.3 Food preference and non-preference:

There was a significant difference between the improved watermelon variety Crimson, sugar baby and the local watermelon variety Saphinga, It worth mentioning that the bugs were strongly attracted to the improved watermelon varieties Crimson (78%), Sugar baby (72%) and to lesser extent to Congo (69%). A significant difference between Ananas and Harshya cucurbit varieties was found, the local varieties to determine the sheltering sites and the nature of the watermelon bug movement were least preferred by the bug (Table 4) the bug preferred the imported improved varieties which may be more susceptible than the local varieties.

Table 4. Mean percentages of food preference of watermelon bug *Coridius viduatus* on different watermelon and pumpkin varieties.

Variety	Mean %
Crimson	78a

Sugar baby	72a
Congo	69ab
Ananas	64b
Saphinga	61b
Harshya	49c
Mean	67.3
SE*±	5.7
C.V**	8.4
LSD***	11.77

263 *SE = Standard error

264 **C.V = Coefficient of variation

265 ***LSD = Least Significant Difference

266

267 4. CONCLUSION

268 The bug aestivation shelters were determined. A regular movement of the watermelon bug
 269 was discovered from aestivation and sheltering site mainly shrubs and trees in addition to
 270 mountain crevices and soil cracks, so control measures may consider this in future. Different
 271 developmental stages were investigated. Local watermelon varieties could be cultivated as
 272 they were less preferred by the bug, however further work should be done to improve their
 273 productivity and tolerance.

274 References:

275 1. TarlaS, Yetiir H, Tarla G. Black Watermelon Bug *Coridius viduatus* (F.)(Heteroptera:
 276 Dinidoridae) in Hatay Region of Turkey. Journal of Basic and Applied Sciences.2013; 9:31-35.

277 2. CoblyLS. An Introduction to the Botony of Tropical Crops; 1979.

278 3. Satyagopal K, Sushil SN, Jeyakumar P, Shankar G, Sharma OP, et al. AESA based IPM
 279 package for Watermelon. National Institute of Plant Health Management, Rajendranagar,
 280 Hyderabad 500 030; 2014.

281

282 4. Aljedani DM. Assessment of Effectiveness of the Imidacloprid and Azadirachtin
 283 on the Black Watermelon Bug. International Journal of Zoological Research. 2018;
 284 14(2):61-70.

285

286 5. Pollard DG.A note on the biology of *Calideaduo decciumpunctata* (Fabr.) in the Sudan.
 287 Entomological Section Research Division, Min. of Agric., Sudan; 1955.

288

289 6. Schumeterer H. Pests of crops in Northeast and central Africa. Gustav Fisher Stuttgart;
 290 1969.

- 291 7. Mustafa NEM, Mariod AA, Matthauss B. Antibacterial Activity of the *Aspongopus viduatus*
292 (Melon bug) oil. J Food Safety. 2008; 28:577-86.
293
- 294 8. MohammedSalih, MHM. Studies on the biology, morphology and host range of the melon
295 bug *Coridius viduatus* (Fabricius) (Hemiptera: Dinidoridae) in the Northern State. M.Sc. thesis
296 U. of Kordofan. Sudan; 2008.
- 297 9. Mariod AA, Matthauss B, Abdelwahab SI. Fatty Acids, Tocopherols of *Aspongopus viduatus*
298 (melon bug) oil during different
299 Maturity stages. Int J Nat Prod Pharm Sci. 2011; 2(1): 20-27
300
- 301 10. Ibrahim IA. Studies on Life tables, Ecology and Efficiency of the aphidophagous
302 coccinellid *Cheilomenes vicina* (Muls). (Coccinellidae: Coleoptera). M. Sc. thesis, University
303 of Khartoum; 1988.
- 304 11. Guddoura E. Insects and disease. Attacking cucurbits .PPD Bull. No.1. plant protection
305 Department .The Ministry of Agriculture, Sudan; (1972).
- 306 12. Dabrowski ZT, Alsaffar, AA, Abdurrahman, AA. Integrated pest management on vegetable
307 crops in the Sudan. Development and implementation strategy. 1994; ICIPE Science
308 press, Nairobi, Kenya David, A. Riger: .24-32.
- 309 13. Mohammed AO. Studies on the Biology, Ecology and Possible control measures of the
310 melon bug *Coridius viduatus* (Fabricius) (Heteroptera: Dinidoridae). M.Sc. thesis Faculty of
311 Natural Resources, University of Kordofan, Sudan; 2003.
- 312 14. Ahmed AI. Studies on the ecology and control of the melon bug *Coridius viduatus*
313 (Fabricius) (Heteroptera: Dinidoridae), M.Sc. Thesis, Faculty of Natural Resources, University
314 of Kordofan, Sudan; 2004.
- 315 15. Adra, A1. Biological, Ecological, and Morphological studies on the melon bug *Coridius*
316 *viduatus* (Fabricius) Hemiptera, Dinidoridae on watermelon in North Kordofan State M.Sc.
317 thesis Faculty of Natural Resources University of Kordofan, Sudan; 2005.
- 318 16. Abdalla MI. Ecology, Biology, Behaviour and control of the melon bug, *Aspongopus*
319 *viduatus* (Fabricius) (Hemiptera: Dinidoridae). Ph.D. thesis, University of Khartoum, Sudan;
320 2010.
- 321 17. Gubarga SM. Pests status, ecology and control. Plant protection department annual
322 report, Ministry of agriculture, Khartoum, Sudan: 2002.
- 323 18. Anonymous .G.TZ project, ISVFF. Tech. pamphlet No (4/1000) April, 1992, No. (5,6,7 and
324 8/15000 may, 1992). Khartoum- Sudan. Press Cairo; (in Arabic language).
- 325 19. Mahgoub I A, Mohammed EE, Ahmed MA. Biology and fecundity of the melon bug
326 *Aspongopus viduatus* (Fabricius) in the laboratory. International Journal of Science,
327 Environment and Technology. 2015; 4 (2) 2015: 414–423.
328
- 329 20. Mohamed AMO. Effect of the entopathogenetic Fungus *Meterium anisopliae*
330 (Metsch), on watermelon bug *Coridius vidautus* (F.) (Heteroptera: Dinidoridae), under
331 Laboratory Conditions. M Sc thesis University of Gezira, Sudan; 2017.
332
333

Appendix




Fig. 1 Watermelon study area North Darfur State- Sudan