

Sg-continuity in Topological ordered spaces

Abstract

Semi generalized closed set in a Topological space was introduced by P.Bhattacharya and B.K.Lahiri in 1987. A subset A of a topological space (X, τ) is a semi generalized closed (sg-closed) set if $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is semi open in (X, τ) . Some authors introduced the notion of sg-continuity in topological spaces. The same notion can be extended to topological ordered spaces. A Topological ordered space is a topological space together with a partial order. In this paper, we introduce and study the notion of semi generalized increasing continuous function (sgi-continuous function), semi generalized decreasing continuous function (sgd-continuous function) and semi generalized balanced continuous function (sgb-continuous function) and study the relationships between them.

Mathematics subject classification: 54A05

Key words: Topological ordered space, increasing set, decreasing set, balanced set and semi generalized closed set.

1. Introduction

The study of Topological ordered spaces (TOS) was introduced by L.Nachbin [2]. It is a triple $(X, \tau, \leq)$ where τ is a topology and $\leq$ is a partial order on X . Let $(X, \tau, \leq)$ be a TOS. For any $x \in X$, $[x, \rightarrow] = \{y \in X / x \leq y\}$ and $[\leftarrow, x] = \{y \in X / y \leq x\}$. A subset A of a TOS $(X, \tau, \leq)$ is increasing if $A = i[A]$ and decreasing if $A = d[A]$ where $i[A] = \bigcup_{a \in A} [a, \rightarrow]$ and $d[A] = \bigcup_{a \in A} [\leftarrow, a]$. The complement of an increasing set is a decreasing set and vice versa. A subset of a TOS $(X, \tau, \leq)$ is balanced set if it is both increasing and decreasing.

2. Preliminaries

In the present paper (X, τ) represent a non-empty topological space on which no separation axioms are assumed unless otherwise mentioned. For a subset A of (X, τ) , the closure is the intersection of all closed sets containing A and semi closure is the intersection of all semi closed sets containing A . They are denoted by $cl(A)$ and $scl(A)$. respectively.

Definition 2.1. A subset A of a topological space (X, τ) is a sg-closed set [1] if $scl(A) \subseteq U$ whenever $A \subseteq U$ and U is semi open in (X, τ) .

The following definition is due to [7].

Definition 2.2. A subset A of a topological ordered space $(X, \tau, \leq)$ is a sgi-closed (resp. sgd-closed, sgb-closed) set if A is sg-closed and increasing (resp. decreasing, balanced).

3. Sg-continuity in Topological ordered spaces

We define new types of semi generalized continuous functions in topological ordered spaces.

A function $f : (X, \tau_1) \rightarrow (Y, \tau_2)$ is sg-continuous [1] if $f^{-1}(V)$ is sg-closed whenever V is a closed set in Y .

The following notions are introduced in a topological ordered space.

Definition 3.1. A function $f : (X, \tau, \leq) \rightarrow (Y, \tau^1, \leq^1)$ is

(1) a semi generalized increasing continuous function (briefly sgi-continuous) if $f^{-1}(V)$ is a sgi-closed set in X whenever V is an i-closed set in Y .

(2) a semi generalized decreasing continuous function (briefly sgd-continuous) if $f^{-1}(V)$ is a sgd-closed set in X whenever V is a d-closed set in Y .

(3) a sg-balanced continuous function (briefly sgb-continuous) if $f^{-1}(V)$ is a sgb-closed set in X whenever V is a b-closed set in Y .

The following examples support the above definitions.

Example 3.2. Let $X = Y = \{a, b, c\}$, $\tau_8 = \{\phi, X, \{a, b\}\}$, $\leq_3 = \{(a, a), (b, b), (c, c), (a, b), (a, c)\}$, $\leq_4 = \{(a, a), (b, b), (c, c), (a, b), (c, a), (c, b)\}$. Then $(X, \tau_8, \leq_3)$ and $(Y, \tau_8, \leq_4)$ are topological ordered spaces. The i-closed sets in Y are ϕ, X and sgi-closed sets in X are $\phi, X, \{c\}, \{b, c\}$. Define $f : X \rightarrow Y$ as $f(a) = b$, $f(b) = c$ and $f(c) = a$ then, f is a sgi-continuous function.

Example 3.3. Let $X = Y = \{a, b, c\}$, $\tau_7 = \{\phi, X, \{a\}, \{b\}, \{b, c\}, \{a, b\}\}$, $\tau_8 = \{\phi, X, \{a, b\}\}$, $\leq_7 = \{(a, a), (b, b), (c, c), (b, a)\}$, $\leq_8 = \{(a, a), (b, b), (c, c)\}$. Then, $(X, \tau_7, \leq_7)$ and $(Y, \tau_8, \leq_8)$ are topological ordered spaces. The d-closed sets in Y are $\phi, X, \{c\}$ and sgd-closed sets in X are $\phi, X, \{c\}, \{b, c\}$. Define $f : X \rightarrow Y$ as $f(a) = b$, $f(b) = a$ and $f(c) = c$ then, f is a sgd-continuous function.

Example 3.4. Let $X = Y = \{a, b, c\}$, $\tau_8 = \{\phi, X, \{a, b\}\}$, $\tau_9 = \{\phi, X, \{b\}, \{c\}, \{b, c\}\}$
 $\leq_1 = \{(a, a), (b, b), (c, c), (a, b), (b, c), (a, c)\}$, $\leq_5 = \{(a, a), (b, b), (c, c), (a, c), (b, c)\}$
 Then, $(X, \tau_8, \leq_5)$ and $(Y, \tau_9, \leq_1)$ are topological ordered spaces. The b-closed sets in Y
 are ϕ, X and sgb-closed sets in X are ϕ, X . Define $f : X \rightarrow Y$ as $f(a) = b$, $f(b) = c$
 and $f(c) = a$. Then, f is a sgb-continuous function.

4. Independency of the functions

Theorem 4.1: The notions sgi-continuity and sgd-continuity are independent.

Proof: The following example proves the theorem.

Let $X = Y = \{a, b, c\}$, $\tau_8 = \{\phi, X, \{a, b\}\}$, $\leq_3 = \{(a, a), (b, b), (c, c), (a, b), (a, c)\}$
 $\leq_4 = \{(a, a), (b, b), (c, c), (a, b), (c, a), (c, b)\}$. Then, $(X, \tau_8, \leq_3)$ and $(Y, \tau_8, \leq_4)$ are topological
 ordered spaces. The i-closed sets in Y are ϕ, X and sgi-closed sets in X are $\phi, X, \{c\}, \{b, c\}$.
 Define $f : X \rightarrow Y$ as $f(a) = a$, $f(b) = c$ and $f(c) = b$ then f is a **sgi-continuous function**.
 The d-closed sets in Y are $\phi, X, \{c\}$ and the sgd-closed sets in X are $\phi, X, \{a, c\}$. Then f is
not a sgd-continuous function.

If we take $X = Y = \{a, b, c\}$, $\tau_8 = \{\phi, X, \{a, b\}\}$, $\leq_4 = \{(a, a), (b, b), (c, c), (a, b), (c, a), (c, b)\}$
 $\leq_5 = \{(a, a), (b, b), (c, c), (a, c), (b, c)\}$. Then, $(X, \tau_8, \leq_4)$ and $(Y, \tau_8, \leq_5)$ are topological ordered
 spaces. The i-closed sets in Y are $\phi, X, \{c\}$ and sgi-closed sets in X are ϕ, X . Define
 $f : X \rightarrow Y$ as $f(a) = a$, $f(b) = a$ and $f(c) = c$. Then, f is a **not a sgi-continuous**
function. The d-closed sets in Y are ϕ, X and sgd-closed sets in X are $\phi, X, \{a, c\}, \{c\}$. Then,
 f is a **sgd-continuous function**.

Theorem 4.2: The notions sgi-continuity and sgb-continuity are independent.

Proof: This can be seen from the following example.

In the topological ordered spaces $(X, \tau_8, \leq_4)$ and $(Y, \tau_8, \leq_5)$
 where $X = Y = \{a, b, c\}$, $\tau_8 = \{\phi, X, \{a, b\}\}$, $\leq_4 = \{(a, a), (b, b), (c, c), (a, b), (c, a), (c, b)\}$ and
 $\leq_5 = \{(a, a), (b, b), (c, c), (a, c), (b, c)\}$, the i-closed sets in Y are $\phi, X, \{c\}$ and sgi-closed sets
 in X are ϕ, X . Define $f : X \rightarrow Y$ as $f(a) = a$, $f(b) = a$ and $f(c) = c$. Then, f is a **not a**
sgi-continuous function. The b-closed sets in Y are ϕ, X and sgb-closed sets in X are
 ϕ, X . Then, f is a **sgb-continuous function**.

On the other hand, in the spaces $(X, \tau_{11}, \leq_7)$ and $(Y, \tau_{11}, \leq_8)$ where $X = Y = \{a, b, c\}$, $\tau_{11} = \{\phi, X, \{c\}, \{c, b\}\}$, $\leq_7 = \{(a, a), (b, b), (c, c), (b, a)\}$ and $\leq_8 = \{(a, a), (b, b), (c, c)\}$, the i -closed sets in Y are $\phi, X, \{a\}, \{a, b\}$ and sgi -closed sets in X are $\phi, X, \{a\}, \{a, b\}$. Define $f : X \rightarrow Y$ as $f(a) = a$, $f(b) = b$ and $f(c) = c$. Then, f is a **sgi-continuous function**. The b -closed sets in Y are $\phi, X, \{a\}, \{a, b\}$ and sgb -closed sets in X are $\phi, X, \{a, b\}$. Then, f is **not a sgb-continuous function**.

Theorem 4.2: The notions sgd -continuity and sgb -continuity are independent.


Proof: This can be observed from the following example.

Consider the spaces $(X, \tau_{11}, \leq_8)$ and $(Y, \tau_{11}, \leq_9)$ where $X = Y = \{a, b, c\}$, $\leq_8 = \{(a, a), (b, b), (c, c)\}$ and $\leq_9 = \{(a, a), (b, b), (c, c), (a, c)\}$. The b -closed sets in Y are ϕ, X and sgb -closed sets in X are $\phi, X, \{a\}, \{b\}, \{a, b\}$. Define $f : X \rightarrow Y$ as $f(a) = c$, $f(b) = b$ and $f(c) = a$. Then, f is a **sgb-continuous function**. The d -closed sets in Y are $\phi, X, \{a\}, \{a, b\}$ and sgd -closed sets in X are $\phi, X, \{a\}, \{b\}, \{a, b\}$. Then, f is **not a sgd-continuous function**.

For the other part, consider the topological ordered spaces $(X, \tau_{11}, \leq_6)$ and $(Y, \tau_{11}, \leq_7)$ where $X = Y = \{a, b, c\}$, $\leq_6 = \{(a, a), (b, b), (c, c), (b, a), (a, c), (b, c)\}$ and $\leq_7 = \{(a, a), (b, b), (c, c), (b, a)\}$. The d -closed sets in Y are $\phi, X, \{a, b\}$ and sgd -closed sets in X are $\phi, X, \{b\}, \{a, b\}$. Define $f : X \rightarrow Y$ as $f(a) = a$, $f(b) = b$ and $f(c) = c$. Then, f is a **sgd-continuous function**. The b -closed sets in Y are $\phi, X, \{a, b\}$ and sgb -closed sets in X are ϕ, X . Then, f is **not a sgb-continuous function**.

Conclusion:

The following results were proved in this paper.



Here the symbol $A \leftrightarrow B$ indicates A and B are independent notions.

References

- [1] Bhattacharyya, P., Lahiri B. K. (1987), *Semi generalized closed sets in topology*. Indian J. Pure. Appl. Math 29:375–382.
- [2] L. Nachbin, *Topology and order*, D. Van Nostrand Inc., Princeton, New Jersey (1965).
- [3] M. K. R. S. Veera Kumar, *Homeomorphisms in topological ordered spaces*, Acta Ciencia Indica, XXVIII (M) (1) (2002), 67-76.
- [4] V. V. S. Ramachandram, B. Sankara Rao and M. K. R. S. Veera Kumar, *g -closed type, g^* -closed type and sg -closed type sets in topological ordered spaces*, Diophantus J. Math., 4(1) (2015), 1-9.
