

Study for Uniform Convergence and Power Series

Abstract: It has been preliminary researched that function series and power series in mathematical analysis course. There are some basic properties and the basic conclusion in the courses. This article is based on the basic theory and properties, for them to make further in-depth study. First of all, as a necessary tool, it has introduced the two properties of definite integral, it is proved that the continuous function sequence limit problem under the definite integral, then it is defined the sequence of functions on subsets of real number set uniformly Cauchy's concept, basis on them several theorem is proved, it is obtained that results of a series of important properties of function terms. Using of these properties, power series of several important theorems are proved, which is about the important properties of the power series again.

Key words: mathematical analysis course; function series; power series; uniform convergence

1. Introduction

This article assumes that the reader is familiar with the basic theory of mathematical analysis course^[1] and its basic results^[1-7]. Basic on these theories and results, the properties of function series^[8-11] is been further studied, it is obtained that the important properties of uniform convergence^[12-15] and power series^[16-18].

Our next theorem shows one can interchange integrals and uniform limits^[1-7]. The adjective “uniform” here is important. We don't prove it, but admits it directly because in the mathematical analysis course^[1] exist its proof.

Discussion 1. To prove Theorem 1 below we merely use some basic facts about integration which should be familiar [or believable] even if that calculus is rusty. Specifically, we use:

(a) If g and h are integrable on $[a, b]$ and if $g(x) \leq h(x)$ for all $x \in [a, b]$, then

$$\int_a^b g(x)dx \leq \int_a^b h(x)dx.$$

We also use the following corollary:

(b) If g is integrable on $[a, b]$, then

$$\left| \int_a^b g(x)dx \right| \leq \int_a^b |g(x)|dx$$

Continuous functions on closed intervals are integrable, as noted mathematical analysis course^[1].

2. The proof of Theorem 1

Now, we begin to prove Theorem 1.

Theorem 1. Let (f_n) be a sequence of continuous functions on $[a, b]$, and suppose

37 $f_n \rightarrow f$ uniformly on $[a, b]$. Then

$$38 \quad \lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b f(x) dx \quad (1)$$

39 **Proof.** By Theorem^[1-7] f is continuous, so the functions are all integrable on $[a, b]$. Let
40 $\varepsilon > 0$. Since $f_n \rightarrow f$ uniformly on $[a, b]$, there exists a number N such that

$$41 \quad |f_n(x) - f(x)| < \frac{\varepsilon}{b-a} \quad \text{for all } x \in [a, b] \text{ and } n > N.$$

42 Consequently $n > N$ implies

$$43 \quad \left| \int_a^b f_n(x) dx - \int_a^b f(x) dx \right| = \left| \int_a^b [f_n(x) - f(x)] dx \right|$$

$$44 \quad \leq \int_a^b |f_n(x) - f(x)| dx \leq \int_a^b \frac{\varepsilon}{b-a} dx = \varepsilon.$$

45 The first \leq follows from Discussion 1(b) applied to $g = f_n - f$ and the second \leq follow

46 from Discussion 1(a) applied to $g = |f_n - f|$ and $h = \frac{\varepsilon}{b-a}$; h happens to be a constant
47 function, but this does no harm.

48 The last paragraph shows that given $\varepsilon > 0$, there exists N such that

$$49 \quad \left| \int_a^b f_n(x) dx - \int_a^b f(x) dx \right| \leq \varepsilon \quad \text{for } n > N.$$

50 Therefore (1) holds.

51 Recall one of the advantages of the notion of Cauchy sequence, A sequence (s_n) of real
52 numbers can be shown to converge without knowing its limit by simply verifying that it is a
53 Cauchy sequence. Clearly a similar result for sequences of functions would be valuable, since
54 it is likely that we will not know the limit function in advance. What we need is the idea of
55 “uniformly Cauchy.”

56 **3. A definition and its properties about the sequence of functions**

57 **Definition 1.** A sequence (f_n) of functions defined on a set $S \subseteq \mathbb{R}$ is uniformly
58 Cauchy on S if

59 for each $\varepsilon > 0$ there exists a number N such that

$$60 \quad |f_n(x) - f_m(x)| < \varepsilon \quad \text{for all } x \in S \text{ and all } m, n > N. \quad (3.1)$$

61 Compare this definition with that of a Cauchy sequence of real numbers and that of
62 uniform convergence. It is an easy exercise to show uniformly convergent sequences of
63 functions are uniformly Cauchy. The interesting and useful result is the converse, just as in
64 the case of sequences of real numbers.

65 **Theorem 2.** Let (f_n) be a sequence of functions and uniformly Cauchy on a set $S \subseteq \mathbb{R}$.

66 Then there exists a function f on S such that $f_n \rightarrow f$ uniformly on S .

67 **Proof.** First we have to “find” f . We begin by showing

68 for each $x_0 \in S$ the sequence $(f_n(x_0))$ is a

69 Cauchy sequence of real numbers.. (3.2)

70 For each $\varepsilon > 0$, there exists N such that $|f_n(x) - f_m(x)| < \varepsilon$ for $x \in S$ and $m, n > N$.

71 In particular, we have

$$72 \quad |f_n(x_0) - f_m(x_0)| < \varepsilon \text{ for } m, n > N.$$

73 This shows $(f_n(x_0))$ is a Cauchy sequence, so (3.1) holds.

74 Now for each x in S , assertion (3.1) implies $\lim_{n \rightarrow \infty} f_n(x)$ exists; this is proved in

75 Theorem^[1-7] which in the end depends on the Completeness Axiom. Hence we define

76 $f(x) = \lim_{n \rightarrow \infty} f_n(x)$. This defines a function f on S such that $f_n \rightarrow f$ uniformly on S .

77 Now that we have “found” f , we need to prove $f_n \rightarrow f$ uniformly on S . Let $\varepsilon > 0$.

78 There is a number N such that

$$79 \quad |f_n(x_0) - f_m(x_0)| < \frac{\varepsilon}{2} \text{ for all } x \in S \text{ and all } m, n > N. \quad (3.2)$$

80 Consider $m > N$ and $x \in S$. Assertion (2) tells us that $f_n(x)$ lies in the open interval

81 $\left(f_m(x) - \frac{\varepsilon}{2}, f_m(x) + \frac{\varepsilon}{2}\right)$ for all $n > N$. Therefore, as a easy fact, the $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ lies

82 in the closed interval $\left[f_m(x) - \frac{\varepsilon}{2}, f_m(x) + \frac{\varepsilon}{2}\right]$. In other words,

$$83 \quad |f(x) - f_m(x)| \leq \frac{\varepsilon}{2} \text{ for all } x \in S \text{ and all } m > N.$$

84 Then of course

$$85 \quad |f(x) - f_m(x)| < \varepsilon \text{ for all } x \in S \text{ and all } m > N.$$

86 This shows $f_m(x) \rightarrow f$ uniformly on S , as desired.

87 Theorem 2 is especially useful for “series of functions.” Let us recall what $\sum_{k=1}^{\infty} a_k$

88 signifies when the a_k ’s are real numbers. This signifies $\lim_{n \rightarrow \infty} \sum_{k=1}^n a_k$ provided this limit exists

89 [as a real number, $+\infty$ or $-\infty$]. Otherwise the symbol $\sum_{k=1}^{\infty} a_k$ has no meaning. Thus the infinite


90 series is the limit of the sequence of partial sums $\sum_{k=1}^n a_k$. Similar remarks apply to series of
 91 functions. A series of functions is an expression $\sum_{k=0}^{\infty} g_k$ or $\sum_{k=0}^{\infty} g_k(x)$ which makes sense
 92 provided the sequence of partial sums converges, or diverges to $-\infty$ or $+\infty$ pointwise. If the
 93 sequence of partial sums $\sum_{k=0}^{\infty} g_k$ converges uniformly on a set S to $\sum_{k=0}^{\infty} g_k$, then we say the
 94 series is uniformly convergent on S .

95 4. Application and examples

96 **Example 1.** Any power series is a series of functions, since $\sum_{k=0}^{\infty} a_k x^k$ has the form
 97 $\sum_{k=0}^{\infty} g_k$ where $g_k(x) = a_k x^k$ for all x .

98 **Example 2.** $\sum_{k=0}^{\infty} \frac{x^k}{1+x^k} g_k$ is a series of functions, but is not a power series, at least not
 99 in its present form. This is a series $\sum_{k=0}^{\infty} g_k$ where $g_0(x) = \frac{1}{2}$ for all x , $g_1(x) = \frac{x}{1+x}$ for all x ,
 100 $g_2(x) = \frac{x^2}{1+x^2}$ for all x , etc.

101 **Example 3.** Let g be the function drawn in Fig. 1,



102 Fig. 1

103 and let $g_n(x) = g(4^n x)$ for all $x \in \mathbb{R}$. Then $\sum_{n=0}^{\infty} \left(\frac{3}{4}\right)^n g_n(x)$ is a series of functions. The

104 limit function f is continuous on \mathbb{R} , but has the amazing property that it is not differentiable at
 105 any point! The proof of the non-differentiability of f is somewhat delicate^[1-7].

106 Theorems for sequences of functions translate easily into theorems for series of

functions. Here is an example.

Theorem 3. Consider a series $\sum_{k=0}^{\infty} g_k$ of functions on a set $S \subseteq R$. Suppose each g_k

is continuous on S and the series converges uniformly on S . Then the series $\sum_{k=0}^{\infty} g_k$ represents a continuous function on S .

Proof. Each partial sum $f_n = \sum_{k=1}^n g_k$ is continuous and the sequence (f_n) converges uniformly on S . Hence the limit function is continuous by Theorem^[1-7].

Recall the Cauchy criterion for series $\sum_{k=1}^{\infty} a_k$ given in paper^[1-7] :

For each $\varepsilon > 0$ there exists a number N such that

$$n \geq m > N \text{ implies } \left| \sum_{k=m}^n a_k \right| < \varepsilon . \quad (*)$$

The analogue for series of functions is also useful. The sequence of partial sums of a series $\sum_{k=0}^{\infty} g_k$ of functions is uniformly Cauchy on a set S if and only if the series satisfies the Cauchy criterion [uniformly on S]:

For each $\varepsilon > 0$ there exists a number N such that

$$n \geq m > N \text{ implies } \left| \sum_{k=m}^n g_k(x) \right| < \varepsilon . \text{ for all } x \in S \quad (**)$$

Theorem 4. If a series $\sum_{k=0}^{\infty} g_k$ of functions satisfies the Cauchy criterion uniformly on a set S , then the series converges uniformly on S by Theorem 2.

Here is a useful corollary.

Theorem 5 (M-test). Let (M_k) be a sequence of nonnegative real numbers where $\sum M_k < \infty$. If $|g_k(x)| \leq M_k$ for all x in a set S , then $\sum g_k$ converges uniformly on S .

Proof. To verify the Cauchy criterion on S , let $\varepsilon > 0$. Since the series $\sum M_k$ converges, it satisfies the Cauchy criterion in Definition^[1-7]. So there exists a number N such that

$$n \geq m > N \text{ implies } \sum_{k=m}^n M_k < \varepsilon .$$

Hence if $n \geq m > N$ and x is in S , then

$$\left| \sum_{k=m}^n g_k(x) \right| \leq \sum_{k=m}^n |g_k(x)| \leq \sum_{k=m}^n M_k < \varepsilon$$

Thus the series $\sum_{k=0}^{\infty} g_k$ satisfies the Cauchy criterion uniformly on S , and Theorem 4 shows it converges uniformly on S .

Example 4. Show $\sum_{n=1}^{\infty} 2^{-n} x^n$ represents a continuous function f on $(-2, 2)$, but the convergence is not uniform.

Solution. This is a power series with radius of convergence 2. Clearly the series does not converge at $x=2$ or at $x=-2$, so its interval of convergence is $(-2, 2)$.

Consider $0 < a < 2$ and note

$$\sum_{n=1}^{\infty} 2^{-n} a^n = \sum_{n=1}^{\infty} \left(\frac{a}{2} \right)^n$$

converges. Since

$$|2^{-n} x^n| \leq 2^{-n} a^n = \left(\frac{a}{2} \right)^n \text{ for } x \in [-a, a],$$

the Theorem 5 (**M-test**) shows the series converges uniformly to a function on $[-a, a]$. By Theorem 3 the limit function f is continuous at each point of the set $[-a, a]$. Since a can be any number less than 2, we conclude f represents a continuous function on $(-2, 2)$.

Since we have $\sup \{ |2^{-n} x^n| \mid x \in (-2, 2) \} = 1$ for each n , the convergence of the series cannot be uniform on $(-2, 2)$ in view of the next example.

Example 5. Show that if the series $\sum g_n$ converges uniformly on a set S , then

$$\lim_{n \rightarrow \infty} \sup \{ |g_n(x)| \mid x \in S \} = 0. \quad (1)$$

Solution. Let $\varepsilon > 0$. Since the series $\sum g_n$ satisfies the Cauchy criterion, there exists N such that

$$n \geq m > N \text{ implies } \left| \sum_{k=m}^n g_k(x) \right| < \varepsilon \text{ for all } x \in S.$$

In particular,

$$n > N \text{ implies } |g_n(x)| < \varepsilon \text{ for all } x \in S.$$

Therefore

$$n > N \text{ implies } \sup \{ |g_n(x)| \mid x \in S \} \leq \varepsilon.$$

This establishes (1).

5. Properties of power series

Now we begin to study the properties of the power series.

Theorem 6. Let $\sum_{n=0}^{\infty} a_n x^n$ be a power series with radius of convergence $R > 0$

[possibly $R = +\infty$]. If $0 < R_1 < R$, then the power series converges uniformly on $[-R_1, R_1]$ to a continuous function.

Proof. Consider $0 < R_1 < R$. A glance at Theorem^[1-7] shows the series $\sum_{n=0}^{\infty} a_n x^n$ and

$\sum_{n=0}^{\infty} |a_n| x^n$ have the same radius of convergence, since β and R are defined in terms of

$|a_n|$. Since $|R_1| < R$, we have $\sum_{n=0}^{\infty} |a_n| R_1^n < \infty$. Clearly we have $|a_n x^n| \leq |a_n| R_1^n$ for all x in

$[-R_1, R_1]$, so the series $\sum_{n=0}^{\infty} a_n x^n$ converges uniformly on $[-R_1, R_1]$ by the Theorem 5

(M-test). The limit function is continuous at each point of $[-R_1, R_1]$ by Theorem 3.

Corollary 7. The power series $\sum a_n x^n$ converges to a continuous function on the open

interval $(-R, R)$.

Proof. If $x_0 \in (-R, R)$ then $x_0 \in (-R_1, R_1)$ for some $R_1 < R$. The theorem shows the

limit of the series is continuous at x_0 .

We emphasize that a power series need not converge uniformly on its interval of convergence though it might.

We are going to differentiate and integrate power series term-by-term, so clearly it would be useful to know where the new series converge. The next lemma tells us.

Lemma 8. If the power series $\sum_{n=0}^{\infty} a_n x^n$ has radius of convergence R , then the power

series

$$\sum_{n=0}^{\infty} n a_n x^{n-1} \quad \text{and} \quad \sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1}$$

also have radius of convergence R .

Proof. First observe the series $\sum_{n=0}^{\infty} na_n x^{n-1}$ and $\sum_{n=0}^{\infty} na_n x^n$ have the same radius of convergence: since the second series is x times the first series, they converge for exactly the same values of x . Likewise $\sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1}$ and $\sum_{n=0}^{\infty} \frac{a_n}{n+1} x^n$ have the same radius of convergence.

Next recall $R = \frac{1}{\beta}$ where $\beta = \limsup |a_n|^{1/n}$. For the series $\sum_{n=0}^{\infty} na_n x^n$, we consider

$$\limsup (n |a_n|)^{1/n} = \limsup n^{1/n} |a_n|^{1/n}.$$

By Theorem^[1-7], we have $\lim n^{1/n} = 1$ so $\limsup (n |a_n|)^{1/n} = \beta$ by Theorem^[1-7]. Hence the series $\sum_{n=0}^{\infty} na_n x^n$ has radius of convergence R .

For the series $\sum_{n=0}^{\infty} \frac{a_n}{n+1} x^n$, we consider $\limsup \left(\frac{|a_n|}{n+1} \right)^{1/n}$. It is easy to show

$\lim (n+1)^{1/n} = 1$; therefore $\lim \left(\frac{1}{n+1} \right)^{1/n} = 1$. Hence by Theorem^[1-7] we have

$\limsup \left(\frac{|a_n|}{n+1} \right)^{1/n} = \beta$, so the series $\sum_{n=0}^{\infty} \frac{a_n}{n+1} x^n$ has radius of convergence R .

Theorem 9. Suppose $f(x) = \sum_{n=0}^{\infty} a_n x^n$ has radius of convergence $R > 0$. Then

$$\int_0^x f(t) dt = \sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1} \quad \text{for } |x| < R. \quad (1)$$

Proof. We fix x and assume $x < 0$; the case $x > 0$ is similar. On the interval $[x, 0]$, the sequence of partial sums $\sum_{k=0}^n a_k t^k$ converges uniformly to $f(t)$ by Theorem 6.

Consequently, by Theorem 1 we have

$$\begin{aligned} \int_x^0 f(t) dt &= \lim_{n \rightarrow \infty} \int_x^0 \left(\sum_{k=0}^n a_k t^k \right) dt \\ &= \lim_{n \rightarrow \infty} \sum_{k=0}^n a_k \int_x^0 t^k dt = \lim_{n \rightarrow \infty} \sum_{k=0}^n a_k \left[\frac{0^{k+1} - x^{k+1}}{k+1} \right] \end{aligned}$$

$$= -\sum_{k=0}^{\infty} \frac{a_k}{k+1} x^{k+1} \quad (2)$$

The second equality is valid because we can interchange integrals and finite sums; this is a basic property of integrals^[1-7]. Since $\int_0^x f(t)dt = -\int_x^0 f(t)dt$. Eq. (2) implies Eq.(1).

The theorem just proved shows that a power series can be integrated term-by-term inside its interval of convergence. Term-by-term differentiation is also legal.

Theorem 10. Let $f(x) = \sum_{n=0}^{\infty} a_n x^n$ have radius of convergence $R > 0$. Then f is differentiable on $(-R, R)$ and

$$f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} \quad \text{for } |x| < R. \quad (1)$$

The proof of Theorem 9 was a straightforward application of Theorem 1 but the direct analogue of Theorem 1 for derivatives is not true^[1-7]. So we give a devious indirect proof of the theorem.

Proof. We begin with series $g(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$ and observe this series converges for $|x| < R$ by Lemma 8. Theorem 9 shows that we can integrate g term-by-term:

$$\int_0^x g(t)dt = \sum_{n=1}^{\infty} a_n x^n = f(x) - a_0 \quad \text{for } |x| < R.$$

Thus if $0 < R_1 < R$, then

$$f(x) = \int_{-R_1}^x g(t)dt + k \quad \text{for } |x| < R_1,$$

where k is a constant; in fact,

$$k = a_0 - \int_{-R_1}^0 g(t)dt.$$

Since g is continuous, one of the versions of the Fundamental Theorem of Calculus^[1-7] shows f is differentiable and $f'(x) = g(x)$. Thus

$$f'(x) = g(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} \quad \text{for } |x| < R.$$

Example 6. Recall

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x} \quad \text{for } |x| < 1. \quad (1)$$

222 Differentiating term-by-term, we obtain

$$223 \quad \sum_{n=1}^{\infty} nx^{n-1} = \frac{1}{(1-x)^2} \quad \text{for } |x| < 1.$$

224 Integrating (1) term-by-term, we get

$$225 \quad \sum_{n=0}^{\infty} \frac{1}{n+1} x^{n+1} = \int_0^x \frac{1}{1-t} dt = -\log_e(1-x)$$

$$226 \quad \log_e(1-x) = -\sum_{n=1}^{\infty} \frac{1}{n} x^n \quad \text{for } |x| < 1. \quad (2)$$

227 Replacing x by $-x$, we find

$$228 \quad \log_e(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots \quad \text{for } |x| < 1. \quad (3)$$

229 It turns out that this equality is also valid for $x=1$ [see Example 7], so we have the interesting
230 identity

$$231 \quad \log_e 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \cdots \quad (4)$$

232 In Eq. (2) set $x = \frac{m-1}{m}$. Then

$$233 \quad \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{m-1}{m} \right)^n = -\log_e \left(1 - \frac{m-1}{m} \right) = -\log_e \left(\frac{1}{m} \right) = \log_e m$$

234 Hence we have

$$235 \quad \sum_{n=1}^{\infty} \frac{1}{n} \geq \sum_{n=1}^{\infty} \frac{1}{n} \left(\frac{m-1}{m} \right)^n = \log_e m \quad \text{for all } m.$$

236 Here is yet another proof that $\sum_{n=1}^{\infty} \frac{1}{n} = +\infty$.

237 To establish (4) we need a relatively difficult theorem about convergence of a power
238 series at the endpoints of its interval of convergence.

239 Let $f(x) = \sum_{n=0}^{\infty} a_n x^n$ be a power series with finite positive radius of convergence R . If

240 the series converges at $x=R$, then f is continuous at $x=R$. If the series converges at $x=-R$, then f
241 is continuous at $x=-R$.

242 **Example 7.** As promised, we return to (3) in Example 1:

$$243 \quad \log_e(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots \quad \text{for } |x| < 1.$$

244 For $x=1$ the series converges by the Alternating Series Theorem^[1-7]. Thus the series represents
245 a function f on $(-1, 1]$ that is continuous at $x=1$ by Abel's theorem. The function $\log_e(1+x)$

is also continuous at $x=1$ so the functions agree at $x=1$. [In detail, if (x_n) is a sequence in $(-1, 1)$ converging to 1, then $f(1) = \lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} \log_e (1 + x_n) = \log_e 2$.] Therefore we have

$$\log_e 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \cdots.$$

Example 8. Recall $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$ for $|x| < 1$. Note that at $x=-1$ the function $\frac{1}{1-x}$ is continuous and takes the value $\frac{1}{2}$. However, the series does not converge for $x=-1$, so Abel's theorem does not apply.

The point of view in our extremely brief introduction to power series has been: For a given power series $\sum a_n x^n$, what can one say about the function $f(x) = \sum a_n x^n$? This point of view was misleading. Often, in real life, one begins with a function f and seeks a power series that represents the function for some or all values of x . This is because power series, being limits of polynomials, are in some sense basic objects.

If we have

$$f(x) = \sum_{n=0}^{\infty} a_n x^n \quad \text{for } |x| < R,$$

then we can differentiate f term-by-term forever. At each step, we may calculate the k th derivative of f at 0, written $f^{(k)}(0)$. It is easy to show $f^{(k)}(0) = k! a_k$ for $k \geq 0$. This tells us that if f can be represented by a power series, then that power series must be

$\sum_{n=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k$. This is the Taylor series for f about 0. Frequently, but not always, the Taylor series will agree with f on the interval of convergence.

6. Conclusions

From the above, we have seen that the properties of the power series are very perfect, it is an extremely rare class of function series; in addition, Cauchy criterion has played important role. Using Cauchy criterion as a tool, not only can derive many properties of number series, it can also be derived a lot of properties of function series in the deep level. In addition, the limit thought is never less important tool in our study.

Reference

- [1] Charles Walmsley. An Introductory Course of Mathematical Analysis[B]. Cambridge University Press , London, February, 2016. PP120-230
- [2] Luming Shen, Chao Ma, Jihong Zhang. On the error-sum function of alternating Lüroth series[J]. Analysis in Theory and Applications . 2006 (3), 78-83

277 [3]Shen L M, Wu J. On the error-sum function of $L(u|^{-})$ roth series[J]. Journal of Mathematical Analysis
278 and Applications . 2007 (2), 78-83

279 [4]Yamin Yang. Equivalence of Gap Sequences and Hausdorff Dimensions of Serf-Similar Sets[A].
280 Department of Mathematical Sciences, Tsinghua University, Beijing, China. Proceedings of the Seventh
281 International Conference on Information and Management Sciences[C]. 2008 , 82-91

282 [5]Castillo E, Gutierrez J M. Nonlinear time modeling and prediction using functional networks, extracting
283 information masked by chaos[J]. Physics Letters . 1998 (3), 67-73

284 [6]Zhou Yongquan, He Dexu, Jiao Licheng. A Neural Computa-tion Structure for Solving Functional
285 Equations with Function-al Networks[J]. Journal of Infor mation and Computational Science . 2005 (1),
286 91-96

287 [7]Indiekofer, K. H., Knopfmacher, A, and Knopfmacher, J. Alternating Balkema-Oppenheim Expansions
288 of Real Numbers[J]. Bull.Soc. Math. Belgique. 1992 (2), 64-68

289 [8]Luming Shen, Chao Ma, Jihong Zhang. On the error-sum function of alternating Lüroth series[J].
290 Analysis in Theory and Applications . 2006 (3), 48-55

291 [9]Castillo E, Cobo A, Gutiérrez J M, etal. Working with differential, functional and difference equations
292 using functional networks[J]. Journal of Applied Mathematics . 1999 (2), 42-47

293 [10]Xiao Er Wu, Yong Shun Liang,Wei Xiao,Jun Huai Du. Hausdorff dimension of continuous functions
294 with at most finite UV points on closed intervals[A]. The 29th China control and decision-making
295 conference proceedings(III)[C]. 2017, 178-182

296 [11]Li Chunguang, Liao Xiaofeng, He Songbai, etal. Function Network Method for the Identification of
297 nonlinear system[C]. System Engineer and Electronics . 2001, 102-108

298 [12]Telyakovskii S A. On the convergence in the metric of L of trigonometric series with rarely changing
299 coefficients[C]. Trudy Mat Inst Steklov . 1991, 132-138

300 [13]Nurcombe, J. R. On the uniform convergence of sine series with quasi-monotone coefficients[J].
301 Journal of Mathematical Analysis and Applications . 1992 (1), 37-43

302 [14]Stanojevic V B. Fourier and trigonometric transforms with complex coefficients regularly varying in
303 mean[J]. Lecture Notes Pure Appl Math . 1994 (2), 38-47

304 [15]M. Ghafarian, A. Ariaei. Free vibration analysis of a system of elastically interconnected rotating
305 tapered Timoshenko beams using differential transform method[J]. International Journal of Mechanical
306 Sciences . 2015 (3), 278-283

307 [16]Manfred Droste, Guo-Qiang Zhang. On transformations of formal power series[J]. Information and
308 Computation. 2003 (2), 85-89

309 [17]Yong-an Huang, Zi-chen Deng, Lin-xiao Yao. Dynamic analysis of a rotating rigid-flexible coupled
310 smart structure with large deformations[J]. Applied Mathematics and Mechanics . 2007 (10), 271-278

311 [18]Du, H., Lim, M. K., Liew, K.M. Power series solution for vibration of a rotating Timoshenko beam[J].
312 Journal of Sound and Vibration . 1994 (1), 49-57