

SYMMETRIC q -GAMMA FUNCTION.

ABSTRACT

In this work we are interested by giving new characterizations of the symmetric q -Gamma function and show that there are intimately related.

Key Words: q -analogue, gamma function, characterization.

1 Introduction

In literature the characterizations of the well known Gamma function are studied by many authors [[2],[4],[11]]. As same as the Gamma function, the characterization of the q -Gamma function was studied by Elmonser, Brahim and Fitouhi in [5], they proved the following results:

Theorem 1 *The q -Gamma function is the unique function $f(x) > 0$ on $]0, +\infty[$ that satisfies the following properties:*

- a) $f(1) = 1$
- b) $f(x+1) = [x]_q f(x)$
- c) $f(x+n) = (1-q)^{[x]_q-x} f(n) [n]_q^{[x]_q} t_n(x)$, where $t_n(x) \rightarrow 1$ as $n \rightarrow \infty$.

The second theorem gives the relationship between three different characterizations of the q -Gamma function:

Theorem 2 *For a q -PG function f , the following properties are equivalent:*

- (C) $\ln f$ is convex on $]0, +\infty[$,
 - (L) $L(n+x) = ([x]_q - x) \ln(1-q) + L(n) + x \ln(n+1) + r_n(x)$,
where $L(x) = \ln f(x+1)$ and $r_n(x) \rightarrow 0$ as $n \rightarrow \infty$,
 - (P) $f(x+n) = (1-q)^{[x]_q-x} f(n) [n]_q^{[x]_q} t_n(x)$,
where $t_n(x) \rightarrow 1$ as $n \rightarrow \infty$.
- A q -PG function f satisfying these properties is equal to $c\Gamma_q(x)$, for some constant c .*

where the a q -PG function (pre- q -gamma function) is a positive function f on $]0, +\infty[$ satisfying the functional equation $f(x+1) = [x]_q f(x)$.

In the present paper we give characterizations of the symmetric q -Gamma function, introduced by K. Brahim and Yosr Sidomou in [3], and we show that there are intimately related.

2 Notations and Preliminaries

We recall some usual notions and notations used in the q -theory (see [6] and [8]). Throughout this paper, we assume $q \in]0, 1[$.

For $a \in \mathbb{C}$, the q -shifted factorials are defined by

$$(a; q)_0 = 1, \quad (a; q)_n = \prod_{i=0}^{n-1} (1 - aq^i) = (1-a)(1-aq)\dots(1-aq^{n-1}), \quad n = 1, 2, \dots \quad (1)$$

$$(a; q)_\infty = \prod_{i=0}^{\infty} (1 - aq^i). \quad (2)$$

We also denote

$$[x]_q = \frac{1 - q^x}{1 - q}, \quad x \in \mathbb{C}, \quad (3)$$

$$\widetilde{[x]}_q = \frac{q^x - q^{-x}}{q - q^{-1}}, \quad x \in \mathbb{C}, \quad (4)$$

$$[n]_q! = \prod_{k=1}^n [k]_q = \frac{(q; q)_n}{(1-q)^n}, \quad n \in \mathbb{N}. \quad (5)$$

and

$$\widetilde{[n]}_q! = \prod_{k=1}^n \widetilde{[k]}_q, \quad n \in \mathbb{N}. \quad (6)$$

One can see that

$$\widetilde{[x]}_q = q^{-(x-1)} [x]_{q^2}. \quad (7)$$

3 The symmetric q -Gamma function:

The q -Gamma function $\Gamma_q(x)$, a q -analogue of Euler's gamma function, was introduced by Thomae [10] and later by Jackson [7] as the infinite product:

$$\Gamma_q(x) = \frac{(q; q)_\infty (1-q)^{1-x}}{(q^x; q)_\infty}, \quad x > 0, \quad (8)$$

where q is a fixed real number $0 < q < 1$.

Recently, K. Brahim and Yosr Sidomou [see [3]] introduced the symmetric q -Gamma function as follows:

$$\widetilde{\Gamma}_q(z) = q^{-\frac{(z-1)(z-2)}{2}} \Gamma_{q^2}(z), \quad z > 0, q > 0, q \neq 1, \quad (9)$$

where

$$\Gamma_q(z) = \begin{cases} \frac{(q, q)_\infty}{(q^x, q)_\infty} (1 - q)^{1-x}, & \text{if } 0 < q < 1, \\ \frac{(q^{-1}, q^{-1})_\infty}{(q^{-x}, q^{-1})_\infty} (1 - q)^{1-x} q^{\frac{x(x-1)}{2}}, & \text{if } q > 1. \end{cases} \quad (10)$$

They proved that it is symmetric under the interchange $q \leftrightarrow q^{-1}$ and satisfies a q -analogue of the Bohr-Mollerup theorem for $q \neq 1$:

Theorem 3 *Let $q > 0$, $q \neq 1$. The only function $f \in C^2((0, \infty))$ satisfying the conditions:*

- (a) $f(1) = 1$.
 - (b) $f(x+1) = \widetilde{[x]}_q f(x)$.
 - (c) $\frac{d^2}{dx^2} \text{Log} f(x) \geq |\text{Log} q|$ for positive x ,
- is the symmetric q -Gamma function.*

Using the relation 9 and the properties of the q -Gamma function [5], we derive the following theorem:

Theorem 4

$$\widetilde{\Gamma}_q(x) = \lim_{n \rightarrow +\infty} q^{-\frac{(x-1)(x-2)}{2}} (1 - q^2)^{[x]_{q^2} - x} \frac{[n]_{q^2}^{[x]_{q^2}} [n]_{q^2}!}{[x]_{q^2} [x+1]_{q^2} \dots [x+n]_{q^2}}, \quad x > 0. \quad (11)$$

4 Characterization of the q -Gamma function:

The first characterization is given by the following theorem:

Theorem 5 *There exists a unique function $f(x) > 0$ on $]0, +\infty[$ that satisfies the following properties:*

- a) $f(1) = 1$
- b) $f(x+1) = \widetilde{[x]}_q f(x)$
- c) $f(x+n) = q^{-\frac{x^2+2nx-3x}{2}} (1 - q^2)^{[x]_{q^2} - x} [n]_{q^2}^{[x]_{q^2}} f(n) t_n(x)$, where $t_n(x) \rightarrow 1$ as $n \rightarrow \infty$.

Proof.

First we prove that $\widetilde{\Gamma}_q(x)$ satisfies conditions (a), (b) and (c).

From theorem 3, the symmetric q -Gamma function satisfies the condition (a) $\widetilde{\Gamma}_q(1) = 1$, and the condition (b) $\widetilde{\Gamma}_q(x+1) = \widetilde{[x]}_q \widetilde{\Gamma}_q(x)$.

As a consequence of the two properties, we get $\widetilde{\Gamma}_q(n) = \widetilde{[n-1]}_q!$

$$(c) \text{ Let } s_n(x) = \frac{\widetilde{\Gamma}_q(x)}{q^{-\frac{(x-1)(x-2)}{2}} (1 - q^2)^{[x]_{q^2} - x} \widetilde{\Gamma}_{n,q}(x)},$$

$$\text{where } \widetilde{\Gamma}_{n,q}(x) = \frac{[n]_{q^2}^{[x]_{q^2}} [n]_{q^2}!}{[x]_{q^2} [x+1]_{q^2} \dots [x+n]_{q^2}} = \frac{[n]_{q^2}^{[x]_{q^2}} \widetilde{[n]}_q!}{q^{nx+x-1} \widetilde{[x]}_q \widetilde{[x+1]}_q \dots \widetilde{[x+n]}_q},$$

$$\text{then } \widetilde{\Gamma}_q(x) = s_n(x) q^{-\frac{(x-1)(x-2)}{2}} (1 - q^2)^{[x]_{q^2} - x} \widetilde{\Gamma}_{n,q}(x) \text{ and } \lim_{n \rightarrow +\infty} s_n(x) = 1.$$

For $n \in \mathbb{N}$ and $x > 0$, we apply (b) n times to get

$$\begin{aligned}
\tilde{\Gamma}_q(x+n) &= \widetilde{[x+n-1]_q \dots [x+1]_q} \widetilde{[x]_q} \tilde{\Gamma}_q(x) \\
&= \frac{\widetilde{[x+n]_q \dots [x+1]_q} \widetilde{[x]_q}}{[x+n]_q} \cdot q^{-\frac{(x-1)(x-2)}{2}} (1-q^2)^{[x]_{q^2}-x} \frac{[n]_{q^2}^{[x]_{q^2}} \widetilde{[n]_q}!}{q^{nx+x-1} \widetilde{[x]_q} \widetilde{[x+1]_q} \dots \widetilde{[x+n]_q}} \cdot s_n(x) \\
&= q^{-\frac{x^2+2nx-3x}{2}} (1-q^2)^{[x]_{q^2}-x} [n]_{q^2}^{[x]_{q^2}} \tilde{\Gamma}_q(n) t_n(x).
\end{aligned}$$

Where $t_n(x) = q^{-x} \frac{\widetilde{[n]_q}}{[x+n]_q} \cdot s_n(x)$. Thus, $\tilde{\Gamma}_q(x+n) = q^{-\frac{x^2+2nx-3x}{2}} (1-q^2)^{[x]_{q^2}-x} [n]_{q^2}^{[x]_{q^2}} \tilde{\Gamma}_q(n) t_n(x)$ and $t_n(x) \rightarrow 1$ as $n \rightarrow +\infty$.

To show uniqueness, we assume $f(x)$ is a function that satisfies (a), (b) and (c). From properties (a) and (b), we have

$$f(n) = \widetilde{[n-1]_q}! \quad (12)$$

$$f(x+n) = \widetilde{[x+n-1]_q} \widetilde{[x+n-2]_q} \dots \widetilde{[x+1]_q} \widetilde{[x]_q} f(x). \quad (13)$$

Combining (12),(13) and (c) together, we have

$$\begin{aligned}
f(x) &= q^{-\frac{x^2+2nx-3x}{2}} (1-q^2)^{[x]_{q^2}-x} \frac{[n]_{q^2}^{[x]_{q^2}} \widetilde{[n-1]_q}!}{\widetilde{[x+n-1]_q} \widetilde{[x+n-2]_q} \dots \widetilde{[x+1]_q} \widetilde{[x]_q}} t_n(x) \\
&= q^{-\frac{(x-1)(x-2)}{2}} (1-q^2)^{[x]_{q^2}-x} \tilde{\Gamma}_{n,q}(x) \cdot s_n(x),
\end{aligned}$$

where $s_n(x) = q^x \frac{\widetilde{[x+n]_q}}{[n]_q} t_n(x) \rightarrow 1$ as $n \rightarrow +\infty$. Therefore $f(x) = \Gamma_q(x)$ and hence f is uniquely determined. This completes the proof.

5 Relationship between characterizations

In what follows, we will adopt the terminology of the following definition.

Definition 1 A function f is said to be a qs -PG function (pre-symmetric- q -gamma function), if f is positive on $]0, +\infty[$ and satisfies the functional equation $f(x+1) = \widetilde{[x]_q} f(x)$.

In the previous section we showed that the property

$$f(x+n) = q^{-\frac{x^2+2nx-3x}{2}} (1-q^2)^{[x]_{q^2}-x} [n]_{q^2}^{[x]_{q^2}} f(n) t_n(x)$$

characterizes the q -gamma function. In this section we will give three properties which are equivalent to one another for a qs -PG function and characterize the symmetric q -gamma function.

Theorem 6 For a q -PG function f , the following properties are equivalent:

(C) $\ln f$ is convex on $]0, +\infty[$,

$$(L) L(n+x) = -\frac{x^2+2nx-3x}{2} \ln q + ([x]_{q^2}-x) \ln(1-q^2) + L(n) + [x]_{q^2} \ln[n+1]_{q^2} + r_n(x),$$

where $L(x) = \ln f(x+1)$ and $r_n(x) \rightarrow 0$ as $n \rightarrow \infty$,

$$(P) \quad f(x+n) = q^{-\frac{x^2+2nx-3x}{2}} (1-q^2)^{[x]_{q^2}-x} [n]_{q^2}^{[x]_{q^2}} f(n) t_n(x),$$

where $t_n(x) \rightarrow 1$ as $n \rightarrow \infty$.

A qs -PG function f satisfying these properties is equal to $c\tilde{\Gamma}_q(x)$, for some constant c .

Proof.

(a) $(P) \Leftrightarrow (L)$. We have

$$\begin{aligned} (P) &\Leftrightarrow f(x+(n+1)) = q^{-\frac{x^2+2nx-3x}{2}} (1-q^2)^{[x]_{q^2}-x} f(n+1) [n+1]_{q^2}^{[x]_{q^2}} t_{n+1}(x), \\ &\quad t_{n+1}(x) \rightarrow 1 \\ &\Leftrightarrow \ln f(x+(n+1)) = -\frac{x^2+2nx-3x}{2} \ln q + ([x]_{q^2}-x) \ln(1-q^2) + \ln f(n+1) \\ &\quad + [x]_{q^2} \ln[n+1]_{q^2} + \ln t_{n+1}(x), t_{n+1}(x) \rightarrow 1 \\ &\Leftrightarrow L(x+n) = -\frac{x^2+2nx-3x}{2} \ln q + ([x]_{q^2}-x) \ln(1-q^2) + L(n) \\ &\quad + [x]_{q^2} \ln[n+1]_{q^2} + r_n(x), r_n(x) \rightarrow 0 \\ &\Leftrightarrow (L). \end{aligned}$$

(b) $(C) \Rightarrow (P)$. Let $m < x \leq m+1$, where $m = 0, 1, 2, \dots$. For any natural n , $n+m-1 < n+m < n+x \leq n+m+1$. The convexity of $\ln f$ gives us (we write $L_m = \ln f(n+m)$)

$$\begin{aligned} \frac{L_m - L_{m-1}}{n+m - (n+m-1)} &\leq \frac{\ln f(n+x) - \ln f(n+m)}{(n+x) - (n+m)} \leq \frac{L_{m+1} - L_m}{(n+m+1) - (n+m)} \\ &\Leftrightarrow (x-m) \ln \widetilde{[n+m-1]_q} \leq \ln \frac{f(n+x)}{f(n+m)} \leq (x-m) \ln \widetilde{[n+m]_q} \\ &\Leftrightarrow \widetilde{[n+m-1]_q}^{x-m} \leq \frac{f(n+x)}{[n+m-1]_q [n+m-2]_q \dots [n]_q f(n)} \leq \widetilde{[n+m]_q}^{x-m} \\ &\Leftrightarrow \widetilde{[n+m-1]_q}^x T_m \leq \frac{f(n+x)}{f(n)} \leq \widetilde{[n+m]_q}^x T_m \frac{\widetilde{[n+m-1]_q}^m}{\widetilde{[n+m]_q}^m}, \end{aligned}$$

$$\text{where } T_m = \frac{\widetilde{[n+m-1]_q} [n+m-2]_q \dots \widetilde{[n]_q}}{\widetilde{[n+m-1]_q}^m} = q^{\frac{m(m-1)}{2}} \frac{[n+m-1]_{q^2} [n+m-2]_{q^2} \dots [n]_{q^2}}{[n+m-1]_{q^2}^m}.$$

Therefore, we have

$$\lim_{n \rightarrow +\infty} q^{nx} \frac{f(n+x)}{f(n)} = \frac{q^{-\frac{x^2-3x}{2}}}{(1-q^2)^x},$$

by the squeezing theorem. If we let

$$t_n(x) = \frac{q^{\frac{x^2+2nx-3x}{2}} f(n+x)}{(1-q^2)^{[x]_{q^2}-x} f(n) [n]_{q^2}^{[x]_{q^2}}},$$

then

$$f(n+x) = q^{-\frac{x^2+2nx-3x}{2}} (1-q^2)^{[x]_{q^2}-x} f(n) [n]_{q^2}^{[x]_{q^2}} t_n(x),$$

where $t_n(x) \rightarrow 1$ as $n \rightarrow \infty$. This proves that f satisfies (P) .

(c) $(P) \Rightarrow (C)$. From the uniqueness part of the proof of the Theorem 5 we have

$$f(x) = f(1) \lim_{n \rightarrow +\infty} q^{-\frac{(x-1)(x-2)}{2}} (1 - q^2)^{[x]_{q^2} - x} \Gamma_{n,q}(x).$$

Using the fact that the limit function of a convergent sequence of convex functions is convex, it suffices to show that $\ln \left(q^{-\frac{(x-1)(x-2)}{2}} (1 - q^2)^{[x]_{q^2} - x} \Gamma_{n,q}(x) \right)$ is convex.

Now

$$\begin{aligned} \ln \left(q^{-\frac{(x-1)(x-2)}{2}} (1 - q^2)^{[x]_{q^2} - x} \Gamma_{n,q}(x) \right) &= -\frac{(x-1)(x-2)}{2} \ln q + ([x]_{q^2} - x) \ln(1 - q^2) \\ &\quad + [x]_{q^2} \ln[n]_{q^2} + \ln([n]_{q^2}!) - \ln[x]_{q^2} - \dots - \ln[x+n]_{q^2}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \left(\ln \left(q^{-\frac{(x-1)(x-2)}{2}} (1 - q^2)^{[x]_{q^2} - x} \Gamma_{n,q}(x) \right) \right)' &= (-x + \frac{3}{2}) \ln q + \left(-2 \frac{\ln q}{1 - q^2} q^{2x} - 1 \right) \ln(1 - q^2) \\ &\quad + \left(-2 \frac{\ln q}{1 - q^2} q^{2x} \ln[n]_{q^2} \right) + \frac{2 \ln q}{1 - q^2} \frac{q^{2x}}{[x]_{q^2}} + \dots \\ &\quad + \frac{2 \ln q}{1 - q^2} \frac{q^{2(x+n)}}{[x+n]_{q^2}}. \end{aligned}$$

And so

$$\begin{aligned} \left(\ln \left(q^{-\frac{(x-1)(x-2)}{2}} (1 - q^2)^{[x]_{q^2} - x} \Gamma_{n,q}(x) \right) \right)'' &= -\ln q - 4 \frac{(\ln q)^2}{1 - q^2} q^{2x} (\ln(1 - q^2) + \ln \frac{1 - q^{2n}}{1 - q^2}) \\ &\quad + 4 \frac{(\ln q)^2}{1 - q^2} \left[\frac{q^{2x} [x]_{q^2} + \frac{q^{4x}}{1 - q^2}}{[x]_{q^2}^2} + \dots \right. \\ &\quad \left. + \frac{q^{2(x+n)} [x+n]_{q^2} + \frac{q^{4(x+n)}}{1 - q^2}}{[x+n]_{q^2}^2} \right] \\ &= -\ln q - 4 \frac{(\ln q)^2}{1 - q^2} q^{2x} (\ln(1 - q^{2n})) \\ &\quad + 4 \frac{(\ln q)^2}{1 - q^2} \left[\frac{q^{2x} [x]_{q^2} + \frac{q^{4x}}{1 - q^2}}{[x]_{q^2}^2} + \dots \right. \\ &\quad \left. + \frac{q^{2(x+n)} [x+n]_{q^2} + \frac{q^{4(x+n)}}{1 - q^2}}{[x+n]_{q^2}^2} \right]. \end{aligned}$$

Then

$$\left(\ln \left((1 - q)^{[x]_q - x} \Gamma_{n,q}(x) \right) \right)'' > 0.$$

This completes the proof.

References

- [1] R. Askey, *The q-Gamma and q-Beta Functions*. Applicable Anal (1978).
- [2] H. Bohr and J. Mollerup, *Laerebog i matematisk Analyse*. Kopenhagen (1922), Vol. III, PP; 149-164.
- [3] K. Brahim, Yosr Sidomou, *On Some Symmetric q-Special Functions*. LE MATHEMATICHE, Vol. LXVIII(2013)-Fasc.II, pp.107-122.

- [4] Detlef Laugwitz and Bernd Rodewald, *A simple characterization of the gamma function*. Amer. Math. Monthly, 94(1987), 534-536.
- [5] Hedi Elmonser ,K. Brahim , Ahmed Fitouhi, *Relationship between characterizations of the q -Gammafunction*. Journal of Inequalities and Special Functions Volume 3 Issue 4(2012), Pages 50-58.
- [6] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd Edition, (2004), Encyclopedia of Mathematics and Its Applications, 96, Cambridge University Press, Cambridge.
- [7] F. H. Jackson, *On a q -Definite Integrals*. Quarterly Journal of Pure and Applied Mathematics 41, (1910), 193-203.
- [8] V. G. Kac and P. Cheung, *Quantum Calculus*, Universitext, Springer-Verlag, New York, (2002).
- [9] H. T Koelink and T. H. Koornwinder, *q -Special Functions*, a Tutorial, in deformation theory and quantum groups with applications to mathematical physics, Contemp. Math. 134, Editors: M. Gerstenhaber and J. Stasheff, J Amer. Math. Soc., Providence, (1992), 141-142.
- [10] J. Thomae, *Beitrage zur Theorie der durch die Heinesche Reihe*, J. reine angew. Math, (70) pp 258-281, 1869.
- [11] Yuan-Yuan Shen, *On characterizations of the Gamma Function*, Mathamatical Association of America, Vol. 68, No. 4(Oct, (1995), pp. 301-305.