

# 1 AGGREGATION OF BASIC REGULAR BLOOD ELEMENTS IN

## 2 CALVES DURING THE PHASE OF MILK FEEDING

### 3 ABSTRACT

4• The aim was to find out aggregation activity of regular blood elements of calves  
5 during the phase of milk feeding. The study was conducted in Kolos farm of  
6 Fatezh district in Kursk region, Russia, spring of 2014. The study used 39  
7 calves of **black and white breed**, taken into investigation on the 11th day of  
8 life. Examination was made on the 11th, 15th, 20th, 25th and 30th days of  
9 calves' life with the usage of biochemical, hematological and statistical methods  
10 of investigation. We estimated the intensity of lipids' peroxidation in plasma,  
11 aggregation of erythrocytes, platelets and neutrophils. During the phase of milk  
12 feeding the calves were noted to have a tendency to the increase of spontaneous  
13 aggregation of erythrocytes. It could be judged by a light tendency for the  
14 increase of summary quantity of erythrocytes in an aggregate, quantity rise of  
15 aggregates themselves and number lowering of disaggregated erythrocytes. All  
16 the calves during milk feeding were noted to have a tendency to strengthening  
17 of platelets' aggregation. So, on the 11th day of life their period of platelets'  
18 aggregation development under collagen impact was equal to  $30.7 \pm 0.12s$ ,  
19 decreasing to some extent during investigation. Analogical platelets'  
20 aggregation state of healthy animals was noted for adenosine diphosphate (to  
21 the end of the phase  $38.1 \pm 0.15s$ ) and ristomicin (to the end of the phase  
22  $46.2 \pm 0.17s$ ). In later period there developed thrombin and adrenaline platelets'  
23 aggregation, also having a tendency to light acceleration during investigation  
24 and being equal to its end to  $51.3 \pm 0.18s$  and  $98.0 \pm 0.34s$ , respectively. During  
25 the phase of milk feeding the calves were also noted to have a little tendency to  
26 strengthening of neutrophils' aggregation. So, their neutrophils' aggregation  
27 during investigation rose with lectin on 4.6%, with concanavalin A - on 6.4%,

28 with phytohemagglutinin - on 3.2%. During the phase of milk feeding the calves  
29 were noted to have a little tendency to strengthening of lipids' peroxidation in  
30 plasma. The calves of the age between 11 to 30 days of life were found to have  
31 little strengthening of regular blood elements' aggregation.

32 **Key words:** phase of milk feeding, calves, aggregation, erythrocytes, platelets,  
33 white blood cells.

## 34 **1. INTRODUCTION**

35 Blood, consisting of regular elements and plasma, continuously circulates along  
36 vessels in a living body [1]. It provides gas exchange and delivery of nutrients  
37 and biologically active substances to tissues [2,3], and also removal of  
38 metabolic waste products out of them [4,5]. The efficiency of hemocirculation,  
39 especially in microcirculation system, mostly depends on regular blood  
40 elements' aggregation [6,7]. Its evidence is under constant control from the side  
41 of a vascular wall [8,9]. It was noted that surplus aggregation of erythrocytes,  
42 platelets and leucocytes can inhibit metabolic processes in a body [10,11]. In  
43 this connection we find to be very urgent the estimation of the degree of regular  
44 blood elements' aggregation in calves at the beginning of their ontogenesis - in  
45 the phase of milk feeding [12]. Given investigations are important for both  
46 fundamental science and practice as abnormalities in the processes of  
47 aggregation and disaggregation in blood play essential role in pathogenesis of  
48 many diseases [13,14]. Both physiology of animals and veterinary science need  
49 precisely adjusted normative indices of basic regular blood elements'  
50 aggregation [15]. These norms are necessary for the estimation of dynamics of

51 cattle state, including calves of milk feeding in case of application of various  
52 impacts on their bodies [16].

53

54 The following aim was put in our investigation - to find out aggregation activity  
55 of regular blood elements in calves during the phase of milk feeding.

## 56 **2. MATERIALS AND METHODS**

57 The research was conducted in strict accordance with ethical principles  
58 established by the European Convent on protection of the vertebrata used for  
59 experimental and other scientific purposes (adopted in Strasbourg in March 18,  
60 1986, and confirmed in Strasbourg in June 15, 2006) and approved by the local  
61 ethic committee of Kursk Institute of Social Education, branch of Russian State  
62 Social University (Record №12 dated December 3, 2015) and the local ethic  
63 committee of All-Russian SII of Physiology, Biochemistry and Animals'  
64 feeding (Record №11, dated December 4, 2015).

65

66 The study used 39 calves of black-many coloured breed, taken into investigation  
67 on the 11th day of life. All the calves were got in autumn. The animals were  
68 kept in Kursk region (Central Russia) in calf-sheds of the farm “Kolos” without  
69 special heating. They received whole milk for feeding in quantity which was  
70 necessary for them. Examination was made five times during the phase of milk  
71 feeding - on the 11th, 15th, 20th, 25th and 30th days of life.

72

73 The activity of the processes of lipids' peroxidation (LPO) in plasma was  
74 estimated by the content of thiobarbituric acid (TBA)-active products with the  
75 help of a set "Agat-Med" and acyl hydroperoxides (AHP). Antioxidant potential  
76 of liquid part of blood was defined by its antioxidant activity (AOA) [17].

77

78 The evidence of erythrocytes' aggregation was defined with the help of a light  
79 microscope in Gorjaev's box. We registered the quantity of erythrocytes'  
80 aggregates, the number of aggregated and disaggregated erythrocytes [18].

81

82 Platelets' aggregation (AP) was estimated with the help of visual micromethod  
83 of AP estimation [19] with the usage of adenosine diphosphate (ADP) ( $0.5 \times 10^{-4}$   
84 M), collagen (dilution 1:2 of basic suspension), thrombin (0.125 un/ml),  
85 ristomicin (0.8 mg/ml) and adrenaline ( $5.0 \times 10^{-6}$  M) in rich in platelets plasma  
86 with standardized platelets' quantity  $200 \times 10^9$  tr. Activity of neutrophils'  
87 aggregation was estimated with the help of a photoelectrocolorimeter. As  
88 inductors we used lectin of wheat foetus in a dose of 32 mkg/ml, concanavalinA  
89 - 32 mkg/ml and phytohemagglutinin - 32 mkg/ml.

90

91 Statistical processing of received data was made with the help of a programme  
92 package "Statistics for Windows v. 6.0", "Microsoft Excel". The results were  
93 processed by Student's criterion (t). Differences in data were considered reliable

94 in case of  $p < 0.05$ .

95

### 96 3. RESULTS AND DISCUSSION

97 Examined calves were noted to have small LPO activity of plasma with a slight  
98 tendency to strengthening during the period of investigation - the content of  
99 AHP in it rose from  $1.44 \pm 0.17$  D<sub>233</sub>/1ml to  $1.47 \pm 0.25$  D<sub>233</sub>/1ml, TBA-active  
100 products - from  $3.59 \pm 0.15$  umol/l to  $3.64 \pm 0.28$  umol/l. It was accompanied by a  
101 tendency to some weakening of plasma AOA from  $33.5 \pm 0.38\%$  on the 11th day  
102 of life to  $33.0 \pm 0.34\%$  on the 30th day of calves' life.

103

104 During the phase of milk feeding the calves were noted to have unexpressed  
105 tendency in the rise of spontaneous erythrocytes' aggregation. It could be judged  
106 by a slight tendency to the increase of summary erythrocytes' quantity in an  
107 aggregate (on 1.9%), quantity rise of aggregates themselves (on 2.4%) and  
108 number lowering of disaggregated erythrocytes (on 2.2%) (table).

109

110 All the calves during milk feeding were noted to have a tendency to  
111 strengthening of platelets' aggregation. So, on the 11th day of life their period of  
112 AP development under the impact of collagen was equal to  $30.7 \pm 0.12$ s,  
113 decreasing to some extent during investigation. Analogical AP state of healthy  
114 animals was noted for ADP (to the end of the phase -  $38.1 \pm 0.15$ s) and  
115 ristomicin (to the end of the phase -  $46.2 \pm 0.17$ s). In later period there developed

116 thrombin and adrenaline AP, having also a tendency to light acceleration during  
117 investigation and being equal to its end to  $51.3 \pm 0.18s$  and  $98.0 \pm 0.34s$ ,  
118 respectively (table).

119 During the phase of milk feeding the calves were also noted to have a small  
120 tendency to strengthening of neutrophils' aggregation. So, during investigation  
121 their neutrophils' aggregation rose with lectin on 4.6%, with concanavalinA - on  
122 6.4%, with phytohemagglutinin - on 3.2% (table).

123 The consumption of milk and beef by the population of the planet increases, and  
124 it dictates the necessity of constant development of this agricultural branch. It  
125 can be achieved in the result of continuation of active scientific investigations of  
126 cattle physiology [15]. In this connection special significance is given to  
127 investigations of calves' blood physiology at the beginning of ontogenesis [20].  
128 So, great attention is devoted to investigations of calves preparing for the  
129 beginning of vegetable feeds' consumption. In our work the calves were noted  
130 to have gradual strengthening of plasma AOA at the age between the 11th and  
131 30th days of life which was accompanied by gradual weakening of LPO  
132 activity. Found facts were supported by the results of earlier investigations [21].  
133 It is known that intensity of freely-radical processes in plasma influences  
134 significantly the morphofunctional state of erythrocytes, platelets and  
135 leucocytes [22,23]. It can explain the slight ability in aggregation of basic  
136 regular blood elements in calves during the phase of milk feeding.

137

138 In the carried-out work special attention is paid to aggregation of uniform  
139 elements of blood. Intra vascular formation of units and success of  
140 microcirculation in many respects depends on her level. In this regard processes  
141 of a metabolism and intensity of growth at animals depend on activity of  
142 aggregation of uniform elements of blood.

143

144 It is evident that the great quantity of electronegative proteins on the surface of  
145 erythrocytes [24] mostly lies in the basis of small activity of erythrocytes'  
146 aggregation in calves at milk feeding. High control over generation of oxygen  
147 active forms in calves provides minimization of oxidative damages of  
148 membrane erythrocyte proteins and globular plasma proteins, participating in  
149 aggregation [25]. In this connection we can consider that the phase of milk  
150 feeding of calves is characterized by optimum of metabolic and receptor  
151 processes in erythrocytes. Received estimation results of erythrocytes'  
152 aggregation are confirmed by the single work containing information about the  
153 tendency to its strengthening in calves of the given age [26]. We should  
154 compare received results with literature data with great caution as in the  
155 previous work the groups were mixed, as far as breed is concerned, but calves  
156 of Simmental breed prevailed. Besides, they were got in autumn, and it also  
157 makes comparison of results difficult.

158

159 Noted in calves during the phase of milk feeding tendency to strengthening of  
160 platelets' aggregative activity is connected with activity increase of their  
161 receptors and postreceptor mechanisms of aggregation. Evidently, in calves'  
162 blood at the age of 11-30 days gradually rises concentration of von Willebrand  
163 Factor - cofactor of platelets' adhesion. It is accompanied by weak number  
164 increase of receptors to it - (GPIb) on platelets' surface. It was pointed in calves  
165 by a tendency to decrease of AP period in response to ristomicin. Found AP  
166 dynamics in response to strong and weak agonists of aggregation can be  
167 explained by physiologically approved activity changes of platelet  
168 phospholipase A<sub>2</sub> and C. They provide functioning of thromboxan and  
169 phosphoinositol ways of platelets' activation [27,28]. In literature there is rather  
170 poor information about platelets' activity in calves of milk feeding [29]. In spite  
171 of the fact that famous sources confirm that calves of milk feeding have a  
172 tendency to strengthening of platelets' aggregation, comparison of these results  
173 with received ones should be done with great caution. It is connected with the  
174 fact that experimental calves in previous investigations were kept in Central  
175 Russia in calf-sheds with special heating, and they received substitutes of whole  
176 milk and fodder concentrated products.

177

178 It is known that activity of neutrophils' aggregation in mammals is provided by  
179 locuses' quantity in their glycoprotein receptors' composition. These receptors  
180 can connect lectins [30]. It is firmly established that phytohemagglutinin can



181 mostly interact with parts of bD-galactose of glycoproteins, lectin of wheat  
182 foetus - with N-acetyl-D-glycosamin и N-acetyl-neuraminic (sialic) acid, and  
183 concanavalin A – with N-glycans containing mannose [11]. That is why, the  
184 state of lectin stimulated neutrophils' aggregation of calves is defined by the  
185 expression level of receptors' adhesion. These receptors have such parts in their  
186 composition. Taking it into consideration, we can consider that found tendency  
187 to growth of neutrophils' aggregation at calves' age of 11-30 days was,  
188 evidently, connected with the rise of sensitivity and density of leucocytes'  
189 glycoprotein receptors. It happens simultaneously with changing of their  
190 composition. Gradual strengthening of lectin - and concanavalin A - induced  
191 neutrophils' aggregation in experimental calves was provided by expression  
192 increase of adhesion receptors on their surface and by some growth of areas  
193 containing N-acetyl-D-glucosamine, N-acetyl-neuraminic acid and mannose.  
194 Strengthening increase of aggregation induced by phytohemagglutinin in calves  
195 between the 11th and the 30th days of life was provided by a tendency to the  
196 rise of areas of glycoproteins, containing bD-galactose [11], in their neutrophils'  
197 receptors. Neutrophils' aggregation was not studied earlier on productive  
198 animals and, moreover, on calves. With the help of available literature sources  
199 containing information about investigations of a human being, it becomes clear  
200 that the role of receptor mechanisms in its realization is great, and that it can be  
201 quickly damaged in case of unfavorable environmental and metabolic  
202 conditions [11,27].

203

204 Noted strengthening of aggregative activity of erythrocytes, platelets and  
205 neutrophils in calves during the phase of milk feeding was mostly caused by  
206 processes of growth and strengthening of environmental impacts on their  
207 background [21]. Sufficient activity of adaptive mechanisms in these conditions  
208 keeps the balance of aggregation and disaggregation in calves' blood on the  
209 level necessary for optimum of blood supply of internal organs [25].

210

#### 211 **4. CONCLUSION**

212 During the phase of milk feeding the calves were noted to have a slight  
213 tendency to strengthening of lipids' peroxidation in plasma. The calves at the  
214 age of 11-30 days were found to have little strengthening of regular blood  
215 elements' aggregation.

#### 216 **REFERENCES**

- 217 1. Medvedev IN, Zavalishina SYu. Platelet Activity in Patients With Third  
218 Degree Arterial Hypertension and Metabolic Syndrome. *Kardiologiya*.  
219 2016;56(1):48.
- 220 2. Medvedev IN, Gromnatskii NI, Golikov BM, Al'- Zuraiki EM, Li VI.  
221 Effects of lisinopril on platelet aggregation in patients with arterial  
222 hypertension with metabolic syndrome). *Kardiologiya*. 2004;44(10):57-59.
- 223 3. Medvedev IN, Gromnatskii NI, Mokhamed A.-ZE. Comparative  
224 Assessment of Effects of Qadropiril and Enalapril on Intravascular Activity

- 225 of Platelets in Hypertensive Patients With Metabolic Syndrome.  
226 Kardiologiya. 2004;44(12):44-46.
- 227 4. Medvedev IN, Gromnatskii NI, Volobuev IV, Osipova VM, Dement'ev VI,  
228 Storozhenko MV. Thrombocytic hemostasis in hypertensive patients with  
229 metabolic syndrome and its correction with lovastatin). Klinicheskaya  
230 meditsina. 2004;82(10):37-41.
- 231 5. Simonenko VB, Medvedev IN, Tolmachev VV. Comparative evaluation of  
232 the influence of sulfhydryl and phosphate ACE inhibitors on thrombocyte  
233 aggregation in patients suffering from arterial hypertension with metabolic  
234 syndrome. Klinicheskaya meditsina. 2007;85(4):24-27.
- 235 6. Medvedev IN. A comparative analysis of normodipin and spirapril effects  
236 on intravascular activity of platelets in patients with metabolic syndrome.  
237 Terapevticheskii Arkhiv. 2007;79(10):25-27.
- 238 7. Kutafina NV, Medvedev IN. Platelet Aggregation in Clinically Healthy  
239 Persons of the Second Coming-of-Age Living in the Kursk Oblast.  
240 Advances in Gerontology.2015;5(4):267-270.
- 241 8. Medvedev IN, Skoryatina IA. Aggregation properties of blood cells and  
242 vascular control over them in patients with arterial hypertension and  
243 dyslipidemia. Russian Journal of Cardiology. 2015;4(120):18-22.
- 244 9. Simonenko VB, Medvedev IN, Mezentseva NI, Tolmachev VV. The  
245 antiaggregation activity of the vascular wall in patients suffering from

- 246 arterial hypertension with metabolic syndrome. *Klinicheskaiia meditsina*.  
247 2007;85(7):28-30.
- 248 10. Medvedev IN, Gamolina OV. Lisinopril effects on platelet activity in  
249 patients with arterial hypertension and impaired glucose tolerance. *Russian*  
250 *Journal of Cardiology*. 2008;3:45-48.
- 251 11. Medvedev IN, Skoryatina IA. The aggregation capacity of neutrophils in  
252 patients with arterial hypertension and dyslipidemia treated with fluvastatin.  
253 *Klinicheskaiia meditsina*. 2015;93(1):66-70.
- 254 12. Solovyova LP, Gorbunova NP, Rybakova GK, Kalysh TV, Barmin SV.  
255 Comparative analysis hemostatic properties of blood in calves and pigs  
256 lacto-vegetarian power. *Modern problems of science and education*. 2015;  
257 6; URL: <http://www.science-education.ru/130-23582>
- 258 13. Medvedev IN, Skoriatina IA. Dynamics of microrheologic properties of  
259 erythrocytes in patients with arterial hypertension and dyslipidemia treated  
260 with atorvastatin. *Klinicheskaiia meditsina*. 2012;90(6):42-45.
- 261 14. Medvedev IN, Lapshina EV, Zavalishina SYu. Experimental methods for  
262 clinical practice: Activity of platelet hemostasis in children with spinal  
263 deformities. *Bulletin of Experimental Biology and Medicine*.  
264 2010;149(5):645-646.
- 265 15. Medvedev IN. Vascular-platelet interaction in pregnant cows. *Bulg. J.*  
266 *Agric. Sci.*2017;23(2):310-314.

- 267 16. Solovyova LP, Gorbunova NP, Rybakova GK, Kalysh TV, Barmin SV.  
268 State functional mechanisms of hemostasis calves pigs and power plant after  
269 traffic stress. Modern problems of science and education. 2015; 2; URL:  
270 <http://www.science-education.ru/131-23568>
- 271 17. Volchegorskij IA, Dolgushin II, Kolesnikov OL, Cejlikman VJe.  
272 Experimental modeling and laboratory assessment of adaptive reactions of  
273 the organism. Cheljabinsk, 2000:167.
- 274 18. Medvedev IN, Maksimov VI, Parakhnevich AV, Zavalishina SYu, Kutafina  
275 NV. Rapid assessment of aggregation abilities and surface properties of  
276 platelets and red blood cells. International Journal of Pharma and Bio  
277 Sciences. 2016 April;7(2):(B)793-797.
- 278 19. Medvedev IN, Savchenko AP, Zavalishina SYu, Krasnova EG, Kumova  
279 TA, Gamolina OV, Skoryatina IA, Fadeeva TS. Methodology of blood  
280 rheology assessment in various clinical situations. Russian Journal of  
281 Cardiology. 2009;5:42-45.
- 282 20. Krasnova EG, Kutafina NV. Basics platelet function. Veterinary, zootechny  
283 and biotechnology. 2015;8:6-18.
- 284 21. Zavalishina SYu. State of the Hemostatic System in Ironn Deficient  
285 Newborn Calves. Russian Agricultural Sciences. 2013;39(4):350-353.
- 286 22. Medvedev IN. Dynamics of violations of intravascular platelet activity in  
287 rats during the formation of metabolic syndrome using fructose models.  
288 Problems of nutrition. 2016;85(1):42-46.

- 289 23. Simonenko VB, Medvedev IN, Tolmachev VV. Effect of irbesartan of the  
290 function of hemocoagulative component of hemostasis in patients with  
291 arterial hypertension during metabolic syndrome. *Klinicheskaiia meditsina*.  
292 2010;88(6):27-30.
- 293 24. Medvedev IN, Skoryatina IA. Erythrocyte aggregation in patients with  
294 arterial hypertension and dyslipidemia treated with pravastatin.  
295 *Klinicheskaiia meditsina*. 2014;92(11):34-38.
- 296 25. Simonenko VB, Medvedev IN, Kumova TA. Pathogenetic aspects of  
297 hypertension in case of metabolic syndrome. *Voenno-meditsinskiizhurnal*.  
298 2010;331(9):41-44.
- 299 26. Belova TA. Functional features erythrocytes at the healthy newborn calves.  
300 *International bulletin of veterinary medicine*. 2010;4:52-55.
- 301 27. Medvedev IN, Skoryatina IA. Fluvastatin effects on blood cell aggregation  
302 in patients with arterial hypertension and dyslipidemia. *Cardiovascular*  
303 *Therapy and Prevention*. 2013;12(2):18-24.
- 304 28. Medvedev IN, Kumova TA, Gamolina OV. Renin-angiotensis system role  
305 in arterial hypertension development. *Russian Journal of Cardiology*.  
306 2009;4:82-84.
- 307 29. Zavalishina SYu. Hemostatic activity of thrombocytes in calves during the  
308 phase of milk feeding. *Agricultural Biology*. 2013;4:105-109.

309 30. Medvedev IN, Skoryatina IA. Pravastatin in correction of vessel wall  
 310 antiplatelet control over the blood cells in patients with arterial hypertension  
 311 and dyslipidemia. Cardiovascular therapy and prevention. 2014;13(6):18-22.

312  
 313  
 314  
 315

**Table. The activity of the processes of lipids' peroxidation in plasma and aggregation of blood elements in calves of dairy nutrition**

Registered parameters	Age of calves (n=39, M±m)				
	11 days	15days	20 days	25 days	30 days
acyl hydroperoxides, D <sub>233</sub> /1ml	1.44±0.17	1.46±0.12 P<95%	1.47±0.20 P<95%	1.47±0.15 P<95%	1.49±0.25 P<95%
TBA-active products, umol/l	3.59±0.15	3.63±0.22 P<95%	3.60±0.26 P<95%	3.62±0.19 P<95%	3.64±0.28 P<95%
AOA, %	33.5±0.38	33.3±0.36 P<95%	33.1±0.34 P<95%	32.9±0.29 P<95%	32.4±0.32 P<95%
sum of all the erythrocytes in an aggregate	40.1±0.19	40.2±0.24 P<95%	40.4±0.29 P<95%	40.6±0.25 P<95%	40.9±0.32 P<95%
quantity of aggregates	8.2±0.12	8.2±0.10 P<95%	8.3±0.16 P<95%	8.4±0.19 P<95%	8.4±0.11 P<95%
quantity of free erythrocytes	245.7±2.19	244.2±2.25 P<95%	241.8±2.01 P<95%	242.0±1.90 P<95%	240.4±2.46 P<95%
AP with ADP, s	39.2±0.16	39.0±0.12 P<95%	38.7±0.13 P<95%	38.4±0.10 P<95%	38.1±0.15 P<95%
AP with collagen, s	30.7±0.12	30.5±0.10 P<95%	30.3±0.09 P<95%	30.1±0.11 P<95%	29.7±0.14 P<95%
APwith thrombin, s	52.7±0.15	52.6±0.10 P<95%	52.2±0.16 P<95%	51.7±0.10 P<95%	51.3±0.18 P<95%
APwith ristomicin, s	47.5±0.12	47.2±0.16 P<95%	46.9±0.22 P<95%	46.6±0.26 P<95%	46.2±0.17 P<95%
APwith epinephrine, s	97.8±0.42	97.4±0.36 P<95%	97.1±0.32 P<95%	98.5±0.45 P<95%	98.0±0.34 P<95%
Aggregation neutrophils with lectin, %	14.5±0.16	14.5±0.17 P<95%	14.7±0.15 P<95%	14.9±0.26 P<95%	15.2±0.22 P<95%
Aggregation neutrophils with concanavalin A, %	14.5±0.10	14.6±0.12 P<95%	14.9±0.16 P<95%	15.1±0.11 P<95%	15.5±0.13 P<95%
Aggregation neutrophils with phytohemagglutinin, %	27.1±0.19	27.2±0.23 P<95%	27.4±0.14 P<95%	27.8±0.26 P<95%	28.0±0.21 P<95%

316  
 317  
 318  
 319

Note: there was found no reliability of indices' changes in comparison with their level at the age of 11 days. We list the values of P – possibility of unmistakable prognosis.