Original Research Article Assessment of sediments pollution by trace metals in the Moloundou swamp, southeast Cameroon

4

5 Abstract

6 The present study aimed to assess the selected trace metal pollutants in the sediments of the 7 Moloundou swamp. Sediments from typical swamp around Moloundou area, southeast Cameroon 8 were collected from various depths of sediment profile (from surface to 120 cm depth). Five sites were chosen for this study, Fe, Co, Ni, Cr, Zn, Cu, and Pb were measured in the sediment. Different trace 9 metals indices (enrichment factor, contamination factor, degree of contamination, ecological risk 10 assessment and geo-accumulation index) were calculated. The results showed that all the swamp is 11 slightly polluted and core 3 is the most polluted. Heavy metal indices give some indication for the 12 pollution of sediments of all cores with Cu. The enrichment factor showed that the source of those 13 14 metals in Moloundou swamp was from natural (Fe, Ni, Zn and Pb) and anthropogenic sources (Cu, Cr 15 and Co). The degree of contamination and contamination factor showed low values along the cores, 16 like the ecological assessment and pollution load index. The geo-accumulation index showed that sediments are more polluted with Cu, Cr and Co. It is urgent to control anthropogenic waste in order to 17 18 avoid probable pollution in this zone.

19 Keywords: Heavy metals indices; Moloundou swamp; sediments; pollution.

20 1. INTRODUCTION

Swamps are considered to play a significant role in catchment hydrodynamics by capturing and storing 21 22 rainwater and then releasing that water slowly as base flow in dry periods [1]. The magnitude of the 23 'sponge' phenomenon of upland swamps is dependent on the volume of water in the sediment profile, 24 which is influenced by the antecedent groundwater level, the magnitude and duration of rainfall, and the presence of any springs or seeps discharging from regional aquifers [2, 3]. This means that the 25 ability of swamps to act as water storage reservoirs and flood attenuators will change under different 26 27 hydrological conditions. Numerous pollutants adsorb to sediments that accumulate at the bottom of 28 reservoirs. These sediments accumulate over the time and can be considered as new pollutant 29 sources to the overlying water [4, 5]. The forms and types of pollutants usually vary from sediments to 30 others because bonding forces vary with respect to grain size [6, 7] which results in different release 31 times and release potentials. Trace metal contamination is a serious threat in aquatic systems due to 32 their toxicity, abundance, persistence in the environment and subsequent accumulation in water 33 habitats [8, 9, 10]. It has been recently reported that contamination of water sources from trace metals 34 includes geological weathering and erosion [5, 11], atmospheric deposition [12], disposal of treated 35 and untreated liquid effluents [13], trace metals containing fertilizers and pesticides [14, 15] and chemicals originating from various urban, industrial and agricultural activities [16, 17, 18]. Moreover, 36 37 few studies have suggested that sediment quality could serve as an indicator for the pollution levels and sediments could act as a screening tool to fingerprint the historical as well as the recent 38 39 contamination in the surrounding environment [19, 20, 21]. In this context, the main aims of the 40 present study were to determine the levels of trace metals in the sediments from the sediments of 41 Moloundou swamp in order to identify their naturally enriched or anthropogenic sources using heavy metals indices, as well as to assess the environmental risk of heavy metal in the investigated area by 42 43 comparing the acquired metal values with Sediment Quality Guidelines (SQGs). This spatial survey is also useful to assess pollution in the Moloundou swamp and to provide basic information for the 44 45 judgment of environmental health risks and management of urgent environmental pollution issues in 46 the sediments.

48 2. EXPERIMENTAL METHODS

49 **2.1 Study area**

50 Moloundou Swamp (01°58'N-02°02'N, 15°25'E-15°29' E), is located at about 28 km Northeast of the 51 Moloundou district, in the East region of Cameroon (Fig. 1). The study area is made up swampy 52 hydromorphic soils. The vegetation cover corresponds to a dense forest. It is composed of Cryogenian 53 tillite and schisto-limestone [22, 23]. Input of materials in this swamp is considered to have been 54 derived from the surrounding areas and transported by surface runoff and rivers which was controlled 55 primarily by precipitation, and established favorable conditions for the development of lacustrine 56 sediments.



Fig. 1. Investigated area and sampling sites inside the sediments of Moloundou swamp

57 2.2 Sampling and analysis of sediments

58 A total of 48 sediment samples were collected from all the selected sites using standard protocol [5]. 59 The sediment samples were taken at five different sites (1; 2; 3; 4 and 5) inside the swamp using a 60 core sampler (raft of wood placed in the middle of the sampling location with PVC pipe), and at each site, composite sediment samples were prepared. In the laboratory, samples were air-dried to reduce 61 the water content. Dried samples were analyzed in 48 representative samples (at 10 cm intervals) for 62 trace metals by inductively coupled plasma-atomic emission (ICP-AES), using the pulp at ALS Global 63 Group (Vancouver, Canada). The samples were pulverized to obtain a homogeneous sample out of 64 which 50-60 g was obtained for the analyses. 0.2 g of rock powder was fused with 1.5 g LiBO₂ and 65 then dissolved in 100 ml 5 % HNO₃. Analytical uncertainties vary from 0.01 % to 6 %. 66

67

68 2.3 Heavy Metals Indices

69 2.3.1 Enrichment Factor (EF)

Iron (Fe) was chosen as a stationary reference element to perform this calculation [24]. The EF values
 2 indicate that the metal is entirely from crustal materials or natural processes; whereas EF values >

72 2 reveal that the sources are more likely to be anthropogenic [25].

73
$$EF = \frac{[M]Sample/[Fe]sample}{[M]Background/[Fe]background}$$

where, M is the metal concentration. The background value is that of average shale [26]. About six
categories are recognized for FE: ≤ 1 background concentration, 1-2 depletion to minimal enrichment,
2-5 moderate enrichment, 5-20 significant enrichment, 20-40 very high enrichment and > 40 extremely
high enrichment [27].

78 2.3.2 Contamination Factor (CF)

The CF is the ratio calculated by dividing the concentration of each metal in the sediment by the baseline or background value [28]. Contamination Factor (CF) = $C_{metal} / C_{background}$, the following expressions are used to describe the contamination factor: CF < 1 (low contamination factor); $1 \le CF <$ 3 (moderate contamination factors); $3 \le CF < 6$ (considerable contamination factors) and CF ≥ 6 (very high contamination factor).

84 2.3.3 Degree of Contamination (DC)

85 The Degree of Contamination (DC) is the sum of all contamination factors for a given site [29]:

$$DC = \sum_{i=1}^{n} CFi$$

where CF is the single contamination factor and n is the amount of the elements present. DC < n, would indicate low degree of contamination; $n \le DC < 2n$, moderate degree of contamination; $2n \le Dc$ < 4n, considerable degree of contamination; and Dc > 4n, very high degree of contamination. For the studied heavy metals, n=7.

90 2.3.4 Ecological Risk Assessment

The ecological risk assessment was carried out potential ecological risk index (RI) for this study. The potential ecological risk index (RI) of the heavy metals is known as the sum of the risk factors and developed for six toxic metals using equations of Hakanson [29] and Zhu et al. [30].

94
$$RI = \sum_{1}^{n} Er$$
 and $Er = Tr \times CF$

where Er is the single index of ecological risk factor, and n is the amount of the heavy metal class, Tr
toxic response factor suggested by Hakanson [29] for six metals Cr (5), Co (5), Cu (5), Pb (5), Ni (5),
Zn (1). Er and RI express the potential ecological risk factor of individual and multiple metals,
respectively. The following expressions was used for the potential ecological risk factor: Er < 40, low

potential ecological risk; $40 \le \text{Er} < 80$, moderate potential ecological risk; $80 \le \text{Er} < 160$, significant potential ecological risk; $160 \le \text{Er} < 320$, high potential ecological risk; and $\text{Er} \ge 320$, very high ecological risk. Furthermore, the potential ecological risk index: RI < 150, low ecological risk; $150 \le \text{RI}$ < 300, moderate ecological risk; $300 \le \text{RI} < 600$, significant ecological risk; and RI > 600, very high ecological risk [29].

104 The pollution load index (PLI) of a single site is the root of number (n) of multiplied together 105 contamination factor (CF) values.

 $PLI = \sqrt[n]{(CF1 \times CF2 \times CF3 \times \dots \times CFn)}$

where, n is the number of metals and CF is the contamination factor. A PLI = 0, indicates perfection; a
 value < 1 indicates no pollution, and values > 1 is polluted area [31].

108 2.3.5 Geo-Accumulation Index (Igeo)

109 The index of geo-accumulation (Igeo) was firstly defined by Muller [32] to determine and define the 110 metal contamination in sediments by comparing current concentrations with preindustrial levels [33].

$$Igeo = Log2 (Cn/1.5Bn)$$

111 where, Cn is the concentration of metals (mg/kg) in sediments, Bn is the geochemical background 112 value (mg/kg) in average shale of element n and 1.5 is the background matrix correction due to 113 anthropogenic influences. The geo-accumulation index (Igeo) was distinguished into seven classes by 114 Buccolieri et al., [34]: Igeo \leq 0, class 0 (unpolluted); 0 < Igeo \leq 1, class 1 (unpolluted to moderately 115 polluted); 1 < Igeo \leq 2, class 2 (moderately polluted); 2 < Igeo \leq 3, class 3 (moderately to strongly 116 polluted); 3 < Igeo \leq 4, class 4 (strongly polluted); 4 < Igeo \leq 5, class 5 (strongly to extremely polluted); 117 and Igeo > 5, class 6 (extremely polluted).

118

119 3. RESULTS AND DISCUSSION

120 3.1 Heavy Metal Pollution

The basic descriptive statistical values and spatial distributional patterns of the studied trace metals 121 122 are presented in Table 1 and Fig. 2. On the average basis, the metals follow a decreasing 123 concentration order Fe > Cu > Cr > Zn > Ni > Co > Pb. Comparison of the average concentrations of heavy metals in the different samples shows that the average Fe concentration is higher (5788 mg/kg) 124 125 than that of other metals. This average of Fe concentration in Moloundou sediments is less than the 126 Average Shale (46700 mg/kg) [26] and UCC [35] reference values. On the other hand, the average 127 concentrations of the other metals in this study are higher than the reference values of Average Shale 128 [26] and UCC [35], These abnormal values can be attributed to the pollutant load provided by the various discharges from agricultural and artisanal mining activities. The maximum concentrations of 129 130 the different elements analysed are recorded in the cores 3 and 5 samples, in particular for Fe (7133 131 mg/kg), Cu (206.67 mg/kg), Pb (23.33 mg/kg), Ni (76.33), Cr (191.33 mg/kg) and Zn (134.67 mg/kg). 132 The high levels of these metals could be attributed to human activities such as ores extraction and 133 agricultural activities inside and around the Moloundou swamp.

Table 1. Heavy metal concentrations (Mean ± SD) in the sediment samples of Moloundou swamp (mg/kg)

Sites	Fe	Со	Ni	Cr	Zn	Cu	Pb
1 (n = 8)	4993 ± 907	35 ± 4.36	67.66 ± 6.81	180 ± 45.83	116.66 ± 20.82	196.66 ± 81.45	16.7 ± 5.94
2 (n = 10)	6546 ± 1176	33.33 ± 7.23	73.33 ± 5.86	186.66 ± 25.17	126.66 ± 11.55	192.66 ± 57.46	12.6 ± 6.83
3 (n = 12)	7133 ± 830	31.66 ± 3.6	66.66 ± 6.43	180 ± 22.72	126.66 ± 11.02	206.66 ± 47.43	23.33 ± 3.76
4 (n = 8)	5163 ± 1056	39.66 ± 2.08	71.33 ± 5.13	180 ± 40	113 ± 6.08	201.66 ± 74.89	12.5 ± 2.26

5 (n = 10)	5240 ± 392	32 ± 2.89	76.33 ± 5.77	191.33 ± 30	134.66 ± 20.82	197.33 ± 80.83	6.43 ± 4.72
Mean (n = 48)	5788 ± 872	32.88 ± 4.03	70.22 ± 6	183.77 ± 32.74	126 ± 14.06	200.22 ± 68.41	15.48 ± 4.70
UCC values	<mark>46700</mark>	<mark>11</mark>	<mark>19</mark>	<mark>35</mark>	<mark>52</mark>	<mark>14</mark>	<mark>17</mark>
136							

137 3.1.1 Copper (Cu)

138 Copper is an essential micronutrient for aquatic life in freshwaters and sediments but it becomes toxic 139 at higher level. It is released to the environment from natural sources such as volcanic eruptions. 140 decaying vegetation, forest fires, and sea spray up to 50 mg/kg [36, 37] and anthropogenic activities, including municipal and industrial wastewater [36, 38]. After a series of natural processes such as 141 weathering and leaching, the water-borne Cu finally accumulates in the sediment and the quantity of 142 Cu contained in the sediment reflects the degree of pollution of the water body. The average 143 144 concentration values were ranging from 192.67 to 206.67 mg/kg with a mean of 200.22 mg/kg (Table 1; Fig.2), relatively lower than those found in previous studies in the paddy fields sediments from 145 mangrove swamp in Pearl River Estuary, China [15], but higher than those measured in the sediments 146 from Turfy Swamps, Northeastern China [18] and sediments from Simbock Lake in Yaounde, 147 Cameroon [5]. However, possible sources of Cu toxicity in the sediments of the Moloundou swamp 148 149 might include agriculture and artisanal gold exploitation around the study area.



150

151 Fig. 2. Distribution maps of the concentration of selected heavy metals (mg/kg) in sediments of

- 152 Moloundou swamp
- 153 3.1.2 Chromium (Cr)

The toxicity of chromium depends on its degree of oxidation. Hexavalent chromium salts are very toxic [39]. Chromium is more toxic in fresh water than in hard water and also more toxic to invertebrates than to fish [40]. The Cr content in Moloundou swamp is about 180 to 191.33 mg/kg, with an average of 183.77 mg/kg (Table 1; Fig.2), which exceeds the French sediment quality standards set by the ATSDR [36] for Cr (100mg/kg). Compared with previously published results, Chromium values of this study are closed to Simbock Lake sediment in Yaounde, Cameroon [5] and higher than those from mangrove swamp in Pearl River Estuary, China [9] and the coastal sediments of the South Sea of Korea [41]. This high content would be justified by the deposition of atmospheric particles and the leaching of chromium-containing in source rock.

163 3.1.3 Zinc (Zn)

Natural background levels of zinc are usually found up to 100 mg/kg in sediments [42]. The concentrations of Zn in the present study ranged from 113 to 134.67 mg/kg with an average of 126 mg/kg (Table 1; Fig.2), which are lower than those reported for the in sediments from mangrove swamp in Pearl River Estuary, China [9], but higher than sediment of Turfy Swamps, Northeastern China [18] and Simbock Lake sediments in Yaounde, Cameroon [5]. However, these Zn levels were similar to those previously found in the coastal sediments of the South Sea of Korea [41]. This high concentration is of anthropogenic origin, which is the mining practiced in this locality.

171 3.1.4 Nickel (Ni)

172 Nickel can enter the environment as a result of the natural weathering and leaching of rocks [43]. The 173 average concentrations of Nickel (Ni) present in the earth's crust are about 17 to 20 mg/kg [26, 35]. The average concentration values (mg/kg) were ranging from 66.66 to 76.33 with a mean of 70.22 174 175 (Table 1; Fig.2), relatively lower than those found in previous studies in the paddy fields sediments 176 from mangrove swamp in Pearl River Estuary, China [9] but higher than those measured in the sediments from Turfy Swamps, Northeastern China [18] and sediments from Simbock Lake in 177 Yaounde, Cameroon [5]. This concentration could be explained by the pollutant load provided by the 178 179 various residues from mining operations, this is agreed with Biney [44] who found same impacts.

180 3.1.5 Cobalt (Co)

181 The average concentrations of cobalt present in the earth's crust are about 20-25 mg/kg [36]. The 182 concentrations of Co (mg/kg) found in the sediments Moloundou sediments ranged from 31.66 to 39.66 with average of 32.88 (Table 1; Fig.2). Compared with previously published results, Co levels 183 184 were lower than those reported for Tigris river, Turkey [45] and higher than sediment of Turfy Swamps, 185 Northeastern China [18]: Simbock Lake in Yaounde, Cameroon [5] and from mangrove swamp in 186 Pearl River Estuary, China [9]. This high concentration could be linked to the presence of iron and 187 manganese oxyhydroxides in Moloundou sediments because in sediments Co is considered to be a 188 low-mobility element and has a high affinity with iron and manganese oxyhydroxide.

189 3.1.6 Lead (Pb)

190 Lead is a non-essential and toxic element released from natural and anthropogenic activities. Major 191 sources include vehicular emissions, volcanoes, airborne soil particles, forest fires, waste incineration, 192 effluents from leather industry, lead containing paints and pesticides [38, 46, 47]. Natural 193 concentration of Pb in the earth's crust varied from 15 to 20 mg/kg [46]. The current study reported 194 that Pb concentrations (mg/kg) in the sediments ranged from 6.43 to 23.33 with an average of 15.43 195 (Table 1: Fig.2). The comparison between Pb concentrations in the sediments of the Moloundou 196 swamp and other areas showed that Pb levels in these sediments had higher levels than those 197 measured in the sediments of Indus and Pakistan [48] rivers and Turfy Swamps, Northeastern China 198 [18] but lower concentrations than those measured in the sediments from the Nile river, Egypt [49, 50]; 199 soils from El Mahalla El-Kobra area, Gharbia governorate, Egypt [51] and Simbock Lake in Yaounde, 200 Cameroon [5]. Most likely, the concentration of Pb reported in this study originated from both natural 201 and anthropogenic sources which drain out their waste from mining [52, 53]. Moreover, use of 202 fertilizers and pesticides is also a common practice in the catchment areas of which could contribute 203 significantly to the presence of Pb in the swamp.

204 3.2 Heavy Metals Indices

205 3.2.1 Enrichment Factor (EF)

The enrichment factors of heavy metals in Moloundou swamp were as shown in Fig. 3 and Table 2. The sequence of EF in the sediments was Cu > Cr > Co > Zn > Ni > Pb. The enrichment factor of heavy metals in all core sediments is between 5-20 range, indicated significant enrichment and that the source of those metals was from anthropogenic activities. Except Cu enrichment factor values are between very high enrichment (for cores 2, 3 and 5) and extremely enrichment (for cores 1 and 4)

(Fig. 3 and Table 2). It is obvious that Cu is most abundant in all core sediment; this could be 211 212 attributed to agricultural wastes and artisanal gold exploitation.



213

Fig. 3. The enrichment factor of heavy metals in the sediments of Moloundou swamp 214

215	Table 2. The Enrichment Factor (EF) of heavy	metals in the sediment	samples of Moloundou
216	swamp		

2	т	b		

Sitoo	Enrichment Factor (EF)								
Siles	Fe	Zn	Pb	Ni	Cr	Co	Cu		
Core 1	1	11.61	7.89	9.41	18.91	17.41	41.31		
Core 2	1	9.61	4.54	7.78	14.96	12.65	30.87		
Core 3	1	8.82	7.72	6.49	13.23	11.03	30.39		
Core 4	1	10.87	5.71	9.59	18.28	19.09	40.97		
Core 5	1	12.77	2.90	10.11	19.15	15.17	39.50		
Mean (n=48)	1	10.82	6.32	8.42	16.65	14.12	36.28		

217

218 3.2.2 Contamination Factor (CF) and Degree of Contamination (DC)

The contamination factor (CF) used to determine the contamination status of the sediment in the 219 220 present study. According to Hakanson classification [29], CF of all metals in the sediments of 221 Moloundou showed moderate contamination factor except for Pb which showed low and Cu which is considerable (Table 3). The degree of contamination (DC) show moderate degree of contamination (7 222 223 \leq DC < 14) in all cores of the sediments in Moloundou swamp (Table 3 and Fig. 4). The sediments were moderately and considerably contaminated by theses metals due to the influence of external 224 discrete sources like industrial activities, agricultural runoff, and other anthropogenic inputs. 225

226	Table 3. The contamination factor (CF), Pollution Load Index (PLI) and degree of Contamination
227	(DC) of Moloundou swamp

Sitos	Contamination factor (CF)								ыі	
Siles	Fe	Zn	Pb	Ni	Cr	Со	Cu		Г С 1	
Core 1	0.11	1.23	0.84	1.00	2.00	1.84	4.37	11.38	1.08	
Core 2	0.14	1.33	0.63	1.08	2.07	1.75	4.28	11.29	1.10	
Core 3	0.15	1.33	1.17	0.98	2.00	1.67	4.59	11.89	1.20	
Core 4	0.11	1.19	0.63	1.05	2.00	2.09	4.48	11.54	1.07	
Core 5	0.11	1.42	0.32	1.12	2.13	1.68	4.39	11.17	0.98	
Mean (n=48)	0.12	1.33	0.77	1.03	2.04	1.73	4.45	11.48	1.11	



229

230 Fig. 4. The degree of contamination (DC) and pollution load index (PLI) of heavy metals in the

231

sediments of Moloundou swamp

232 **3.2.3 Ecological Risk Assessment**

233 3.2.3.1 Ecological Risk Factor (Er) and Ecological Risk Index (RI)

The Er and RI of the heavy metals in the investigated cores is given in Table 4 and Figure 5. All analysed metals showed low potential ecological risk, it varied from 1.19 to 22.96 (Er < 40). The RI of the studied trace metals in sediments of Moloundou swamp ranged from 43.24 to 47.36 It is clear that all the cores showed low ecological risk index, this indicates low polluted according to Hakanson [29].

238

Table 4. Pollution indices (Er and RI) in the sediment of Moloundou swamp

Sites -	Ecological Risk (Er)							
	Zn	Pb	Ni	Cr	Со	Cu	RI	
Core 1	1.23	4.18	4.98	4.00	9.21	21.85	45.44	
Core 2	1.33	3.15	5.39	4.15	8.77	21.41	44.20	
Core 3	1.33	5.83	4.90	4.00	8.33	22.96	47.36	
Core 4	1.19	3.13	5.25	4.00	10.44	22.41	46.41	
Core 5	1.42	1.61	5.61	4.25	8.42	21.93	43.24	
Mean (n=48)	1.33	3.87	5.16	4.08	8.65	22.25	45.35	

239



240

Fig. 5. The ecological risk factor (Er) and the ecological risk index (RI) of heavy metals in the sediments of Moloundou swamp

243

244 3.2.3.2 Pollution Load Index (PLI)

Pollution severity and its variation along the rivers was determined with the use of pollution load index. This index is a quick tool in order to compare the pollution status of different places [54]. The values of Pollution Load Index indicated that core 5 <1 therefor considered unpolluted, while as cores 1, 2, 3 and 4 considered polluted because of the value of PLI >1 (Table 3 and Fig. 4). These results attributed principally to anthropogenic sources, such as agriculture and ores extraction.

250 3.2.4 Geo-Accumulation Index (Igeo)

The geo-accumulation index is a quantitative measure of the degree of pollution in sediments [55]. Table 5 presents the geo-accumulation index for the quantification of heavy metal accumulation in the Moloundou swamp. The Igeo grades for the study area sediments varies from metal to other and between cores (Table 5 and Fig. 6). Iron, Zinc, Lead and Nickel, and remain in grade 0 (unpolluted) at all core which suggesting that the study area sediments are in background value with respect to this metal. The Igeo for cobalt, chromium and copper while reach grade 1 at all cores (from unpolluted to moderately polluted). This may be due to the agriculture and mining exploitation.

258	Table 5. The geo-accumulation index (Igeo) of heavy metals in the sediments of
259	Moloundou swamp

Sites		Geo-accumulation Index (Igeo)								
0103	Fe	Zn	Pb	Ni	Cr	Со	Cu			
Core 1	-1.15	-0.09	-0.25	-0.18	0.12	0.09	0.46			
Core 2	-1.03	-0.05	-0.38	-0.14	0.14	0.07	0.46			
Core 3	-1.00	-0.05	-0.11	-0.18	0.12	0.05	0.49			
Core 4	-1.14	-0.10	-0.38	-0.16	0.12	0.14	0.48			
Core 5	-1.13	-0.02	-0.67	-0.13	0.15	0.05	0.47			
Mean (n=48)	-1.09	-0.05	-0.29	-0.16	0.13	0.06	0.47			

260



261

Fig. 6 The geo-accumulation index (Igeo) of heavy metals in the sediments of Moloundou

263 **swamp**

264

265 4. CONCLUSION

The aims of the present study were to assess the heavy metals levels and indices in the sediments of Moloundou swamp (Southeast-Cameroon). The distribution of these metals in the sediments is almost uniform over the whole swamp and the change in concentration was due to the release of these metals from different anthropogenic and natural sources.

In the present investigation, concentrations of Cu were higher than the safe recommended values,
which suggested sediments of Moloundou swamp are polluted by Cu and might create an adverse
effect on the swamp ecosystem.

Evaluation of metal toxicity based on heavy metals indices revealed that the sediments are seriously
contaminated with Cu preferentially. The others metals (Fe, Co, Ni, Cr, Zn and Pb) are not negligible,
but not more than the threshold toxicity values. Heavy metal levels and distribution was found higher
at that cores sediments which were in vicinity of artisanal ores extraction and agricultural areas.
Government authorities must ensure strict enforcement of industrial and artisanal effluents to save the
swamp from further degradation.

279 **REFERENCES**

Fryirs, K., Freidman, B., Williams, R., Jacobsen, G. 2014. Peatlands in eastern Australia?
 Sedimentology and age structure of temperate highland peat swamps on sandstone (THPSS) in the
 Southern Highlands and Blue Mountains of NSW, Australia. *The Holocene*, *24*(11), 1527-1538.

- 283 2. Baird, C., Cann, M. 2008. Environmental Chemistry Solutions Manual (Vol. 2). Macmillan.
- 3. Kvaerner, J., Kløve, B. 2008. Generation and regulation of summer runoff in a boreal flat fen.
 Journal of Hydrology, *360*(1-4), 15-30.
- 4. Młynarczyk, N., Bartoszek, M., Polak, J., Sułkowski, W. W. 2013. Forms of phosphorus in sediments from the Goczałkowice Reservoir. *Applied geochemistry*, *37*, 87-93.
- 5. Ekoa, B. A. Z., Ngueutchoua, G., Ndjigui, P.D. 2018. Mineralogy and geochemistry of sediments
 from Simbock Lake, Yaoundé area (southern Cameroon): provenance and environmental implications. *Arabian Journal of Geosciences*, *11*(22), 710.
- 291 6. Zahran, M. A. E. K., El-Amier, Y. A., Elnaggar, A. A., Mohamed, H. A. E. A., El-Alfy, M. A. E. H.
 2015. Assessment and distribution of heavy metals pollutants in Manzala Lake, Egypt. *Journal of*293 *Geoscience and Environment Protection*, *3*(6), 107-122.
- 7. Mimba, M. E., Ohba, T., Fils, S. C. N., Nforba, M. T., Numanami, N., Bafon, T. G., Festus T. A.,
 Suh, C. E. 2018. Regional geochemical baseline concentration of potentially toxic trace metals in the
 mineralized Lom Basin, East Cameroon: a tool for contamination assessment. *Geochemical transactions*, *19*(1), 11.
- 298 8. Fu, W., Li, H., Lubow, S., Li, S., Liang, E. 2014. Effects of dust feedback on vortices in 299 protoplanetary disks. *The Astrophysical Journal Letters*, *795*(2), L39.
- 9. Chai, M., Li, R., Tam, N. F. Y., Zan, Q. 2018. Effects of mangrove plant species on accumulation of
 heavy metals in sediment in a heavily polluted mangrove swamp in Pearl River Estuary, China.
 Environmental geochemistry and health, 6: 1-15.
- 10. El-Amier, Y. A., Al-Hadithy, O. N., Kadhim, O. G., El-Alfy, M. A. E. H. 2018. Evaluation of Water and Sediment Quality of the Tigris River, Baghdad City, Iraq. *American Journal of Earth and Environmental Sciences*, *1*(1), 10-19.
- 11. Hanif, N., Eqani, S. A., Ali, S. M., Cincinelli, A., Ali, N., Katsoyiannis, I. A., Zafar I. T., Bokhari, H.
 2016. Geo-accumulation and enrichment of trace metals in sediments and their associated risks in the
 Chenab River, Pakistan. *Journal of Geochemical Exploration*, *165*, 62-70.

12. Demirak, A., Yilmaz, F., Tuna, A. L., Ozdemir, N. 2006. Heavy metals in water, sediment and
 tissues of Leuciscus cephalus from a stream in southwestern Turkey. *Chemosphere*, *63*(9), 1451 1458.

- 312 13. Zheng, N. A., Wang, Q., Liang, Z., Zheng, D. 2008. Characterization of heavy metal concentrations
 313 in the sediments of three freshwater rivers in Huludao City, Northeast. *Environmental Pollution 154*(1),
 314 135-142.
- 14. Iqbal, J., Shah, M. H. 2014. Occurrence, risk assessment, and source apportionment of heavy
 metals in surface sediments from Khanpur Lake, Pakistan. *Journal of Analytical Science and Technology*, *5*(1), 28.
- 15. Cai, W., Huang, H., Li, C., Zhang, J., Zeng, H. 2018. Effects of Humus on Heavy Metals
 Absorption of Maize in Heavy Metal Contaminated Soil. In *IOP Conference Series: Materials Science*and Engineering (Vol. 394, No. 5, p. 052061). IOP Publishing, 2018, July.
- 16. Silveira, C. S., Brandão, V. S., Bernedo, A. V. B., Mantovano, J. L. 2016. Geochemistry of river
 suspended sediments in tropical watersheds: anthropogenic and granite-gneiss sources, SE Brazil.
 International Journal of River Basin Management, *14*(4), 385-391.
- 17. El-Alfy, M., Lashin, A., Abdalla, F., Al-Bassam, A. 2017. Assessing the hydrogeochemical
 processes affecting groundwater pollution in arid areas using an integration of geochemical equilibrium
 and multivariate statistical techniques. *Environmental Pollution*, 229, 760-770.
- 18. Wang, H., Nie, L., Xu, Y., Lv, Y. 2017. The effect of highway on heavy metal accumulation in soil in turfy swamps, Northeastern China. *Water, Air, & Soil Pollution, 228*(8), 292.
- Xiao, R., Bai, J., Huang, L., Zhang, H., Cui, B., Liu, X. 2013. Distribution and pollution, toxicity and
 risk assessment of heavy metals in sediments from urban and rural rivers of the Pearl River delta in
 southern China. *Ecotoxicology 22*(10), 1564-1575.
- 20. Zahra, A., Hashmi, M. Z., Malik, R. N., Ahmed, Z. 2014. Enrichment and geo-accumulation of
 heavy metals and risk assessment of sediments of the Kurang Nallah—Feeding tributary of the Rawal
 Lake Reservoir, Pakistan. Science of the Total Environment 470, 925-933.
- 21. El-Amier, Y. A., Elnaggar, A. A., El-Alfy, M. A. 2017. Evaluation and mapping spatial distribution of
 bottom sediment heavy metal contamination in Burullus Lake, Egypt. *Egyptian Journal of Basic and Applied Sciences*, 4(1), 55-66.
- 22. Alvarez, P. 1995. Evidence for a Neoproterozoic carbonate ramp on the northern edge of the
 Central African Craton: relations with Late Proterozoic intracratonic troughs. *Geologische Rundschau*,
 84(3), 636-648.
- 341 23. Vicat, J.P., Pouclet, A., Koumbou, C., Semé Mouangué A. 1997. The fissural volcanism
 342 Neoproterozoic series of lower Dja, Yokadouma (Cameroon) and Nola (RCA) Geotectonic
 343 significance. *C.R. Acad. Sci.* Paris 325: 671-677. (*In French*).
- 344 24. Agunbiade, F. O., Olu-Owolabi, B. I., Adebowale, K. O. 2009. Phytoremediation potential of
 345 Eichornia crassipes in metal-contaminated coastal water. *Bioresource Technology*, *100*(19), 4521346 4526.
- Liaghati, T., Preda, M., Cox, M. 2003. Heavy metal distribution and controlling factors within
 coastal plain sediments, Bells Creek Catchment, southeast Queensland, Australia. *Environment International, 29*, 935–948. doi:10.1016/S0160-4120(03)00060-6.
- Turekian K., Wedepohl K. 1961. Distribution of the elements in some major units of the earth's
 crust. *American Geological Society. Bulletin, 72*, 175–182.
- 352 27. Sutherland, R. A. 2000. Bed sediment-associated trace metals in an urban stream, Oahu, Hawaii.
 353 *Environmental geology*, *39*(6), 611-627.
- 28. Tomlinson, D. L., Wilson, J. G., Harris, C. R., Jeffrey, D. W. 1980. Problems in the assessment of heavy-metal levels in estuaries and the formation of a pollution index. *Helgoländer meeresuntersuchungen*, *33*(1), 566.
- 357 29. Hakanson, L. 1980. An ecological risk index for aquatic pollution control. A sedimentological
 358 approach. *Water research*, *14*(8), 975-1001.
- 359 30. Zhu, W., Bian, B., Li, L. 2008. Heavy metal contamination of road-deposited sediments in a 360 medium size city of China. *Environmental monitoring and assessment*, *147*(1-3), 171-181.

- 361 31. Harikumar, P. S., Nasir, U. P., Rahman, M. M. 2009. Distribution of heavy metals in the core
 362 sediments of a tropical wetland system. *International Journal of Environmental Science & Technology*,
 363 6(2), 225-232.
- 364 32. Muller, G. 1969. Index of geo-accumulation in sediments of the Rhine River. *Geojournal 2*(3), 108-118.
- 366 33. Chakravarty, M., Patgiri, A. D. 2009. Metal pollution assessment in sediments of the Dikrong River,
 367 NE India. *Journal of Human Ecology*, *27*(1), 63-67.
- 368 34. Buccolieri, A., Buccolieri, G., Cardellicchio, N., Dell'Atti, A., Di Leo, A., Maci, A. 2006. Heavy
 369 metals in marine sediments of Taranto Gulf (Ionian Sea, southern Italy). *Marine chemistry*, *99*(1-4),
 370 227-235.
- 371 35. Wedepohl K. 1995. The composition of the continental crust. Geochemica and *Cosmochimica* 372 *Acta, 59*, 1217-1232.
- 373 36. Agency for Toxic Substance and Disease Registry (ATSDR). Toxicological Profile for Copper.
 374 2004. Available at: <u>http://www.atsdr.cdc.gov/toxprofiles/tp132.pdf</u>
- 375 37. Saleem, M., Iqbal, J., Shah, M. H. 2013. Study of seasonal variations and risk assessment of
 376 selected metals in sediments from Mangla Lake, Pakistan. *Journal of Geochemical* 377 *Exploration 125*, 144-152.
- 378 38. Ullah, K., Hashmi, M. Z., Malik, R. N. 2014. Heavy-Metal Levels in Feathers of Cattle Egret and
 379 Their Surrounding Environment: A Case of the Punjab Province, Pakistan. *Archives of environmental* 380 *contamination and toxicology 66*(1), 139-153.
- 381 39. Ramade, F. 1993. Encyclopedic dictionary of ecology and environmental sciences. International
 Bediscience.
- 40. Moore, J. W., Ramamoorthy, S. 1984. Heavy Metals in Natural Waters: Applied Monitoring and Impact Assessment, *Springer-Verlag; New York*, 28-246.
- 41. Song, Y., Choi, M. S., Lee, J. Y., Jang, D. J. 2014. Regional background concentrations of heavy
 metals (Cr, Co, Ni, Cu, Zn, Pb) in coastal sediments of the South Sea of Korea. *Science of the Total Environment*, 482, 80-91.
- World Health Organization (WHO). 2001. Environmental Health Criteria 221 Zinc. *Geneva, Switzerland*: World Health Organization.
- 43. Chau, Y. K., Kulikovsky-Cordeiro, O. T. R. (1995). Occurrence of nickel in the Canadian environment. *Environmental Reviews*, *3*(1), 95-120.
- 44. Biney, C. A. 1995. Environmental Baseline Studies-Limnology, Sakumo II Lagoon. Ghana Coastal
 Wetlands Management Project, Accra.
- 45. Varol, M., Şen, B. 2012. Assessment of nutrient and heavy metal contamination in surface water and sediments of the upper Tigris River, Turkey. *Catena*, *9*2, 1-10.
- 46. Agency for Toxic Substance and Disease Registry (ATSDR). Toxicological Profile for Lead. 2007.
 Available at: <u>http://www.atsdr.cdc.gov/toxprofiles/tp13.pdf</u>
- 47. Abdullah, M., Fasola, M., Muhammad, A., Malik, S.A., Boston, N., Bokhari, H., Kamran, M. A.,
 Shafqat, M. N., Alamdar, A., Khan, M., Ali, N., Eqani, S.A. 2015. Avian feathers as a non-destructive
 bio-monitoring tool of trace metals signatures: A case study from severely contaminated areas. *Chemosphere, 119*, 553-561.
- 402 48. Tariq, J., Ashraf, M., Jaffar, M., Afzal, M. 1996. Pollution status of the Indus River, Pakistan, 403 through heavy metal and macronutrient contents of fish, sediment and water. *Water research 30*(6), 404 1337-1344.
- 49. Rifaat, A. E. 2005. Major controls of metal's distribution in sediments off the Nile Delta Egypt.
 Egyptian Journal of Aquatic Research 31(2), 16-28.

- 407 50. El-Amier, Y.A., Abd El-Gawad, A.M. 2016. Assessing the Sediment Pollution using Heavy Metals
 408 Indices in the Nile River Branches in Egypt, *Journal of Environmental Science and Pollution Research*,
 409 2(3), 107-112.
- 410 51. Mahmoud, E., Abd El-Kader, N. 2015. Heavy metal immobilization in contaminated soils using 411 phosphogypsum and rice straw compost. *Land Degradation & Development*, *26*(8), 819-824.

412 52. Eqani, S. A. M. A. S., Malik, R. N., Alamdar, A., Faheem, H. 2012. Status of organochlorine 413 contaminants in the different environmental compartments of Pakistan: a review on occurrence and 414 levels. *Bulletin of environmental contamination and toxicology*, *88*(3), 303-310.

415 53. Ali, Z., Malik, R. N., Qadir, A. 2013. Heavy metals distribution and risk assessment in soils affected 416 by tannery effluents. *Chemistry and Ecology 29*(8), 676-692.

417 54. Adebowale, K. O., Agunbide, F. O., & Olu-Owolabi, B. 2009. Trace metal concentration, site 418 variations and partitioning pattern in water and bottom sediments from coastal area: A case study of 419 Ondo Coast, Nigeria. *Environmental Research Journal*, *3*(2), 46-59.

55. Singh, M., Ansari, A. A., Müller, G., Singh, I. B. 1997. Heavy metals in freshly deposited sediments
of the Gomati River (a tributary of the Ganga River): effects of human activities. *Environmental Geology*, *29*(3-4), 246-252.