

Alteration in the balance between type 1 and type 2 immune responses among lead exposed population

Abstract

Several immunotoxic effects of lead were reported. However, overt cytotoxicity of major immune cell types was not evident. It was suggested that lead might rather affect the balance between T-helper 1 and T-helper 2 immune responses cytokines. Such hypothesis was investigated in the current study and the whole blood lead (B-Pb), interleukin-4 (IL-4) and Gamma interferon (IFN- γ) levels were measured in the sera of subjects of two groups, a group environmentally exposed to lead and a control group. The results showed significantly higher levels of blood lead and serum IL-4 and a significantly lower level of IFN- γ in the subjects of exposed group when compared to the control group. Also, a statistically significant negative correlation was found between B-Pb level and serum IFN- γ in all recruited (80) subjects. It is recommended to consider lead exposure as a potential cause of immunotoxic effects even with no overt immune cell cytotoxicity. .

Key Words: Lead, environmental exposure, immune response, immunotoxic, blood lead level, interleukin-4, Gamma interferon.

1. Introduction and Background

Lead is an abundant toxic metal that is detectable in practically all phases of the inert environment and in all biological systems (**Goyer, 2001**). In Egypt, lead smelting facilities are a major source of environmental pollution by lead. In 1999, there were 27 lead smelters in Egypt in seven governorates: Cairo, Qalioubiya, Giza, Alexandria, Dakahliya, Minya and Sharkiya. Most of them are secondary smelting facilities (**Chemonics International, 1999**). Lead is known to have several toxic effects on different body systems. However, the neurological and immune systems appear to be most sensitive to lead. Recent literature showed that the immune system may even be impaired by lead levels below those required for neurotoxicity to occur. Thus, many people may suffer from suboptimal immune function due to undiagnosed lead exposure (**Lawrence and McCabe, 2002**). Many studies have shown that lead exposure can lead to immunodeficiencies (**Dietert and McCabe Jr., 2007**). Though, more recently, the spectrum of immunopathologies affected **by lead has and reports have suggested that**

lead exposure contributes also to increased risk of atopy, asthma and some forms of autoimmunity (*Dietert and Piepenbrink, 2006*).

Lead exposure at low to moderate levels, **however**, appears not to produce overt cytotoxicity of major immune cell types. Instead, immune-associated health effects result from disrupted regulation and shifts in functional capacity. Both in vivo and in vitro studies suggest that lead exposure at low to moderate levels causes a significant shift in the production of Th1 versus Th2 cytokines with the bias toward the latter. Production of IL-4, IL-6 and frequently IL-10 is elevated while production of IFN- γ is decreased (*Dietert and McCabe Jr., 2007*). However, there were other studies that reported exceptions to Pb-induced biasing in favor of Th2. These controversial findings together with the apparent gap of information as regards the effects in environmentally exposed populations were the triggers of the current study.

2. Aim of the Study

The overall objective of this study is to investigate the effect of environmental exposure to lead on the immune system, particularly the effect on the balance between type 1 (T-helper 1) and type 2 (T-helper 2) immune responses. This entails studying the effect of environmental

exposure to lead on blood lead level and investigating the balance between IFN- γ and IL-4 and its relation to lead exposure and blood lead levels in the study subjects.

3. Subjects and Methods

The study involved 80 subjects that were recruited into two groups;

a) An “Exposed Group” that consisted of 46 subjects who were environmentally exposed to lead as a result of living a residential area (Al-Molliha) that surrounds one of the oldest and largest secondary lead smelters in Cairo, Egypt. The production from the smelter, however, was stopped 15 years prior to the study conductance. The subjects were selected as a purposive sample from about 10,000 populations living within 500 meters from the smelter. **B) A “Control Group”** composed of 34 subjects recruited from a rural area (Abu Rawash) at the borders of Greater Cairo and are not exposed to lead, neither environmentally nor occupationally.

Twenty eight subjects of the exposed group were females and 18 were males. Age ranged from 11 to 70 years (with a mean of 41.30 and a standard deviation of 16.08 years). On the other hand, control group consisted of 11 females and 23 males with their age ranging from 19 to 26 years (with a mean of 38.94 and a standard deviation of 12.43 years).

The whole studied population was subjected to interview, history taking, medical examination and investigation in the form of whole blood lead level, serum interleukin-4 (IL-4) and serum Gamma interferon (IFN- γ). For performing such investigations, a 7.5 mL peripheral venous blood sample was collected by venipuncture from non-fasting subject. From this sample, 2.5 mL was collected in tubes containing EDTA as an anticoagulant and was used to determine blood lead level and 5 mL was collected in blood tubes without additives or anticoagulants and was used for analysis of IL-4 and IFN- γ .

Blood samples used for assessment of blood lead level were wet-ashed by heating in the presence of mineral acid, and subjected to analysis by atomic absorption spectrometry (AA Series Spectrometer manufactured by Thermo Electron Corporation, USA) using flame photometer.

Blood collected in tubes without additives or anticoagulants (5 mL) was allowed to clot then centrifuged to obtain serum, which was stored at - 20°C until use for cytokine levels determination. Plasma levels of IL-4 and IFN- γ were measured using an in-vitro enzyme-linked immunosorbent assay (ELISA), according to the manufacturer's instructions (Invitrogen, Belgium).

Study data were coded, entered and processed using the statistical package SPSS version 15.

4. Results and Discussion

There was a **statistically significant difference** between exposed and control groups as regards blood lead level (with higher levels in exposed group subjects), serum Interlukin-4 (IL-4) (with higher levels in exposed group subjects), serum Interferon Gamma (IFN- γ) (with lower levels in exposed group subjects) (**Table - 1 and Figure - 1**).

When studied in all subjects, there was **no statistically significant correlation** between blood lead level and serum Interlukin-4 (IL-4).

However, a **statistically significant negative correlation** was found between blood lead level (on one hand) and serum Interferon Gamma (IFN- γ) and the IFN- γ /IL-4 ratio (on the other hand) (**Table - 2 and Figure - 2**).

The results of the present work come in agreement with the findings of several studies that investigated the shift in the balance between Th1 and Th2 immune responses in relation to lead exposure. **Kaminska et al., 1998** have proven that when the blood leukocyte of cows were induced in vitro to produce IFN- γ and TNF- α in the presence of different concentrations of

lead, a significant reduction in IFN- γ titers but enhanced production of TNF- α was observed.

Hemdan et al., 2005 also studied the influence of lead on the in vitro cytokine production of human peripheral mononuclear blood cell. The results showed that expression of interferon- γ (IFN- γ), interleukin-1 β (IL-1 β), and tumor necrosis factor- α (TNF- α) was reduced whereas IL-4, IL-6, and IL-10 secretion was increased in the presence of lower lead doses. **Chen et al., 1999** have also demonstrated that lead suppressed Th1 type responses (delayed-type hypersensitivity and IFN- γ production) and enhanced Th2-type response (IL-4 and increased TNF- α production) from macrophages in mice.

In humans, **Yucesoy, et al. 1997** reported that chronic lead exposure in lead-exposed workers resulted in significant suppression of the serum IL-1 β level and IFN- γ , but did not alter IL-2 and TNF- α level.

Valentino et al., 2007 also found that low lead-exposed workers had significantly higher plasma IL-10 level, T cell cross-regulator factor produced by Th2 cells leading to reduced Th1 cytokine production, and higher plasma TNF- α level compared to non-exposed workers.

The results of the present work and the previously mentioned agreeing studies were not in agreement with two exceptional studies. One of them

conducted by **Goebel and colleagues, 2000** who reported a local bias towards Th1 response with increased IFN- γ production in the intestinal tract of specialized autoimmune diabetes prone strain of mice treated with PbCl₂ via oral administrations in drinking water.

The other study was conducted by **Mishra et al., 2003** on lead occupationally-exposed workers. Investigators also found a significant increase in IFN- γ production in phytohaemagglutinin (PHA) stimulated culture supernatant of peripheral blood mononuclear cells (PBMCs) obtained from lead exposed individuals as compared with unexposed controls with a positive correlation between blood lead level and IFN- γ production in culture.

The findings of these two studies, however, should not be considered contradictory to the previously mentioned studies. This is because, **in the study of Mishra and colleagues**, the authors themselves attributed the results to the very high lead levels (mean B-Pb level of 88.269 ± 98 $\mu\text{g/dl}$) considered in the study. Whereas, in **Goebel and colleagues' work**, the results as a whole suggest that with some genotypes or additional disease conditions, an imbalance might occur in the direction of a gut-associated Th1 environment.

In the present work, and although the correlation between B-Pb level and serum IL-4 was not statistically significant ($r=0.141$, $p=0.212$), the sample size may have played a role in that as IFN- γ /IL-4 ratio had a statistically significant negative correlation to B-Pb level ($r=-0.284$, $p=0.011$). This later correlation may serve as a better proof for the hypothesis of the present work as IFN- γ /IL-4 ratio can be taken as an individual index of Th1/Th2 balance in each subject recruited in the study.

5. Recommendation

Consider lead exposure, even in low to moderate levels, as a potential cause of immunotoxic effects even with no overt immune cell cytotoxicity.

Table-1: Blood lead level and levels of serum IL-4 and serum IFN- γ in exposed and control groups


Parameter	Exposed Group		Control Group		Statistical difference	
	Range (median)	Mean $\pm$ SD	Range (median)	Mean $\pm$ SD	Z	p
Blood Lead (Pb) ($\mu\text{g/dl}$)	2.8 - 63.5 (15.5)	20.28 $\pm$ 13.99	4.40 – 28 (11)	12.52 $\pm$ 5.61	2.75	0.006*
Interlukin-4 (IL-4) (pg/ml)	0.07 - 166.6 (26.4)	41.26 $\pm$ 38.12	11 – 105 (15.9)	20.16 $\pm$ 16.14	3.675	<0.001*

Interferon Gamma (IFN-γ) (pg/ml)	2.89 - 85.33 (6.33)	8.96 $\pm$ 12.38	11 - 28 (19.79)	19.52 $\pm$ 3.5	6.89 2	<0.001 *
*Statistically significant correlation						

Table-2: Correlation between blood lead level and levels of serum IL-4 and serum IFN- γ in all study subjects

Blood Lead (Pb) (μg/dl)		Parameter
<i>p</i>	<i>r</i>	
0.212	0.141	Interlukin-4 (IL-4) (pg/ml)
0.011*	-0.284	Interferon Gamma (IFN-γ) (pg/ml)
0.018*	-0.264	IFN-γ/IL-4
*Statistically significant correlation		

Figure-1: Blood lead level and levels of serum IL-4 and serum IFN- γ in exposed and control groups







Figure-2: Correlation between blood lead level and levels of serum IL-4 and serum IFN- γ in all study subjects



6. References

- Chemonics International. (1999). Lead Smelters and Lead Foundries in Egypt. Available on: http://pdf.usaid.gov/pdf_docs/PNACY045.pdf (August 23, 2014).
- Chen, S., Golemboski, K.A., Sanders, F.S., & Dietert, R.R. (1999). Persistent effect of in utero meso 2, 3-dimercaptosuccinic acid (DMSA) on immune function and lead induced immunotoxicity. *Toxicology*, 132, 67-79.
- Dietert R.R., & McCabe Jr. M.J. (2007). Lead immunotoxicity. In R. Luebke, R. House and I. Kimber (Ed.). *Immunotoxicology and immunopharmacology* (3rd Ed.) (pp. 207-225). Boca Raton: CRC Press.
- Dietert R.R., & Piepenbrink M.S. (2006). Lead and immune function. *Critical Review Toxicology*, 36 (4), 359-385.
- Goebel C., Flohé S.B., Kirchhoff K., Herder, C., & Kolb, H. (2000). Orally administered lead chloride induces bias of mucosal immunity. *Cytokine*, 12(9), 1414-1418.
- Goyer R.A., & Clarkson T.W. (2001). Toxic effects of metals. In C.D. Klaassen (Ed.). *Casarett and Doull's toxicology: the basic science of poisons* (6th Ed.) (pp. 827–834). New York, NY: McGraw-Hill.

- Hemdan N.Y., Emmrich F., Adham K., Wichmann G., Lehmann I., El-Massry A., ..., & Sack U. (2005). Dose-dependent modulation of the in vitro cytokine production of human immune competent cells by lead salts. *Toxicological Science*, 86 (1), 75-83.
- Kaminska T., Filar J., Madej E., Szuster-Ciesielska A., & Kandefer-Szerszen M. (1998). Modification of bovine interferon and tumor necrosis factor production by lead in vivo and in vitro *Archivum Immunologiae et Therapiae Experimentalis [Arch Immunol Ther Exp (Warsz)]*, 46, 323–328.
- Lawrence D.A., & McCabe Jr M.J. (2002). Immunomodulation by metals. *International Immunopharmacology*, 2 (2-3), 293-302.
- Mishra K.P., Singh V.K., & Rani R. (2003). Effect of lead exposure on the immune response of some occupationally exposed individuals. *Toxicology*, 188, 251- 259.
- Valentino M., Rapisarda V., & Santarelli L. (2007). Effect of lead on the levels of some immunoregulatory cytokines in occupationally exposed workers. *Human & Experimental Toxicology*, 26, 551-556.
- Yucesoy B., Turhan A., Ure M., & Karakaya A. (1997). Effects of occupational lead and cadmium exposure on some immunoregulatory cytokine levels in man. *Toxicology*, 123: 143–147.