

Modulation of Immunological and Hematological disturbances of diabetes mellitus by Diets containing combined leaves of *Vernonia amygdalina* and *Gongronema latifolium*

ABSTRACT

Background: Immunological and hematological alterations often occur in diabetes mellitus as a result of oxidative stress. The study investigated the effects of consumption of diets containing *Vernonia amygdalina* leaves on some biomarkers of immunology and hematology in Streptozotocin -induced diabetic wistar rats with the view to evaluating its involvement in the management of immunological and hematological complications among diabetics. **Method:** The design consisted of eighty rats randomly divided into eight groups of 10 rats per group. Groups 2, 3, 4, 5, 6, 7 and 8 were diabetic, while group 1 was normal control. Group 1 and 2 were fed with control diet. Group 3 to 7 were fed with diets containing *Vernonia amygdalina*, or *Gongronema latifolium* or combined leaves of *Vernonia amygdalina* and *Gongronema latifolium*. Group 8 was fed with the control diet but treated with Insulin. Feed and water were given *ad libitum* for 28 days. **Results:** Results showed that diabetic rats consuming diets of *Vernonia amygdalina*, *Gongronema latifolium* leaves or the combined leaves diets all had significant ($p < 0.5$) increase in the RBC, hemoglobin, and lymphocyte counts relative to the diabetic control. All the diets resulted in significant ($p < 0.5$) reduction in the level of WBC, platelets, neutrophil, and CD4 cell count relative to the diabetic control. The results for *Vernonia amygdalina* and *Gongronema latifolium* diets were similar to insulin on the measured parameters and their levels were not significantly different ($p < 0.5$) when compared to the normal control. Diets containing combined leaves of *Vernonia amygdalina* and *Gongronema latifolium* performed better than insulin and the measured parameters were numerically equals to the normal control. **Conclusion:** The findings of the present study indicate that consumption of diets containing *Vernonia amygdalina* or *Gongronema latifolium* leaves or in combination at the levels in this study have significant role in ameliorating hematological and immunological disturbances associated with diabetes mellitus with the combined leaf diet performing better due to positive synergy among the leaves bioactive agents and may be recommended as dietary therapies for management of immunological and hematological complications in diabetes mellitus.

Key words: Diabetes, Diet, *Vernonia amygdalina*, *Gongronema latifolium*, immunological, hematological, cardiovascular risk.

INTRODUCTION

Diabetes mellitus (DM) is a major global health problem [1],[2]. According to estimates of the World Health Organisation, the number of people suffering from diabetes worldwide is increasing at an alarming rate. There were 346 million people suffering from diabetes worldwide in 2011 [3]. It is predicted that about 366 million people are likely to be diabetic by the year 2030 [4]. Report has it that low and middle –income countries will bear the brunt of the increase and that Africa will contribute significantly to the rise [4]. Cure eludes physicians and many sufferers especially in poor countries of the world cannot meet the cost

of conventional drugs. Therefore, it has become desirable to investigate alternative sources of medicament.

Many reports have been made in recent years regarding the relationship between diabetes and dietary pattern [5]. Many of these studies have reported that vegetable intake reduces the onset of diabetes and improves plasma glucose control, blood lipids and renal function in diabetic patients thus reducing the risk of cardiovascular events [5]. There are few studies on vegetarian diets in diabetes at present, considering the enormous number of vegetables and plant products in the world. Therefore, novel plants and plant components that may have significant outcome and provide an alternative to drug therapy are desirable. There is scarcity of literature with regards to the implication of combining several plants or plants components in diet for diabetes management. This study assessed the effect of consumption of diets containing combined leaves of *Vernonia amygdalina* and *Gongronema latifolium* on hematological and immunological parameters of Streptozotocin induced experimental diabetic Wistar rats, with the view to evaluating their potential role in the management of hematological and immunological complications common among diabetics.

The underlying cause of the disturbances in the immunity and hematology in diabetes mellitus is mainly due to oxidative damage [6]. Although almost all organisms possess antioxidant defense and repair systems that have evolved to protect them, these systems are insufficient to protect completely in many disease conditions. For this reason, green leafy vegetables have been widely investigated for their protective action in diabetes mellitus since they contain valuable antioxidant, especially antioxidant vitamins including ascorbic acid, α -tocopherol, β -carotene and phenolics [7], [8].

Vernonia amygdalina Del popularly called bitter leaf is a species of *Vernonia* that belongs to the family Compositae (Asteraceae). *Gongronema latifolium* Benth is of the family (Asclepiadaceae). It is a tropical rainforest plant, a climber with tuberous base limited in distribution to wet and dry forest of tropical Africa [9], [10], Guinea Bissau and western Cameroun [11]. These plants are popularly used in the preparation of soups in many African countries.

MATERIALS AND METHODS

Collection and processing of plant materials

Fresh but matured leaves of *Vernonia amygdalina* Del and *Gongronema latifolium* were collected from the Endocrine Research Farm, University of Calabar, and from University of Calabar Staff Village, Calabar in March, 2011. These leaves were authenticated by a Taxonomist and Voucher Specimens were deposited in the herbarium in the Department of Botany, University of Calabar. The leaves were selected to remove extraneous materials, washed and rinsed with distilled water and dried under shade until the leaves were well dried. Dried leaves of *Vernonia amygdalina* Del and *Gongronema latifolium* were milled separately using commercial feed mill machine (Artec model 40) to powder and sieved with 1mm mesh to obtain fine leaf powder. Fine leaf powder were packaged separately in a well-labeled amber container and stored in the refrigerator at temperature 2- 8°C until used for the preparation of rat chow.

Formulation of experimental diets

Feed ingredients include: leaf powder, soybean meal, maize meal, Garri, mineral premix, vitamin premix, L-lysine L-methionine and corn oil. These feed ingredients were purchased

from Victory Livestock Ltd, an accredited Livestock feeds/ vaccines/drug dealers, located at 79, Aka road, Uyo, Akwa Ibom State. Standard rat chows (growers) were formulated according to the nutritional requirement of rat [12]. (Table 1). Seven (7) different diets were formulated namely: Control, VA-5%; GL-5%, VA+GL-5%, VA-7.5%, GL-7.55, VA+GL-7.5%. Control diet differed from the other five diets because it did not contain leaf powder, but had all the other feed ingredients contained in the other diets. The other five diets contained leaf powder at five (5%) or seven and a half (7.5%) percent. The percentage composition and nutrient analysis of the experimental diets are shown table 1.

Animals

Eighty (80) albino rats of Wistar strain (female only) weighing between 83-121g were purchased from the animal house of the Faculty of Basic Medical Science, University of Uyo, Uyo, and transported in well ventilated cages to the animal house of the Department of Biochemistry, University of Calabar, Cross River State, where they were kept throughout the duration of the experiment. The animals were allowed to acclimatize for two weeks. They were housed in well ventilated cages (wooden bottom and wire mesh top) and kept under controlled environmental conditions of temperature ($25 \pm 5^{\circ}\text{C}$), relative humidity ($50 \pm 5\%$) and twelve hour light/dark cycle. Approval was granted by the Ethic committee of the College of Basic Medical Science, University of Calabar and the animals were kept under the care of a trained animal technician and cared for according to [13]. Animals were allowed free access to water and chow over a two weeks adaptation period and closely monitored.

Experimental design and induction of experimental diabetes mellitus

The design consisted of eighty (80) female rats divided into 7 groups of diabetic and 1 group of normal rats with 10 animals in each group. The rats in the diabetic groups were subjected to an overnight fast (12hrs) prior to induction of diabetes. The weight of individual rats were measured and noted. Diabetes mellitus was induced in the diabetic groups by intraperitoneal injection of 55mg/kg body weight of Streptozotocin, (STZ) (sigma St. Louis, MO. USA) reconstituted in 0.1%M sodium citrate buffer. The pH of the buffer was adjusted to 4.5. Rats whose fasting blood glucose concentration were higher or equal to 200 mg/dl three days after the induction were confirmed diabetic and recruited in the study. Blood glucose concentration was determined using one touch Glucometer (Lifescan, Inc. 1995, Milpas, Galifornia, U.S.A) with blood obtained from the tail vein of the rats. The seven diabetic groups were randomly assigned to the seven experimental diets: control, VA-5%, GL-5%, VA+GL-5%, VA-7.5%, GL-7.5% and VA+GL-7.5%. Insulin, a standard chemotherapeutic agent was introduced for comparison. Insulin dose used was 5U/kg body weight (b.w), given subcutaneously (s.c) according to [14]. It was given once per day post prandial. That is, the group that received insulin was fed with the control diet. Rats in the normal group were fed with the control diet. Feed and water (Tap water) were given *ad-libitum*. Treatment lasted for 28 days. Daily recording of feed intake, weight of animals and blood glucose concentration were kept.

Collection of sample for analysis

At the end of the 28 days, food and water were withdrawn. They rats fasted overnight. They were then euthanized under chloroform vapor and sacrificed. Whole blood was collected via cardiac puncture using sterile syringes and needles. The blood was emptied into EDTA sample bottles. The samples were used for analysis within 12h of collection.

Determination of full blood count (FBC) using automated hematology analyzer, KX2IN (non-cyanide hemoglobin analysis method)

Full blood counts including PCV (HCT), HB, RBC, WBC, platelet count, differential WBC (lymphocytes and mixed), red cell indices MCHC, MCH and MCV), were estimated using the Sysmex® Automated Analyzer KX-2IN, Sysmex Corporation, Kobe-Japan.

CD₄⁺ count

The CD₄⁺ lymphocyte was estimated by flow cytometry [15] using the cyflow automated cell counter (Parlec, Germany). Ten microlitres of CD₄⁺ PE antibody was mixed with 5ml of EDTA anticoagulated whole blood in a test tube. The mixture was incubated in the dark chamber for 15 min at room temperature of 22 -28⁰C. During incubation, the content of the tube was mixed every five min, eight hundred microlitres of buffer was added, mixed and plugged into the counter. After, counting the CD₄⁺ cells, monocytes and noise were separated gated and the result was recorded.

Statistical analysis

The results were analyzed for statistical significance by one-way ANOVA using the SPSS statistical program and least square test (LSD) between group using MS excel programme. Student T tests were used to assess differences between individual variables. All data were expressed as mean ± SEM. P value <0.05 was considered significant.

RESULTS AND DISCUSSION

The effects of consumption of diet containing combined leaves of *Vernonia amygdalina* and *Gongronema latifolium* on some hematological and immunological parameters of diabetic rats are shown in Table 2 and 3 respectfully. The results in Table 2 showed that the diabetic untreated rats (diabetic control) had significantly higher (P<0.05) level of WBC (10.60 ±2.00 x 10³/μl) relative to the normal control (6.87 ±2.28 x 10³/μl). The diabetic rats treated with diets and insulin had white blood cells count that were significantly lower (P<0.05) compared to the diabetic control. The effect of *Vernonia amygdalina* diet on WBC appeared to be dose dependent, better at the higher inclusion level (7.5%). *Gongronema latifolium* diet also had dose dependent effect on WBC, but better at lower level of inclusion (5%). Combination of *Vernonia amygdalina* and *Gongronema latifolium* in the diet exerted an effect that brought the WBC to a level numerically equal to those of normal control and this treatment was better than insulin particularly at 5% inclusion level. RBC, HGB and HCT as well as red blood cell function indices: MCV, MCH and MCHC of the diabetic control were all significantly lower (p<0.05) compared to the normal control, but the Platelet of the diabetic control was significantly higher (p<0.05) compared to the normal control (Table 2). Treatment with diets and insulin significantly increased (P<0.05) the RBC, HGB and HCT as well as red blood cell function indices: MCV, MCH and MCHC of the diabetic treated rats (Table 2). The increased in the RBC, HGB and HCT as well as red blood cell function indices: MCV, MCH and MCHC following consumption of the diets were not dose dependent. However, the RBC, HGB and HCT tended to increase more at higher inclusion level of *Vernonia amygdalina* (7.5%), but at lower inclusion level of *Gongronema latifolium* (5%). The MCV, MCH and MCHC tended to increase more at lower level of inclusion of *Vernonia amygdalina* but no pattern was followed by *Gongronema latifolium*.

Table 3 showed that the diabetic control had neutrophil, basophil, monocyte and CD₄⁺ cell count that were significantly higher (P<0.05) compared to the normal control (30.33±4.95%,

1.81±0.00%, 5.40±0.00%, and 51.53±0.77 x 10⁶/l respectively) compared to the normal control (23.00±1.93%, 0.00±0.00%, 4.67±0.33% and 15.70±0.36 x 10⁶/l respectively). The esenophil and lymphocyte cell counts for the diabetic control rats (1.33±0.55% and 43.33±4.21 respectively) were significantly lower (p<0.05) compared to the normal control (2.67±0.56 and 70.67±2.19 respectively). Treatment with the diets and insulin significantly lower (P<0.05) the neutrophil, monocyte and the CD₄⁺ cell count but increased the esenophil, and lymphocyte cell count relative to the diabetic control. The *Vernonia amygdalina* diet had dose dependent effect on the neutrophil and better at low inclusion level (5%). The effect of *Gongronema latifolium* diet on the neutrophil was not dose dependent but was better at the high inclusion level (7.5%). Combination of *Vernonia amygdalina* and *Gongronema latifolium* in the diet exerted better effect on the neutrophil than using *Vernonia amygdalina* or *Gongronema latifolium* alone. This brought the neutrophil to a level that was numerically equal to those of the normal control; and this treatment was better than insulin particularly at 5% inclusion level. The *Vernonia amygdalina* diet had no dose dependent effect on esenophil and lymphocyte. Effect on esenophil was similar at the two levels of inclusion but on lymphocyte was better at the lower level of inclusion. The *Gongronema latifolium* diet also had no dose dependent effect on esenophil, lymphocyte, monocyte and CD₄⁺ cell count. Effect on esenophil was similar at the two levels of inclusion, on lymphocyte was better at higher dose, on monocyte it was similar at the two levels and also similar at the two levels for CD₄⁺ cell count. Combination of *Vernonia amygdalina* and *Gongronema latifolium* in the diet exerted better effect on the esenophil, lymphocyte, monocyte and CD₄⁺ cell count than using either *Vernonia amygdalina* or *Gongronema latifolium* alone. This brought their levels to be numerically equal to those of the normal control; and this treatment was also better than insulin particularly at 5% inclusion level.

Type 1 diabetes is characterized by overt destruction of pancreatic islet cells. Individual sufferers depend on insulin. Many patients with type 1 diabetes fail to meet recommended glycemic goals regardless of the recognition of optimal glycemic control as a key component for improving clinical outcomes and quality of life [16]. Patients and physician related barriers to the adoption of insulin therapy include fear and anxiety about infecting insulin, concerns about side effects, the availability and high cost of insulin and personal health beliefs in regard to the use of insulin [16]. There is an unmet need for an alternative therapy that provides optimal glycemic control, well tolerated, and improves patient adherence. There is upsurge of interest in leafy vegetables in view of the abundance of natural bioactive agents. In this study, we developed a model of Type 1 diabetes in rats and investigated the effect of consumption of diets containing combined leaves of *Vernonia amygdalina* and *Gongronema latifolium* on hematological and immunological parameters and compared to insulin which is the standard therapy. The diabetic control rats were observed to have alterations in hemoglobin (Hb), red blood cell (RBC) and white blood cell (WBC) count. The significant (p<0.05) reduction in the hemoglobin, and RBC cell and an increase in the white blood cells count, neutrophil, lymphocyte, platelet and CD₄⁺ cells of the diabetic untreated rats (diabetic control) compared to the normal control and the diabetic treated rats may be a manifestation of the diabetic condition. As it pertains to hematology of the diabetic, terms such as anaemia, atherosclerosis resulting from increased platelet aggregation, glycosylation of hemoglobin (RBC) and of recent, even white blood cells have been discussed extensively [17], [18]. The normal physiologic response following the perception of an insult by the body is an increase in WBC count. It is likely that the damage caused by diabetes contributed to the observed increase in WBC count. This is in agreement with [19] who reported that leucocytosis may occur in hepatic damage. White blood cell count is increased in obesity [20] and is a risk factor for atherosclerosis [21]. An elevated WBC count is present in impaired

glucose tolerance (IGF) [21] and is associated with macro- and micro angiopathic complications in type 2 diabetes [22]. High WBC count has been shown to be associated with insulin resistance, decrease in insulin action [23], and is predictive of the development of type 2 diabetes [24], [21], coronary artery disease [25], stroke [25], and diabetes micro- and macrovascular complications [22].

Red blood cell count, hemoglobin concentration, packed cell volume alongside with erythrocyte function indices: MCV, MCH and MCHC all reduced significantly ($P<0.05$) in the diabetic control compared to the normal group and treatment groups. Diabetes mellitus is associated with oxidative stress. The reduction in red blood cell count may be due to an increase in the production of lipid peroxides that lead to hemolysis of RBC [26]. The major pathological consequences of free radical induced membrane lipid peroxidation include increased membrane rigidity, decreased cellular deformability, reduced erythrocyte survival, and lipid fluidity [27]. It may also be due to reduction in the synthesis of erythropoietin, which stimulates stem cells in the bone marrow to produce red blood cells. Erythropoietin is produced by the kidney. Kidney damage at several levels is a complication of diabetes. Changes in the kidneys that occur with diabetes range from diabetic nephropathy all the way to chronic kidney disease (CKD).

Low hemoglobin concentration in diabetic control group may be associated with kidney damage due to diabetes. Low hemoglobin concentration was associated with low postprandial c-peptide concentration and low β -cell responsiveness [28]. Some studies suggest that β -cell dysfunction occurs due to hypoxic damage to the pancreatic islet cells [29], [28], which progressed to diabetes and low hemoglobin concentrations contributing to development of cardiovascular events in patients with diabetes [30]. Diabetes mellitus is associated with a more rapid decline in glomerular filtration rate than that of other kidney diseases [31] and diabetic nephropathy and diabetic retinopathy result in increased susceptibility to low hemoglobin level [32]. Low hemoglobin level is associated with increased cardiovascular mortality and chronic kidney disease (CKD) in patients with diabetes mellitus [33]. The decrease in MCV, MCH and MCHC values, observed in the diabetic control group is an indication of abnormal hemoglobin synthesis, failure of blood osmoregulation, and plasma osmolality [34] that may be associated with diabetes mellitus.

The level of platelet was significantly higher ($P<0.05$) in the diabetic control compared to the normal group and treatment groups. Platelets are small discoid blood cells that circulate and participate in hemostasis. Primary plug formation due to platelets seals the vascular defects and provides the required phospholipid surface for the recruited and activated coagulation factors [35]. In response to stimuli generated by the endothelium of blood vessels, platelets change shape, adhere to subendothelial surfaces, secrete the contents of intracellular organelles, and aggregate to form a thrombus [35]. These pro-aggregatory stimuli include thrombin, collagen, epinephrine, ADP (dense storage granules), and thromboxane A₂ (activated platelets) [35]. The increase in the level of platelet in the diabetic control group may be due to inflammatory responses caused by diabetes and correlate with the level and sustenance of high blood glucose level [36], [30]. Platelets assume an important role in signaling of the development of advanced atherosclerosis in diabetes [35], [36], [30].

We observed that consumption of all the diets and administration of insulin significantly reduced ($P<0.05$) WBC and platelet but significantly increased ($P<0.05$) RBC, HGB, PCV, MCV, MCH and MCHC compared to diabetic control. For red blood cells, this gives an indication that the diets could stimulate the formation or secretion of erythropoietin, which stimulates stem cells in the bone marrow to produce red blood cells [37]. This hormone

stimulation enhances rapid synthesis of RBC which is supported by the improved level of MCH and MCHC [38]. This may also suggest the restoration of oxygen carrying capacity of the blood. Although the mechanism of this effect is not well known, it is assumed that the antioxidants in the leaves [39] might have help to reduce oxidative stress of diabetes and thus improve the metabolism and well being of the red cells. The leaves rich content of blood formatting minerals and vitamins [39] might have also help to supplement those lost due to urinary excretion and this might have promoted the formation of RBC and hemoglobin. Red blood cell counts can be a factor in erythropoietin process [40]. The increased circulating erythropoietin concentrations [40] might have been able to elicit and enhance the production and expression of red cells antioxidant [41] and enhance their ability to lower lipid peroxidation level [42] that causes hemolysis of erythrocytes.

Antioxidants prevent the expression of hepcidin, a 25 amino acid peptide hormone, in the liver which is increased dramatically by inflammation and because of chronic disease [43]. Released hepcidin is bound to iron efflux protein ferroportin [44] and act as a negative regulator of body iron homeostasis, inhibiting the release of iron recycled from senescent red blood cells by reticuloendothelial macrophages [45] and the absorption of dietary iron by intestinal enterocytes [46]. Thus, the diets may participate directly in amelioration of the effect of diabetes on RBC indices and correction of the anemic status in the diabetic rats by attenuating proinflammatory cytokines production [47] which is central to the expression of hepcidin. The result we obtain is in line with the findings of [48] who reported the reversal of anaemia in cadmium toxicity after the supplementation of diet, and [6] who also reported reversal of anaemia in diabetic rats treated with some leafy vegetables.

Some parameters such as WBC and neutrophil responded to the leaves in the diet. The mechanism is not known but may relate to the level of the bioactive agents found. Failure of the leaf diet to influence the parameter at low dose level might be due to inadequate amount of the phytochemicals / nutrients needed to reverse the changes in those parameters. Combining the two leaves resulted in better performance indicating a positive synergy. The two leaves contain varying levels of flavonoids, tannins, saponins, polyphenols, alkaloids, hydrocyanic acid, vitamin A, E, C, riboflavin, thiamin, niacin, iron, selenium, and chromium. Combination of the two leaves in diet tended to synergies each other and the result obtained was better with combined leaves than using either of the leaves. Combination of the leaves in a diet at 5% impacted better than insulin on all parameters measured.

Table 3 shows that the neutrophil, basophil, monocyte and CD_4^+ cell count were significantly increased ($P<0.05$) in the diabetic control group compared to the treatment group and the normal control, while the lymphocyte was significantly reduced ($P<0.05$) in the diabetic control group compared to the treatment group and the normal control. Polymorphonuclear cells, including monocytes as well as lymphocytes make up the peripheral blood leukocytes. In a state of hyperglycemia, polymorpho- and mononuclear leukocytes can be activated by advanced glycation end products [49], oxidative stress [50], angiotensin II [51], and cytokines [52]. The release of cytokines, such as $TNF-\alpha$ [53], transforming growth factor-1 [54], superoxide [55], nuclear factor κB (NF- κB) [56], monocyte chemoattractant protein 1, interleukin- 1β , and others [53], may activate leukocytes to participate in the pathogenesis of diabetic micro- and macrovascular complications. Studies showed that diabetes in mice was accompanied by moderate neutrophilic leukocytosis and prolonged circulation times of neutrophils and monocytes, and a shortened circulation time of lymphocytes, which increases the susceptibility to infection [57]. The raised leukocyte count reflected low-grade inflammation [57]. The mechanism responsible for leukocytosis in diabetes is largely unknown but recent evidence shed light on how elevated WBC increase cardiovascular and



stroke risk linking inflammation mediated by neutrophil to insulin resistance [58]. We suggest that the raised leukocyte count may be inflammatory response to diabetes. The decrease in the WBC count, in diabetic treated rats shows the anti-inflammatory property of the *Vernonia amygdalina*, *Congronema latifolium* and the combined leaf diets. Lymphocytes and CD₄⁺ cells are the main effectors cells of the immune system. The increase in the lymphocyte count in rat placed on *Vernonia amygdalina*, *Congronema latifolium* and combined leaf diets may be an indication of immunostimulation. The reduction in CD₄⁺ cell counts might indicate reduce stimulation of immune cells recruitment in inflammatory reactions. Similar anti-inflammatory activities have been reported for *Azardiracta indica* (neem) leaf extract [59], [60]. The extract effect was attributed to inhibition of immune cells migration and phagocytosis, particularly for macrophage and neutrophils in respect to inflammatory stimuli. The extract was also seen to inhibit the induction of inducible nitric oxide synthase, prostaglandins E₂ (PGE₂) and interleukin 1 (IL-I) productions [59], thus controlling the increased vascular permeability associated with inflammatory reactions.

CONCLUSION

The findings of the present study indicate that consumption of diets containing *Vernonia amygdalina* or *Gongronema latifolium* leaves or combination have significant role in ameliorating hematological and immunological disturbances associated with diabetes with the combined leaf diet performing better due to positive synergy among the leaves bioactive agents. This could be due to the prevention or inhibition of lipid peroxidative system by their antioxidant, maintenance of cellular integrity, and attenuation of pro-inflammatory cytokine production. Consumption of diet containing combined leaves of *Vernonia amygdalina* and *Congronema latifolium* at 5% was better than using insulin and may be recommended as a dietary therapy for management of immunological and hematological complications in diabetes.

REFERENCES

1. Mahsud MAJ, Khan A, Hussain J: Hematological Changes in Tobacco using Type 2 Diabetic Patients. *Gomal J Med Sci* 2010; **8**:8-11.
2. King H, Aubert RE, Herman WH: Global burden of diabetes, 1995-2025: prevalence, numerical estimates, and projections. *Diabetes Care* 1998, **21**:1414- 31.
3. World Health Organization August 2011. Available from: <http://www.who.int/mediacentre/factsheets/fs312/en/> [Last accessed on 2011 Nov 17].
4. World Health Organisation (WHO) (2008). Diabetes in the WHO country and regional data, 2008.(website:ww.who.int/diabetes/facts/wporldfigures/en/index.html).
5. Keiko Takahashi, Chiemi Kamada, Hidenori Yoshimura, Ryota Okumura, Satosh Iimuro, Yasuo Ohashi, Atsushi Araki, Hiroyuki Umegaki, Takashi Sakurai, Yukio Yoshimura,Hideki Ito: Effects of total and green vegetable intakes on glycated hemoglobin A1c and triglycerides in elderly patients with type 2 diabetes mellitus: The Japanese Elderly Intervention Trial. *Geriatrics & Gerontology International* 2012, (**12**)50–58l.
6. Saliu JA, Elekofehinti OO, Komolafe K, Obboh G: Effect of some green leafy vegetables on the hematological parameters of diabetes rat. *J. Nat. Pro. Plant. Resour* 2012, **2** (4): 482-485.
7. Chu YF, Sun X, Wu X, Liu R. H. Antioxidant and antiproliferative activities of vegetable. *J. Agric Food Chem* 2002, **50**, 6910- 6916.

- 364 8. Oboh G, JBT: Rocha, leading edge. *Antioxid Research* 2007, 35-64
- 365 9. Ugochukwu NH , Babady N E : Antihyperglycemic effect of aqueous and ethanolic extracts of
366 *Gongronema latifolium* leaves on glucose and glycogen metabolism in livers of normal and
367 streptozotocin-induced diabetic rats. *Life science* 2002, **73**: 1925-1938.
- 368 10. Ugochukwu NH, Babady NE, Cobourne M, Gasset SR: The effect of *Gongronema latifolium*
369 extracts on serum lipid profile and oxidative stress indices in hepatocytes of diabetic rats. *Journal of*
370 *Biosciences* 2003, **28**: 1-5
- 371 11. Francis VU, Nelson CO, Simon JU: Influence of *Gongronema Latifolium* Leaf-Extracts Treatment
372 on Some Hepatic Enzymes Activity In Rats. *Journal of Natural Sciences Research* 2013, **3** (5)32-35
- 373 12. National Research Council: *Nutrition requirement of laboratory animals*, USA. 1995. 38
- 374 13. Canadian Council on Animal Care. *Guide to the care and use of experimental animals*. Vol. 2.
375 Ottawa, Ontario, Canada: CCAC, 1993
- 376 14. Sonia B, Scrivinasan BP: Investigations into the antidiabetic activity of *Azardiracta indica*. *Indian*
377 *Journal of Pharmacology* 1999, **31**: 138- 141
- 378 15. Center for Disease Control and Prevention: Revised guidelines for performing CD⁺₄ T-cell
379 determinations in persons infected with human immunodeficiency virus (HIV). *MMWR Recomm. Rep.*
380 , 1997 **46**(RR-21): 1-29.
- 381 16. Priscilla A. Hollander: Evolution of a Pulmonary Insulin Delivery System (*Exubera*) for Patients
382 With Diabetes *MedGenMed* 2007, **9**(1):45-51
- 383 17. Thomas M, Tsalamandris C, MacIsaac R, Jerums G: Anemia in diabetes: An emerging
384 complication of microvascular disease. *Current Diabetes Reviews* 2005, **1**:107-126
- 385 18. Colwell JA, Nesto RW: The platelets in diabetes: Focus on prevention of ischemic events.
386 *Diabetes Care* 2003, **26**: 2181-2188
- 387 19. Finlayson NDC, Hayes PC, Simpson KJ: Diseases of the liver and biliary system. in: Davidson's
388 principles and practice of medicine. Eds. Haslet C, Chilvers E. R, Hunker J. A. A. Boon N. A., 18th ed.
389 1999 Churchill Living Sone, Edinburgh. Pp 683-736.
- 390 20. Kullo IJ, Hensrud DD, Allison TG: Comparison of numbers of circulating blood monocyte in men
391 grouped by body mass index (< 25, 25 to <30, > or = 30. *Am. J. Cardiol* 2002, **89**:1441-1443
- 392 21. Ohshita K, Yamane K, Hanafusa M, Mori H, Mito K, Okubo M, Itara H, Kohno N. (2004).
393 Elevated white blood cell count in subjects with impaired glucose tolerance. *Diabetes Care*; **27**: 491-
394 496
- 395 22. Tong PC, Lee KF, So WY, Ng MH, Chan WB, Lo MK, Chan NN, Chan JC: (2004). **White**
396 blood cell count is associated with macro and microvascular complications in Chinese patients with
397 type 2 diabetes. *Diabetes Care*, **27**:216-22
- 398 23. Vojarova B, Weyer C, Lindsay RS, Pratley RE, Bogardus C, Tataranni PA: High white blood cell
399 count is associated with a worsening of insulin sensitivity and predicts the development of type 2
400 diabetes. *Diabetes* 2002, **51**(2):455-61.
- 401 24. Barbara GB, Małgorzata S , Joanna G, Beata K, Justyna S, Mieczysław W, Anelli S: Changes in
402 inflammatory biomarkers after successful lifestyle intervention in obese children.
403 *Endokrynologia Polska/Polish Journal of Endocrinology* 2011, **62**:499- 505.

410

- 411 25. Lee CD, Folsom AR, Nieto FJ, Chambless LE, Shahar E, Wolfe DA: White blood cell count and
412 incidence of coronary heart disease and ischemic stroke and mortality from cardiovascular
413 disease in African- American and white men and women: atherosclerosis risk in communities
414 study. *Am J Epidemiol* 2001, **154**:758-64.
- 415 26. Arun GS, and Ramesh KG: Improvement of insulin sensitivity by perindopril in spontaneously
416 hypertensive and streptozotocin diabetic rats. *Indian J Pharmacol* 2002,**34**:156- 64.
- 417 27. Kolanjiappan K, Manoharan S, Kayalvizhi M: Measurement of erythrocyte lipids, lipid peroxidation,
418 antioxidants and osmotic fragility in cervical cancer patients. *Clin Chim Acta* 2002, **326**:143-9.
- 419 28. Rhodes CJ: Type 2 diabetes-a matter of beta-cell life and death? *Science* 2005, **307**:380–384
- 420 29. Jansson L: The regulation of pancreatic islet blood flow. *Diabetes Metab Rev* 1994, **10**:407–416.
- 421 30. Stevens PE: Anaemia, diabetes and chronic kidney disease: where are we now? *J Ren Care* 2012,
422 **38**(Suppl 1):67–77
- 423 31. Rossing K, Christensen PK, Hovind P, Tarnow L, Rossing P, Parving HH: Progression of nephropathy
424 in type 2 diabetic patients. *Kidney Int.* 2004, **66**:1596–1605.
- 425 32. Ranil PK, Raman R, Rachepalli SR: Anemia and diabetic retinopathy in type 2 diabetes mellitus. *J*
426 *Assoc Physicians India*, 2010, **58**:91–94.
- 427 33. New JP, Aung T, Baker PGI: The high prevalence of unrecognized anaemia in patients with
428 diabetes and chronic kidney disease: a population-based study. *Diabet Med*, 2008 **25**:564–569.
- 429 34. Stookey JD, Burg M, Sellmeyer DE, Greenleaf JE, Arief A, Van Hove L : A proposed method for
430 assessing plasma hypertonicity in vivo. *Eur J Clin Nutr* 2007, **61**: 143-6
- 431 35. Al-Khoury S, Afzali B, Shah N, Covic A, Thomas S, Goldsmith DJ: Anaemia in diabetic patients with
432 chronic kidney disease: prevalence and predictors. *Diabetologia* 2006, **49**:1183– 1189.
- 433 36. Herzog CA, Mangrum JM, Passman R: Sudden cardiac death and dialysis patients. *Semin Dial*
434 **21**:300–307.
- 435 37. Ohlsson A, Aher SM: *Cochrane Database Syst Rev* 2012, **9**: CD004863. Early erythropoietin for
436 preventing red blood cell transfusion in preterm and/or low birth weight infants.
- 437 38. Abu-Zaiton AS: Antidiabetic activity of *Ferula asafoetida* extract in normal and alloxan induced
438 diabetic rats. *Pak J Biol Sci* . 2010, **13**:97-100.
- 439 39. Atangwho IJ, Ebong PE, Eyong EU, Williams IO, Eteng MU, Egbung GC: Comparative chemical
440 composition of leaves of some antidiabetic medicinal plants: *Azadirachta indica*, *Vernonia*
441 *amygdalina* and *Gongronema latifolium*. *African Journal of Biotechnology* 2009, **8**(18):
442 4685-4689
- 443 40. Inomata H, Itoh M, Imai H, Sato T: Serum erythropoietin levels as a novel marker of
444 severity of, diabetic nephropathy. *Nephrolog* 1997, **7**:426-9
- 445 41. Digicayliogu M, Lipton MA: Erythropoietin- mediated neuroprotection involves cross-talk
446 between Jak 2 and NF-kappa signalling cascades. *Nature* 2001, **412**: 641-647
- 447 42. Mahmoud AM: Influence of rutin on biochemical alterations in hyperammonemia in rats.
448 *ExpToxicol Pathol*; 2012, **64**:783-9.
- 449 450
- 451

- 452 43. Nicolas G, Chauvet C, Viatte L, Danan JL, Bigard X, Devaux I: The gene encoding the iron
453 regulatory peptide hepcidin is regulated by anemia hypoxia and inflammation. *J Clin Invest*
454 2002,**110**:1037-44.
- 455 44. Nemeth E, Tuttle MS, Powelson J, Vaughn MB, Donovan A, Ward DM: Hepcidin regulates cellular
456 iron efflux by binding to ferroportin and inducing its internalization. *Science* 2004, 306:2090-3.
- 457 45. Knutson MD, Oukka M, Koss LM, Aydemir F, Wessling-Resnick M: Iron release from macrophages
458 after erythrophagocytosis is up-regulated by ferroportin 1 overexpression and down-regulated by
459 hepcidin. *Proc Natl Acad Sci USA* 2005,**102**:1324-8.
- 460 46. Laftah AH, Ramesh B, Simpson RJ, Solanky N, Bahram S, Schumann: Effect of hepcidin on
461 intestinal iron absorption in mice. *Blood* 2004, **103**:394-4.
- 462 47. Mahmoud AM, Ahmed OM, Ashour MB, Abdel-Moneim A: Hesperidin and naringin attenuate
463 hyperglycemia-mediated oxidative stress and proinflammatory cytokine production in high fat
464 fed/strptozotocininduced type 2 diabetic rats. *J Diabetes Complications* 2012, **26**:483-90.
- 465 48. Eteng MU, Onwuka FC, Akpanyung EO, Oguchukwu NC, Bassey SC, Nwankpa P: Reversal of
466 cadmium induced toxicity following dietary supplementation with garlic, ginger and cabbage in male
467 wistar rats. *J. Nat.Prod. Plant Resour.* 2012, **2** (1): 169- 174.
- 468 49. Pertynska-Marczewska M, Kiriakidis S, Wait R, Beech J, Feldmann M, Paleolog EM: Advanced
469 glycation end products upregulate angiogenic and pro-inflammatory cytokine production in
470 human monocyte/ macrophages. *Cytokine* 2004, **28**:35- 47.
- 471
472 50. Shurtz-Swirski R, Sela S, Herskovits AT, Shasha SM, Shapiro G, Nasser: Involvement of peripheral
473 polymorphonuclear leukocytes in oxidative stress and inflammation in type 2 diabetic patients.
474 *Diabetes Care* 2004, **24**:104-10.
- 475 51. Lee P, Peng H, Gelbart T, Beutler E: The IL-6- and lipopolysaccharideinduced transcription of
476 hepcidin in HFE-, transferrin receptor 2-, and beta 2-microglobulindeficient hepatocytes. *Proc*
477 *Natl Acad Sci* 2004 **101**:9263-5.
- 478 52. Scherberich JE: Proinflammatory blood monocytes:main effector and target cells in systemic and
479 renal disease;background and therapeutic implications. *Int J Clin Pharmacol Ther* 2003,**41**:459- 64.
- 480 53. Shanmugam N, Reddy MA, Guha M, Natarajan R: High glucose-induced expression of
481 proinflammatory cytokine and chemokine genes in monocytic cells. *Diabetes* 2003, **52**:1256 64.
- 482 54. Korpinen E, Groop PH, Fagerudd JA, Teppo AM, Akerblom HK, Vaarala O: Increased secretion of
483 TGF-beta1 by peripheral blood mononuclear cells from patients with type 1 diabetes mellitus
484 with diabetic nephropathy. *Diabet Med* 2001,**18**:121-5.
- 485 55. Kedziora-Kornatowska KZ: Production of superoxide and nitric oxide by granulocytes in non-
486 insulin-dependent diabetic patients with and without diabetic nephropathy. *IUBMB Life* : 1999,
487 **48**:359-62..
- 488 56. Hofmann MA, Schiekofer S, Kanitz M, Klevesath MS, Joswig M, Lee V: Insufficient glycemic
489 control increases nuclear factor-kappa B binding activity in peripheral blood mononuclear cells
490 isolated from patients with type 1 diabetes. *Diabetes Care* 1998, **21**:1310-6.
- 491 57. Kozlov IA, Novitski VV, Baikov AN: Kinetics of blood leukocytes in mice with alloxan diabetes.
492 *Biull Eksp Biol Med* 1995,**120**:33-5.
- 493 58. Talukdar S, Oh DY, Bandyopadhyov: Neutrophils mediate insulin resistance in mice fed high fat diet
494 through secreted elastase. *Nat. Med* 2012, **18**,1407-1412.
- 495 59. Kaur G, Sarwar AM, Ather M: Nimbidum suppresses functions of macrophages and neutrophils:
496 relevance to its anti-inflammatory mechanism. *Phytotherapy Research* 2006, **18**(5): 419-424.

60. Udeinya II, Mbah AU, Chijioke GP, Shu EN: An antimalarial extract from neem leaves is antiretroviral. *Trans. Res. Soc. Trop. Med. Hyg* 2004, **98** (7). 435-437.

Table 1: Percentage composition and nutrient analysis of diets

Feed ingredients	Diets:						
	Control	VA-5%	GL-5%	VA+GL-5%	VA-7.5%	GL-7.5%	VA+GL-7.5%
Soybean meal(%)	33.78	31.03	31.03	31.03	30.53	30.53	30.53
Garri(%)	26	25	25	25	25	25	25
Maize meal(%)	38	37	37	37	35	35	35
L-Lysine(%)	0.18	0.18	0.18	0.18	0.18	0.18	0.18
L-Methionine(%)	0.17	0.17	0.17	0.17	0.17	0.17	0.17
Min/vitamin (%)	0.25	0.25	0.25	0.25	0.25	0.25	0.25
DCP(%)	2.00	2.00	2.00	2.00	2.00	2.00	2.00
Bone meal(%)	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Corn oil (%)	0.25	0.25	0.25	0.25	0.25	0.25	0.25
V. amygdalina(%)	-	5	-	2.5	7.5	-	3.75
G. latifolium(%)	-	-	5	2.5	-	7.5	3.75
Analysis:							
CP	18.40	18.31	18.40	18.34	18.47	18.46	18.48
CFAT	4.30	4.01	4.14	4.07	3.97	4.16	4.06
CFIBRE	3.71	4.27	4.15	4.21	4.58	4.41	4.49
ME	3219	3214	3218	3216	3213	3218	3216

Composition of premix:(nutrient in Amount in 2.5kg)Vit A(I.U) 12,000,000,vit D₃(I.U) 2,500,000, Vit E(mg) 20,000,vit K₃(mg) 2,000,vit B1(mg) 2,000,vit B1 (mg) 5,000,Vit B6(mg) 4,000,vit B12(mg) 15,niacin(mg) 30,000,Pantotheic acid (mg) 11,000,Folic acid(mg) 1,500,Biotin(mg) 60,Choline chloride(mg)220,000,Antioxidant(mg) 1,250,Manganase (mg) 50,000, Zinc(mg) 40,000, Iron(mg) 20,000,Copper,(mg) 3,000,Iodine (mg) 1,000,Selenium(mg) 200,Cobalt(mg) 200. **Manufactured by Megabiotics Nigeria LTD.**

Table 2: Effect of consumption of diet containing combined leaves of Vernonia amygdalina and Gongronema latifolium on hematology of diabetic rats

Treatment	WBC (10 ³ /UL)	RBC (10 ⁶ /UL)	HGB (gm/dl)	HCT (%)	PLT (10 ³ /UL)	MCV (fl)	MCH (pg)	MCHC (g/dl)
NC	6.87 ±2.28 ^a	7.01 ±0.13 ^a	15.40 ±0.04 ^a	45.13 ±0.34 ^a	6.19 ±19.08 ^a	72.75 ±1.02 ^a	25.11 ±0.68 ^a	35.20 ±5.32 ^a
DC	10.60 ±2.00 ^b	6.13 ±0.14 ^b	12.30 ±0.27 ^b	30.13 ±0.73 ^b	7.15 ±39.61 ^b	50.15 ±3.20 ^b	16.25 ±3.78 ^b	29.20 ±2.05 ^b
5%VA	8.43 ±0.24 ^c	6.55 ±0.25 ^a	12.76 ±0.27 ^a	41.83 ±0.89 ^a	5.61 ±76.53 ^a	72.82 ±1.00 ^a	26.65 ±1.00 ^a	32.00 ±2.01 ^a
5%GL	7.20 ±1.26 ^a	7.00 ±0.22 ^a	18.23 ±0.26 ^a	45.13 ±1.44 ^a	6.77 ±58.08 ^a	70.30 ±0.28 ^a	22.13 ±0.32 ^a	35.34 ±1.25 ^a
5%VA+GL	6.87 ±3.10 ^a	6.91 ±0.15 ^a	15.89 ±1.02 ^a	45.00 ±0.04 ^a	5.59 ±1.72 ^a	73.84 ±0.01 ^a	20.13 ±0.02 ^a	36.00 ±2.25 ^a

7.5%VA	6.55 ±1.27 ^a	6.58 ±0.09 ^a	13.10 ±0.13 ^a	45.10 ±0.62 ^a	6.80 ±13.19 ^a	70.25 ±0.10 ^a	26.00 ±1.40 ^a	30.21 ±1.00 ^a
7.5%GL	8.33 ±0.44 ^c	7.64 ±0.18 ^a	18.40 ±0.43 ^a	46.83 ±1.38 ^a	6.28 ±60.32 ^a	71.25 ±0.50 ^a	26.17 ±0.50 ^a	34.21 ±3.41 ^a
7.5%VA+GL	7.21 ±2.21 ^a	6.23 ±2.39 ^a	13.00 ±0.11 ^a	44.65 ±0.01 ^a	6.11 ±3.21 ^a	70.00 ±4.21 ^a	26.00 ±2.32 ^a	31.76 ±0.00 ^a
INSD	7.43 ±0.33 ^a	6.48 ±0.33 ^a	12.83 ±0.58 ^a	41.73 ±1.45 ^a	6.04 ±18.21 ^a	67.28 ±0.28 ^a	20.05 ±1.26 ^a	30.11 ±1.25 ^b

Means within the same column with different superscript are significantly different (P<0.05)

512

Table 3: Effect of consumption of diet containing combined leaves of *Vernonia amygdalina* and *Gongronema latifolium* on differential white blood and CD4⁺ cells count of diabetic rats

Treatment	Neutrophil(%)	Esenophil(%)	Basophil(%)	lymphocyte(%)	monocyte(%)	CD4 ⁺ cell count(10 ⁶ /L
NC	23.00 ±1.93 ^a	2.67 ±0.56 ^a	0.00 ±0.00	70.67 ±2.19 ^a	4.67±0.33 ^a	15.70 ±0.36 ^a
DC	30.33 ±4.95 ^b	1.33±0.55 ^b	1.81 ±0.00	43.33 ±4.21 ^b	5.40±0.00 ^b	51.53±0.77 ^b
5%VA	23.66 ±0.91 ^a	3.00 ±0.73 ^a	0.00 ±0.00	73.33 ±1.64 ^a	0.00 ±0.00	0.00 ±0.00
5%GL	24.33 ±4.07 ^a	1.66±0.76 ^a	0.00 ±15.38	58.66 ±14.15 ^a	4.00 ±0.00 ^a	19.29±0.47 ^a
5%VA+ GL	22.95±1.06 ^a	2.78 ±1.43 ^a	0.00 ±0.00	70.77 ±2.30 ^a	4.00±1.21 ^a	13.33 ±3.42 ^a
7.5%VA	29.00 ±1.34 ^b	3.50 ±1.11 ^a	0.00 ±0.0	67.50±0.22 ^a	0.00 ±0.00	18.95±0.42 ^a
7.5%GL	22.00 ±2.22 ^a	1.66 ±0.21 ^a	0.00 ±0.00	61.33 ±2.43 ^a	4.00 ±0.00 ^a	20.25±0.77 ^a
7.5%VA+ GL	23.01 ±1.21 ^a	1.98 ±1.00 ^a	0.00 ±0.00	64.76 ±1.32 ^a	4.00 ±0.00 ^a	19.00±0.10 ^a
INSD	26.00 ±5.17 ^b	5.00 ±0.73 ^d	0.00 ±0.00	59.00 ±0.22 ^a	0.00 ±0.00	17.87 ±0.51 ^a

Means within the same column with different superscript are significantly different (P<0.05)

516

517