

New Formulas For One-Dimensional Differential Transform

Abstract

In this work, we introduced new useful formulas for one-dimensional differential transform and applied the differential transform method to selected linear ordinary differential equations. This study showed that this method is powerful and efficient in finding series solutions for linear differential equations and capable of reducing the size of calculations comparing with other methods.

Keywords: *Differential transforms, Differential transform method, Taylor series method, Differential equations.*

1. Introduction

The idea of differential transform presented by G. E. Pukhov [1] in 1982 and the concept of differential transform proposed first by Zhou [2] in 1986 when applied to solve linear and nonlinear initial value problems in electric circuit analysis. The (Pukhov-Zhou) differential transform method presented as a new method based on Taylor series[3-14] and consider as a semi-analytical technique uses the Taylor series to construct the solutions of differential equations in the form of a power series. This method represents an iterative procedure for obtaining analytic series solutions of differential equations and useful for obtaining exact and approximate solutions of linear ordinary differential equations [4-6] and system of linear ordinary differential equations [8-9]. The main aims in this work are to introduce new useful algorithms for one-dimensional differential transform and applied the differential transform method to selected linear ordinary differential equations.

2. The One-Dimensional Differential Transform

In this section, we introduce the concept of one-dimensional differential transform and review some basic fundamental theorems [4-13]. To do that, we assume that $f(x)$ be a $C^\infty(I)$ function in an open interval of R , and x_0 be any point of I . Then the Taylor series of $f(x)$ about x_0 can be represented by

$$f(x) = \sum_{k=0}^{\infty} \left[\frac{f^{(k)}(x)}{k!} \right]_{x=x_0} (x-x_0)^k \quad (2.1)$$

Note that, throughout this work, we assume that $f(x)$ is an analytic function at every point of I .

Definition (1): Let $f(x)$ be an analytic function about x_0 , then the k^{th} order differential transform of $f(x)$ is defined as

$$D_T \{f(x)\} := \left[\frac{f^{(k)}(x)}{k!} \right]_{x=x_0} . \quad (2.2)$$

37

38 In definition (2.1), $D_T \{f(x)\}$ represents the one-dimensional differential transform of $f(x)$ about x_0 and it
39 usually denoted by $F(k)$ and throughout our work, we will take $x_0 = 0$, which reduces the definition (2.1) to

40

$$41 \quad D_T \{f(x)\} = F(k) := \frac{f^{(k)}(0)}{k!}. \quad (2.3)$$

42

43 **Definition (2):** The inverse differential transform of $F(k)$ denotes by $D_T^{-1}\{F(k)\}$ and it defines as

44

$$45 \quad D_T^{-1}\{F(k)\} = f(x) := \sum_{k=0}^{\infty} F(k)x^k \quad (2.4)$$

46

47 For example $D_T \{e^{\alpha x}\} = \alpha^k / k!$, $D_T \{\cos \alpha x\} = (\alpha^k / k!) \cos(k\pi/2)$, respectively, where, α is a constant.

48

49 **Theorem (1):** Let $f(x)$ and $g(x)$ be analytic functions, with differential transforms $F(k)$ and $G(k)$
50 respectively, then

51

$$52 \quad D_T \{\alpha f(x) + \beta g(x)\} = \alpha D_T \{f(x)\} + \beta D_T \{g(x)\}, \quad (2.5)$$

53

$$54 \quad D_T^{-1}\{\alpha F(k) + \beta G(k)\} = \alpha D_T^{-1}\{F(k)\} + \beta D_T^{-1}\{G(k)\}, \quad (2.6)$$

55

56 where α and β are constants. The proof of the linearity property follows immediately from the definitions (1)
57 and (2).

58

59 **Theorem (2):** Let $f(x)$ be an analytic function, with $D_T \{f(x)\} = F(k)$, then

60

$$61 \quad D_T \left\{ \frac{d^n f(x)}{dx^n} \right\} = \frac{(k+n)!}{k!} F(k+n) \quad (2.7)$$

62

63 Hence,

64

$$65 \quad D_T \left\{ \frac{d^n f(x)}{dx^n} \right\} = D_T \left\{ \sum_{k=0}^{\infty} \frac{f^{(k+n)}(0)}{k!} x^k \right\} = D_T \left\{ \sum_{k=0}^{\infty} \frac{(k+n)!}{k!} \frac{f^{(k+n)}(0)}{(k+n)!} x^k \right\}$$

$$66 \quad = \frac{(k+n)!}{k!} \frac{f^{(k+n)}(0)}{(k+n)!} = \frac{(k+n)!}{k!} F(k+n). \quad (2.8)$$

67

68 **Theorem (3):** Let $F(k)$ be the differential transform of $f(x) = x^m$, then $F(k) = \delta_{km}$, where, δ_{km} is the

69

70 Kronecker delta. The proof of this theorem follows immediately from identity $x^m \equiv \sum_{k=0}^{\infty} \delta_{km} x^k$.

Theorem (4): Let $f(x)$ and $g(x)$ be analytic functions and

$$f(x) = \int_0^x g(t) dt, \quad (2.9)$$

then the differential transform of the function $f(x)$ is given by

$$D_T \{f(x)\} = \frac{G(k-1)}{k}, \quad (2.10)$$

where $G(k)$ is the differential transform of $f(x)$. Hence, applying the Leibniz's rule on the equation (2.9) gives, $f'(x) = g(x)$, and this leads to the relation

$$\frac{g^{(k)}(0)}{k!} = \frac{(k+1)!}{k!} \frac{f^{(k+1)}(0)}{(k+1)!} = (k+1) \frac{f^{(k+1)}(0)}{(k+1)!}.$$

This can be written in the form

$$\frac{f^{(k)}(0)}{k!} = \frac{1}{k} \frac{g^{(k-1)}(0)}{(k-1)!},$$

which his yields the formula (2.10).

Theorem (5): Let $f_1(x)$ and $f_2(x)$ be analytic functions, with $f(x) = f_1(x) \cdot f_2(x)$, then

$$D_T \{f(x)\} = \sum_{n=0}^k F_1(n) F_2(k-n), \quad (2.11)$$

where $F_1(k)$ and $F_2(k)$ are the differential transforms of the functions $f_1(x)$ and $f_2(x)$ respectively. To prove (2.11), we first write

$$\begin{aligned} f_1(x) f_2(x) &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{f_1^{(n)}(0)}{n!} \frac{f_2^{(m)}(0)}{m!} \frac{x^{n+m}}{n!m!} = \sum_{k=0}^{\infty} \sum_{n=0}^k \frac{f_1^{(n)}(0)}{n!} \frac{f_2^{(k-n)}(0)}{(k-n)!} \frac{x^k}{n!(k-n)!} \\ &= \sum_{k=0}^{\infty} \sum_{n=0}^k \frac{f_1^{(n)}(0)}{n!} \frac{f_2^{(k-n)}(0)}{(k-n)!} \frac{x^n x^{k-n}}{n!(k-n)!} \end{aligned}$$

Next, applying the differential transform on both sides of the above equation yields the formula (2.11).

3. New Formulas

In this section, we introduce new basic formulas for the one-dimensional differential transform.

Theorem (6): Let $f(x)$, be an analytic function, with $D_T \{f(x)\} = F(k)$, then

$$D_T \{x^m f^{(n)}(x)\} = \sum_{i=0}^k \delta_{im} \frac{(k+n-i)!}{(k-i)!} F(k+n-i) \quad (3.1)$$

To prove the formula (3.1), we make use of (8) and $D_T \{x^m\} = \delta_{km}$. This enables us to write

$$F_1(k) = D_T \{x^m\} = \delta_{km} \quad F_2(k) = D_T \{f^{(n)}(x)\} = \frac{(k+n)!}{k!} F(k+n). \quad (3.2)$$

Then, the formula (2.11) leads to

$$D_T \{x^m f^{(n)}(x)\} = \sum_{i=0}^k F_1(i) F_2(k-i) \quad (3.3)$$

Finally, if we substitute (3.2) into (3.3), we can establish the formula (3.1). Note that, for $m=0$, the formula (3.1) reduces to the formula (2.7).

Corollary (1): Let $f(x)$, be an analytic function, with $D_T \{f(x)\} = F(k)$, then

$$D_T \{x^n f^{(n)}(x)\} = \prod_{i=0}^{n-1} (k-i) F(k). \quad (3.4)$$

To prove (3.4), we put $m=n$ into the formula (3.1). This gives

$$D_T \{x^n f^{(n)}(x)\} = \frac{k!}{(k-n)!} F(k). \quad (3.5)$$

Next making use, of the identity

$$k! = \frac{(k-n+1)!}{(k-n+1)(k-n)(k-n-1)(k-n-2)\dots(k-1)}, \quad (3.6)$$

we can easily establish the formula (3.4).

Corollary (2): Let $f(x)$ be an analytic function, with $D_T \{f(x)\} = F(k)$, then

$$D_T \{e^{\alpha x} f^{(n)}(x)\} = \sum_{i=0}^k \frac{\alpha^i}{i!} \frac{(k+n-i)!}{(k-i)!} F(k+n-i) \quad (3.7)$$

To prove (3.7), we make use of (2.2) and (2.7) to write

$$F_1(k) = D_T \{e^{\alpha}\} = \frac{\alpha^k}{k!}, \quad F_2(k) = D_T \{f^{(n)}(x)\} = \frac{(k+n)!}{k!} F(k+n). \quad (3.8)$$

142

143 Then, the formula (2.11) leads to

144

$$D_T \{e^{\alpha} f^{(n)}\} = \sum_{i=0}^k F_1(i) F_2(k-i) = \sum_{i=0}^k \frac{\alpha^i}{i!} \frac{(k+n-i)!}{(k-i)!} F(k+n-i) \quad (3.9)$$

146

147 **Theorem (7):** Let $f(x)$ be an analytic function, with $D_T \{f(x)\} = F(k)$, then

$$D_T \left\{ \left(x f^{(n)}(x) \right)' \right\} = \frac{(k+1)(k+n)!}{k!} F(k+n) \quad (3.10)$$

149

150 We can establish the proof of the formula (3.9) by using (2.4), (6) and (2.11) as following

151

$$\begin{aligned} D_T \left\{ \left(x f^{(n)}(x) \right)' \right\} &= D_T \{x f^{(n+1)}(x)\} + D_T \{f^{(n)}(x)\} = \frac{(k+n)!}{(k-1)!} F(k+n) + \frac{(k+n)!}{k!} F(k+n) \\ &= \frac{(k+n)!}{(k-1)!} \left[1 + \frac{1}{k} \right] F(k+n) = \frac{(k+1)(k+n)!}{k!} F(k+n). \end{aligned}$$

155

156 Note that, for $n=1$ the formula (3.10), reduce to the formula

157

$$D_T \left\{ \left(x f'(x) \right)' \right\} = (k+1)^2 F(k+1) \quad (3.11)$$

159

160 **Theorem (8):** Let $f(x)$ be an analytic function with $D_T \{f(x)\} = F(k)$, then

161

$$D_T \left\{ \left(x^m f^{(n)}(x) \right)' \right\} = \frac{(k+1)(k+n-m+1)!}{(k-m+1)!} F(k+n-m+1) \quad (3.12)$$

163

164 We can establish the proof of the formula (3.12) by using (2.5) and (3.1) as

165

$$\begin{aligned} D_T \left\{ \left(x^m f^{(n)}(x) \right)' \right\} &= D_T \{x^m f^{(n+1)}(x)\} + m D_T \{x^{m-1} f^{(n)}(x)\} \\ &= \frac{(k+n+1-m)!}{(k-m)!} F(k+n+1-m) + m \frac{(k+n-m+1)!}{(k-m+1)!} F(k+n-m+1) \end{aligned}$$

169

170 Now by setting $h = m-1$, we can write

171

$$D_T \left\{ \left(x^m f^{(n)}(x) \right)' \right\} = \frac{(k+n-h)!}{(k-h-1)!} F(k+n-h) + (h+1) \frac{(k+n-h)!}{(k-h)!} F(k+n-h)$$

173

$$\begin{aligned}
 &= \frac{(k+n-h)!}{(k-h-1)!} F(k+n-h) + \frac{(h+1)}{(k-h)} \frac{(k+n-h)!}{(k-h-1)!} F(k+n-h) \\
 &= \left(1 + \frac{(h+1)}{(k-h)} \right) \frac{(k+n-h)!}{(k-h-1)!} F(k+n-h) \\
 &= \left(\frac{(k+1)}{(k-h)} \right) \frac{(k+n-h)!}{(k-h-1)!} F(k+n-h) = \frac{(k+1)(k+n-h)!}{(k-h)!} F(k+n-h).
 \end{aligned}$$

This yields the formula (3.12).

4. Numerical Examples

In this section, we apply the differential transform method to selected linear differential equations and compare our results with the results obtained by the Taylor series method.

Example (1)

Consider the initial value problem

$$y'' - 2xy' + 2ny = 0, \quad (4.1)$$

$$y(0) = c_0, \quad y'(0) = c_1, \quad (4.2)$$

The differential equation (4.1) is known as Hermit equation, where n is usually a non-negative integer. To solve the initial value problem (4.1-2), we apply the differential transform to both sides of (4.1). This gives

$$D_T \{y''\} - 2D_T \{xy'\} + 2nD_T \{y\} = 0. \quad (4.3)$$

Next, making use of the formulas (2.7) and (3.1) enables us to find

$$(k+1)(k+2)Y(k+2) - 2kY(k) + 2nY(k) = 0. \quad (4.4)$$

This leads to the iteration formula

$$Y(0) = c_0, \quad Y(1) = c_1, \quad Y(k+2) = \frac{2(k-n)}{(k+1)(k+2)} Y(k), \quad k = 0, 1, 2, \dots \quad (4.5)$$

Now, the iteration formula (4.5) gives

$$Y(2) = \frac{-2n}{2!} c_0, \quad Y(3) = \frac{-2(n-1)}{3!} c_1, \quad Y(4) = \frac{2^2 n(n-2)}{4!} c_0, \quad (4.6)$$

$$Y(5) = \frac{2^2(n-1)(n-3)}{5!}c_1, \quad Y(6) = \frac{-2^3n(n-2)(n-4)}{6!}c_0.$$

Therefore, this leads to the series solution

$$y(x) = c_0 \left(1 - \frac{2n}{2!}x^2 + \frac{2^2n(n-2)}{4!}x^4 - \frac{2^3n(n-2)(n-4)}{6!}x^6 \dots \right) + c_1 \left(x - \frac{2(n-1)}{3!}x^3 + \frac{2^2(n-1)(n-3)}{5!}x^5 + \dots \right). \quad (4.7)$$

This is the Hermit polynomial given in ref. [15].

Example (2)

Consider the initial value problem

$$(1-x^2)\frac{d^2y}{dx^2} - 2x\frac{dy}{dx} + n(n+1)y = 0, \quad (4.8)$$

$$y(0) = c_0, \quad y'(0) = c_1, \quad (4.9)$$

The self-adjoint equation (4.8) is known as the Legendre equation, where, c_0 and c_1 are constant and n is an integer. To solve the initial value problem (4.8-9), we apply the differential transform to both sides of (4.8). This gives

$$D_T \{y''\} - D_T \{(x^2y')'\} + D_T \{n(n+1)y\} = 0. \quad (4.10)$$

Next, making use of the formulas (2.7) and (3.12) enables us to find

$$(k+1)(k+2)Y(k+2) - k(k+1)Y(k) + n(n+1)Y(k) = 0, \quad (4.11)$$

This leads to the iteration formula

$$Y(0) = c_0, \quad Y(1) = c_1 \quad (4.12)$$

$$Y(k+2) = \frac{1}{(k+1)(k+2)} [k(k+1) - n(n+1)]Y(k), \quad k = 0, 1, 2, \dots$$

Now, the iteration formula (4.12) gives

$$Y(2) = -\frac{n(n+1)}{2!}c_0, \quad Y(3) = -\frac{(n-1)(n+2)}{3!}c_1, \quad Y(4) = \frac{(n-2)n(n+1)(n+3)}{4!}c_0, \quad (4.13)$$

$$Y(5) = \frac{(n-3)(n-1)(n+2)(n+4)}{5!}c_1, \quad Y(6) = -\frac{(n-4)(n-2)n(n+1)(n+3)(n+5)}{6!}c_0,$$

248

$$Y(7) = -\frac{(n-5)(n-3)(n-1)(n+2)(n+4)(n+6)}{7!}c_1, \dots$$

250

251 Therefore, this leads to the series solution

252

$$y(t) = c_0 \left[1 - \frac{n(n+1)}{2!}x^2 + \frac{(n-2)n(n+1)(n+3)}{4!}x^4 - \frac{(n-4)(n-2)n(n+1)(n+3)(n+5)}{6!}x^6 + \dots \right] +$$

254

$$c_1 \left[x - \frac{(n-1)(n+2)}{3!}x^3 + \frac{(n-3)(n-1)(n+2)(n+4)}{5!}x^5 - \right.$$

256

$$\left. \frac{(n-5)(n-3)(n-1)(n+2)(n+4)(n+6)}{7!}x^7 + \dots \right] \quad (4.14)$$

258

259 This result is identical to the result obtained by the Taylor series procedure applied in ref. [16].

260

261 **Example (3)**

262 Consider the initial value problem

263

$$y' - 2xy = 0, \quad (4.15)$$

265

$$y(0) = c_0. \quad (4.16)$$

267

268 To solve the initial value problem (4.15-16), we make use of the formulas (7), (11) and (13). This gives

269

$$(k+1)Y(k+1) - 2 \sum_{i=0}^k \delta_{i1} Y(k-i) = 0. \quad (4.17)$$

271

272 Now, we can write the iteration formula

$$Y(0) = c_0, \quad Y(k+1) = \frac{2}{k+1} \sum_{i=0}^k \delta_{i1} Y(k-i), \quad k = 0, 1, 2, \dots \quad (4.18)$$

274

275 The iteration formula (41) for $k = 0, 1, 2, \dots$ gives

276

$$Y(0) = c_0, \quad Y(1) = 0, \quad Y(2) = c_0, \quad Y(3) = 0,$$

278

$$Y(4) = \frac{c_0}{2!}, \quad Y(5) = 0, \quad Y(6) = \frac{c_0}{3!}, \dots$$

280

281 Therefore, the close form solution can be easily in the form written as

282

$$y(x) = \sum_{k=0}^{\infty} Y[k]x^k = c_0 \left(1 + x^2 + \frac{1}{2!}x^4 + \frac{1}{3!}x^6 + \frac{1}{4!}x^8 + \dots \right) = c_0 \sum_{k=0}^{\infty} \frac{1}{k!}x^{2k}.$$

This represents the Taylor series expansion of the exact solution $y(x) = c_0 e^{x^2}$.

5. Conclusion

In this work, we introduced the concept of differential transform and studied some of their properties. As a contribution, we introduced and proved new formulas for the one-dimensional differential transform and applied the differential transform method to selected linear ordinary differential equations. This study showed that the differential transform method is powerful and efficient techniques in finding analytical solutions for linear differential equations. It also showed that this technique is capable of reducing the size of calculations comparing with the Taylor series method.

References

1. Pukhov G., Differential Transforms And Circuit Theory, Circuit Theory and Applications, VOL. 10, 265-276, 1982.
2. Zhou J.K., Differential Transformation and its Applications for Electrical Circuits, Huarjung University Press, Wuuhahn, China, 1986.
3. Bervillier C., Status of the differential transformation method, Original Research Article Applied Mathematics and Computation, Volume 218, Issue 20, 15 June, 10158-10170, 2012.
4. Batiha K. and Batiha B., A new Algorithm for Solving Linear Ordinary Differential Equations, World Applied Sciences Journal 15 (12): 1774-1779, 2011.
5. Erturk V., Application of Differential Transformation Method to Linear Sixth-Order Boundary Value Problems, Applied Mathematical Sciences, Vol. 1, No. 2, 51-58, 2007.
6. Hassan I. and Erturk V., Solutions of Different Types of the Linear and Nonlinear Higher-Order Boundary Value Problems by Differential Transformation Method, European Journal of Pure and Applied Mathematics, Vol. 2, No. 3, 426-447, 2009.
7. Erturk V., Approximate Solutions of a Class of Nonlinear Differential Equations by Using Differential Transformation Method, International Journal of Pure and Applied Mathematics, Vol. 30 No. 3, 403-409, 2006.
8. Thongmoona M. and Pusjuso S., The Numerical Solutions of Differential Transform Method and the Laplace Transform Method for a System of Differential Equations, Nonlinear Analysis: Hybrid Systems 4, 425-431, 2010.
9. Mirzaee F., Differential Transform Method for Solving Linear and Nonlinear Systems of Ordinary Differential Equations, Applied Mathematical Sciences, Vol. 5, No. 70, 3465-3472, 2011.
10. Biazar J. and Eslami M., Differential Transform Method for Quadratic Riccati Differential Equation, International Journal of Nonlinear Science, Vol.9, No.4, 444-447, 2010.
11. Ali J., One Dimensional Differential Transform Method for Some Higher Order Boundary Value Problems in Finite Domain, Int. J. Contemp. Math. Sciences, 263-272, 2012.
12. Raslan K. and Zain F. Abu Sheer, Differential Transform Method for Solving Non-linear Systems of Partial Differential Equations, International Journal of Physical Sciences, Vol. 8 (38), 1880-1884, 16, 2013.
13. Chang S. and Chang I., A new Algorithm for Calculating One-Dimensional Differential Transform of Non linear Functions, Applied Mathematics and Computation 195, 799-808, 2008.

- 324 14. Zill D., A First Course in Differential Equations With Applications, Prindle, Weber and Schmidt Publishers,
325 1982.
- 326 15. Apostol T., Mathematical Analysis, Addison-Wesley Publishing Company, 1974.
- 327 16. S. Ross L., Differential Equations, John Wiley & Sons, Inc., 1984.
- 328