

Keywords: *Morula* fruit, storage temperature, cultivars, chilling injury, postharvest shelf life

Fruits are living tissues and are diverse in morphology, structure, composition and general physiology. Due to high moisture content, active metabolisms, tender nature and rich in nutrients, they are vulnerable to dehydration, physiological disorders, environmental stress, mechanical injury and pathological breakdown therefore usually considered to be highly perishable [1]. Therefore, these characteristics limit the storage life of the fruits and vegetables and cause significant deterioration following harvest [2]. Postharvest loss can occur at any point in production and market chain. It is estimated that the magnitude of these

31 losses due to inadequate postharvest handling and storage in fresh fruits and vegetables is
 32 relatively high but varies in accordance with the country and commodity [1]. In a hungry and
 33 increasingly competitive world, reducing postharvest food losses is a major agricultural goal.
 34 In order to reduce these losses, postharvest technology which delays senescence and maintain
 35 quality must be applied. These technology systems include temperature management,
 36 controlled atmosphere and modified atmosphere storages.



37 Temperature being the most important environmental factor that influences the deterioration
 38 of harvested horticultural commodities, most perishable commodities last longest at
 39 temperature near 0 °C. Temperature influences how other internal and external factors
 40 influence the commodity [3]. Temperature management is the most effective tool for
 41 maintaining quality and safety, and for extending the postharvest life of fresh horticultural
 42 commodities. After harvest, most fruits including marula remain alive and temperature during
 43 storage plays an important role in their quality. Ripening of the fruit can be slowed down to
 44 extend shelf life by reducing the fruit temperature. Low temperature storage is considered the
 45 most effective method of extending postharvest life and maintaining the quality of the fruits
 46 [4]. However, most fruits that originated from the tropical and subtropical regions are chilling
 47 sensitive when exposed to low temperatures that are above the freezing point [5]. Such crops
 48 including marula when stored below its critical temperature generally 10-13 °C for most
 49 varieties are injured. Damage often is induced by a very brief exposure, but may not become
 50 apparent for several days or until transfer to warmer temperature depending on the duration
 51 and severity of chilling. Due to sensitivity of tropical and some sub-tropical produce to
 52 chilling injury, their transport to distant markets is generally only successful by air.
 53 Therefore, advances in the improvement of fruits to resist low temperature storage could be
 54 enhanced to prolong their shelf life even their trade over long distance. However, it has been
 55 suggested that the temperature range of 13 °C is adequate to avoid the onset of chilling injury
 56 in tropical and some sub-tropical horticultural produce for at least three weeks
 57 [6]. There is hardly any literature available on the storage temperature of marula fruit, except
 58 it was reported that marula juice stores well at 4 °C [7]. Therefore, the objective of this study
 59 was to evaluate the effects of storage temperature on postharvest quality of marula fruit.



60 2.0 MATERIALS AND METHODS

61 2.1 Experimental site

The experiment was conducted in the Crop Science and Production Department Laboratories of the Botswana College of Agriculture between January and April 2015. Botswana College of Agriculture is located at Sebele (latitude 24 34⁰S and longitude 25 57⁰E, altitude of 994 m above sea level) and 10 km from Gaborone.

2.2 Experimental design

The experimental design was a completely randomized design (CRD). Fresh marula fruits were picked randomly at the green mature stage (physiological maturity) from 10 different marula trees around Sebele. The fruits were stored at 0, 4, 8 and 12°C. Four refrigerators were set at different temperatures of 0, 4, 8 and 12 °C ± 1°C. Fresh fruits were collected when still firm, green, uniform and free from bruises and defects as judged subjectively based on epidermal colour. Fruits were washed to remove soil and other external material. Three hundred fruits were stored in each storage temperature. All analyses were done in triplicate.

2.3 Dependent variables determined

The dependent variables analysed were fruit development, fruit chlorophyll content, fruit anthocyanin and carotenoid content, fruit juice soluble solids content (SSC), total titratable acidity (TTA), vitamin C content, chilling injury (incidence and severity) and storage life of the fruit in cold storage and after storage at room temperature (25 ± 3 °C).

2.3.1 Titratable acidity

The fruit and skin was cut and 100 g of sample weighed, then 200 ml of distilled water was added to the sample. The mixture was then blended for five minutes. The sample was homogenized and filtered through 5-layers of cheese cloth. 20 ml of filtrate (juice) was then placed in a 50-ml conical flask and two drops of 1% phenolphthalein indicator was added. The samples were titrated with 0.1 N NaOH to end point and this was done in triplicates. The results were expressed as total titratable acidity equivalents using the formula given below.

Total titratable acidity (g/100 ml juice) = $\frac{\text{ml of base} \times \text{normality base} \times 1 \times 100 \times 2}{\text{ml sample}}$

ml sample

2.3.2 Soluble solids content

89 Soluble solids content (SSC) was determined using a refractometer. Ten fruits per treatment
90 per replication were pinched using a razor blade and the juice squeezed directly into the hand
91 refractometer (0-32°) and then the average sugar content was determined in °Brix.

92 2.3.3 Vitamin C (ascorbic acid) determination

93 Vitamin C content was determined using 2, 6-dichloroindophenol titrimetric method [8].

94 Ascorbic acid reduces oxidation-reduction indicator dye (2, 6-dichloroindophenol) to
95 colourless solution. At end point, excess unreduced dye is rose pink in acid solution. Vitamin

96 C was extracted and titration performed in the presence of metaphosphoric acid-acetic acid
97 solution to maintain proper acidity for reaction and to avoid autooxidation of ascorbic acid at
98 high pH [8].

99 2.3.4 Fruit chlorophyll, anthocyanin and carotenoid contents

100 Anthocyanin plus carotenoids and chlorophyll content of the fruit skin were measured from
101 two discs (11 mm diameter) per fruit (10 marula fruits per replicate/treatment), cut from the
102 yellowish and greenest portions using a cork borer and the scalpel. The 10 discs were then
103 extracted in 4 ml of 0.1 N HCL in methanol at 25 ± 2 °C in the dark for 24 hours. Absorbance
104 of extracts was measured using a UV Visible spectrophotometer (UV-1602 I PC, Shimadzu,
105 RSA Pty Ltd.Co.Reg.NO: 90/07641/07) at various wavelengths of 530, 620, 645, 650 and 663
106 nm.

107 Anthocyanins plus carotenoid content was determined according to an established method
108 [9], using a molar extinction coefficient of 4.62×10^4 [10].

109 Anthocyanins plus carotenoids (nmole/cm² marula skin) = $\frac{[(A_{530}-A_{620}) - 0.1(A_{650}-A_{620})]}{4.62 \times 10^4}$

110

111 Skin chlorophyll content was measured as absorbance at 663 nm [11]. The following
112 equation was used to calculate the relative total chlorophyll content [11].

113 Chlorophyll (mg/cm² of morula skin) = $24.88 \times A_{653}$

114 2.3.5 Chilling injury (incidence and severity)

115 Chilling injury was evaluated daily for incidence and severity of chilling in storage. Chilling
116 injury (CI) incidence was determined from a sample of 20 fruit/replicate/treatment. Fruit

showing symptoms of CI were counted and expressed as a percentage. Chilling severity was evaluated on a predetermined scale form: 0 being no injury; 1 being slight injury (where 1/3 of the fruit showed some injury symptoms); 2 moderate injuries (where 2/3 of the fruit showed some injury symptoms) and 3 was severe injury (determined by more than 2/3 of the fruit showing injury depending on the peel damage)



Chilling injury index was calculated by the following equation;

$$CII = [(no\ of\ fruits\ with\ no\ injury \times 0) + (no\ of\ fruits\ with\ slight\ injury \times 1) + (no\ of\ fruits\ with\ moderate\ injury \times 2) + (no\ of\ fruits\ with\ severe\ injury \times 3)] \div no\ of\ fruits\ sampled.$$

2.3.6 Fruit development

Fruit development was measured from 10 tagged fruits every week starting from one week after full bloom to 21 weeks after full bloom using a digital electronic vernier calliper.

2.3.6 Data analysis

Analysis of variance was performed on the data collected using the general linear models, procedure of the Statistical Analysis System (SAS) Programme. Multiple comparisons among means was performed using the Protected Least Significant Difference at $P=0.05$.




3.0. RESULTS

3.1. Fruit development




The results of this study showed that marula fruit growth and development showed a typical simple sigmoid growth curve (Figure 1). Fruit diameter increased with time and reached maximum diameter at 18 weeks after full bloom (Figure 1).




138

139 3.2 Chilling injury

140 The results of this study showed that storage temperature significantly ($P \leq 0.05$) influenced
 141 the incidence (Figure 2) and severity (Figure 3) of chilling injury of marula fruit. All marula
 142 fruit stored at temperatures $\leq 8^\circ\text{C}$ developed chilling injury, but the severity was
 143 significantly ($P \leq 0.01$) greater on fruit stored at 0 and 4°C (Figure 3). Marula fruit storage at
 144 8°C had severity index of ≥ 2 (67 % or above of the fruit is chilled) after 15 days of storage,
 145 while fruit stored at 0 and 4°C attained a severity index ≥ 2 after 7 and 8 days, respectively
 146 (Figure 3). Marula fruit stored at 12°C did not suffer from chilling injury (Figure 2, 3).




147




148

149 3.3 Titratable acidity

150 During fruit growth and development, titrat^{gh} acidity increased with time and a reached a
 151 maximum after 21 weeks after full bloom and decreased thereafter (Figure 4). During storage
 152 for 8 and 16 days total titratable acidity significantly ($P \leq 0.05$) decreased with increase in
 153 storage temperature (Figure 5). Storage temperature and duration of storage significantly ($P \leq$
 154 0.05) influenced the decrease in fruit titratable acidity (Figure 5). Fruit stored for 8 days had
 155 significantly ($P \leq 0.05$) higher titratable acidity than fruit stored for 16 days at 0, 4, 8 and 12
 156 °C, respectively (Figure 5).






157





3.4 Total soluble solids

During fruit growth and development, the fruit total soluble solids content increased with time (Figure 6). During storage at room temperature (25 ± 3 °C) fruit soluble solids content (SSC) increased and reached a maximum after six days in storage, thereafter the fruit SSC decreased (Figure 7). Storage temperature and duration of storage significantly ($P \leq 0.05$) influenced the marula fruit SSC (Figure 8). As storage temperature increased from 0 to 12 °C, fruit SSC significantly ($P \leq 0.05$) increased, irrespective of storage duration (Figure 8). Fruit stored for 16 days had significantly ($P \leq 0.05$) higher SSC than fruit stored for 8 days (Figure 8).




3.5 Fruit anthocyanin and carotenoid content (colour)

The fruit colour (anthocyanin and carotenoid) content increased significantly ($P \leq 0.05$) with increase in storage temperature (Figure 9). After 22 days of storage at lower temperatures ($\leq 12^\circ\text{C}$), maximum fruit colour developed in fruit stored at 12°C (Figure 9). Though fruit stored at $25 \pm 3^\circ\text{C}$, developed the best colour, but it was not significantly different from fruit stored at 8 and 12°C (Figure 9). Maximum fruit colour was attained after 8 and 25 days of storage at 25 ± 3 and 12°C , respectively (Figure 9).




3.6 Fruit chlorophyll content

The results of the current study showed that during fruit growth and development, chlorophyll content was high but significantly ($P \leq 0.05$) decreased as fruit maturation increased (Figure 10). In storage, storage temperature significantly ($P \leq 0.05$) affected fruit chlorophyll content (Figure 11). Increase in storage temperature from 0 to 25 °C significantly ($P \leq 0.05$) increased the degradation of fruit chlorophyll content of marula fruit (Figure 11).



187



188

189 3.7 Vitamin C content

190 The vitamin C content of marula fruit was significantly ($P \leq 0.05$) influenced by storage
 191 temperature (Figure 12). Retention of marula fruit vitamin C content was significantly ($P \leq$
 192 0.05) increased as storage temperature was lowered (Figure 12). The vitamin C content of
 193 marula fruit stored at $25 \pm 3^\circ\text{C}$ for 8 days was 1.35 times lower than the vitamin C content of
 194 fruit stored at 0°C for 25 days (Figure 12).

195 4.0 DISCUSSION

196 4.1 Fruit Development

The results of this study showed that marula fruit displayed a simple sigmoid growth curve. The increase in fruit size was attributed to cell division in the first four weeks after full bloom and then followed by cell elongation and expansion [12, 13, 14, 15]. Fruit can increase in mass or volume by 100-fold or more from fertilization to maturity and such changes follow simple or double-sigmoid growth curve depending on the fruit species [13, 16].

4.2 Effect of storage temperature on postharvest quality of marula fruits

Marula is subtropical in origin, however, most fruits that are of tropical or subtropical in origin are reported to be chilling sensitive [5, 17, 18]. These fruits are injured after a period of exposure to chilling temperature below 10-15°C, but above their freezing temperatures. The results of this study showed that storage temperature significantly influenced the development of chilling injury (incidence) and severity of chilling injury in marula fruit. As storage temperature decreased below 12°C, the incidence and severity of chilling injury increased. Chilling injury symptoms of skin pitting, poor colour development, poor aroma and surface lesions developed in marula fruit stored at temperatures $\leq 12^{\circ}\text{C}$, though severity varied with storage temperature and duration of storage. Marula fruit stored at 0, 4 and 8 °C developed chilling injury symptom after 7 and 9 days of storage, but severity of chilling was high in fruit stored at 0 and 4 °C. Marula fruit stored at 12 °C did not develop chilling injury. Similar results have been reported [2, 17], who all reported of chilling injury symptoms such as scalding, surface pitting, poor aroma, uneven ripening, poor colour development and increased susceptibility to diseases in mango stored at temperatures below 12 °C. Chilling injury can be related to the enzymatic browning of activities of polyphenoloxidase (PPO) and peroxide (POD) and the increase of phenolic compounds [19]. Chilling injury is known to significantly change the microstructure of the tissue which in severe cases may lead to tissue breakdown due to failure to carry normal metabolic processes [20]. Various physiological, biochemical alteration and cellular dysfunction occur in chilling sensitive species in response to chilling stress [20]. These alterations include increased membrane permeability and alteration of activities of membrane proteins. If chilling stress is prolonged, these alterations cascade to development of chilling injury symptoms such as skin surface pitting, sunken or surface lesions, uneven ripening, pulp discoloration, greyish-scald discoloration of the skin, water-soaking of the tissue, off-flavour, susceptibility to fungal decay, reduced aroma, carotenoids [21, 22] and become more serious when fruits are transferred to room temperature probably because they satisfactorily ripen at 21-24 °C. Factors that have been reported to influence the susceptibility of horticultural produce to chilling injury include

environmental conditions, maturity when picked and postharvest handling techniques, and duration of exposure to the chilling temperature [2, 23].

4.3 Marula fruit changes during ripening

Fruit ripening is a highly coordinated, genetically programmed, and an irreversible phenomenon involving a series of physiological, biochemical, and organoleptic changes that lead to the development of a soft and edible ripe fruit with desirable quality attributes. A wide spectrum of biochemical changes such as increased respiration, chlorophyll degradation, biosynthesis of carotenoids, anthocyanins, essential oils, and flavor and aroma volatiles, increased activity of cell wall hydrolases, and a transient increase in ethylene production are some of the major changes involved during fruit ripening [24, 25]. The results of the current study showed that as storage temperature increased from 25°C, SSC content, carotenoid and anthocyanin biosynthesis and chlorophyll degradation increased, but titratable acidity decreased, irrespective of storage temperature, indicating fruit ripening occurred in all the storage temperatures. The fruit colour change during ripening is due to unmasking of preformed pigments by degradation of chlorophyll and/or biosynthesis of carotenoids and anthocyanins and their accumulation in vacuoles [26, 27]. The increase in SSC and decrease in titratable acidity after three weeks of marula fruit storage at different temperatures is attributed to increased glucogenesis and hydrolysis of polysaccharides especially starch [26,27]. Decreased acidity and accumulation of sugars (SSC) and organic acids in marula fruits results in an excellent sugar/acid blend for edibility as fresh fruit or for processing of marula into various products. In general, total acidity decreases during ripening, though the content of one or more acids may increase. Total acidity decreases with ripening due to their utilization as respiratory substrates especially in the Krebs (TCA) cycle [16].

4.4 Vitamin C content

Loss of vitamin content especially vitamin C, pro-vitamin A, thiamine (vitamin B1) and nicotinic acid is common in horticultural produce poorly handled. Vitamin C includes both ascorbic acid ($C_6H_8O_6$) and its oxidation product dehydroascorbic acid ($C_6H_6O_6$) both being antiscorbutic. The two forms of vitamin C are largely interchangeable via the unstable monodehydroascorbic acid being catalyzed by any of the oxidases such as ascorbic acid oxidase, cytochrome oxidase, o-diphenol oxidase, p-diphenol oxidase and peroxidase [16, 28]. The balanced oxidation and reduction of ascorbic and dehydroascorbic acids leads to no loss of vitamin C. Loss can occur by irreversible conversion of dehydroascorbic acid to 2,3-dioxo-L-gulonic acid, which is then further metabolized. The reaction is pH dependent, being slow in acid pH, rapid at neutral pH and extremely rapid at alkaline pH [28]. The results of the current study showed that the loss of vitamin C was significantly reduced when marula fruit was stored at lower temperatures $\leq 12^\circ C$. Storage of fruits and vegetables at temperatures below $5^\circ C$ is reported to decrease the loss of vitamin C [28, 29]. The rate of loss of vitamin C is higher with higher storage temperature, an effect associated with loss of acidity [28]. Marula fruit according to the current study have very high levels of vitamin C in the range of 586-793 mg/100 g. Mangoes stored at $24-26^\circ C$ for 6 days has been reported to result in a drop in vitamin C content from the initial 71 mg/100g to 63.9 mg/100g [28].

5.0 CONCLUSION

Postharvest management of marula fruits is important for their successful marketing. The study showed that the most critical factor affecting the postharvest shelf life of marula fruits is storage temperature. It was concluded that in order to extend the shelf life and marketing period of marula fruits, the fruits should be stored at $12^\circ C$ and 90-95% RH, because the fruit will not suffer from chilling injury and will undergo normal ripening process. Marula fruit could initially be stored at $8^\circ C$ for 9 days and transferred to $12^\circ C$ to extend the marketing period and availability of the fruit beyond the harvest period.

REFERENCES

1. Kader, AA. Increasing food availability by reducing postharvest losses of fresh produce. *Acta Horticulturae*. 2005; 682: 2169 – 2175.

- 283 2. Emongor, VE. The effects of temperature on storage life of mango (*Mangifera indica*
284 L.). American Journal of Experimental Agriculture. 2015; 5(3): 252-261.
- 285 3. Kader, AA. Postharvest technology of horticultural crops - An overview from farm to
286 fork. 2013; 1: 1-8.
- 287 4. Will, RBH, McGlasson, B, Graham, D, Joyce, D. Postharvest: an introduction to the
288 physiology and handling of fruits, vegetables and Ornamentals. 4th revised edition,
289 Sydney, Australia: UNSW Press; Wallingford, Oxon, UK; New York: CAB
290 International. 1998; 262.
- 291 5. Gross, KC, Wang, CY and Sailtveit, M. The commercial storage of fruits, vegetables
292 and florist and nursery crops. <http://www.ba.ars.usda.gov/hb66/index.html>. 2002.
- 293 6. Hatton, TT. Reduction of chilling injury with temperature manipulation. In: Wang,
294 CY (Ed.). Chilling injury of horticultural crops. CRC Press, Boca Raton, FL. 1990;
295 269-280.
- 296 7. Dube, S, Dlamini, NR, Shereni, I, Sibanda, T. Extending the shelf Life of fresh
297 marula (*Sclerocarya berria*) juice by altering its physico-chemical parameters. In:
298 Jose CJ (Ed.). Biochemical Testing. 2012; 181-192.
- 299 8. AOAC. Association of Official Analytical Chemists Official Methods of Analysis.
300 Arlington, Virginia, USA. 1996; 17.
- 301 9. Creasy, LL. The effect of temperature on anthocyanin synthesis of New York State.
302 Horticultural Society. 1966; 11:93-96.
- 303 10. Zapsalis C, Francis FJ. Cranberry anthocyanins. Journal of Food Science. 1965; 30:
304 396-399.
- 305 11. Douglas JB. An evaluation of harvest indices for McIntosh apples in two orchards.
306 HortScience. 1983; 18(3): 216-218.
- 307 12. McGlasson, WB, Adato, I. Relationship between the capacity to Ripen and
308 Ontogeny in tomato fruits. Australian Journal of Plant Physiology. 1977; 4:451-458.
- 309 13. Harker, FR, Redgwell, RJ, Hallet, IC. (1987). Texture of fresh fruits.
310 Fruit. Horticultural Reviews. 1987; 20:12-224.

- 311 14. Emongor, VE. Thinning activity of benzyladenine on “Empire” apples: Application,
312 timing, and storage. PhD, Faculty of Agriculture, University of Guelph, Guelph
313 Ontario, Canada. 1995; 174.
- 314 15. Seymour, GB, Ostergard L, Chapman, NH. Fruit development and ripening. Annual
315 Review of Plant Biology. 2013; 64-219-241.
- 316 16. Emongor VE. Postharvest Physiology and Technology Manual. 2010; 256.
- 317 17. Tasneem A. Postharvest treatments to reduce chilling injury symptoms in stored
318 mangoes. Master of Science Thesis, Department of Bioresource Engineering,
319 Macdonald Campus, McGill University. 2004; 105.
- 320 18. Lim CS, Kang SM, Cho JL, Gross KC. Antioxidizing enzyme activities in chilling
321 sensitive and chilling tolerant pepper fruit as affected by stage of ripeness and storage
322 temperature. Journal of the American Society for Horticultural Science. 2009; 134:
323 156-163.
- 324 19. Trejo-Márquez MA, Ramírez-Viullatoro G, Camacho de la Rosa NA. Polyphenol
325 oxidase and peroxidase activities in mangoes stored at chilling temperature. Acta
326 Horticulturae. 2010; 864:53-55.
- 327 20. Wang CY. Physiological and biochemical responses of plants to chilling stress.
328 HortScience. 1982; 17: 173-187.
- 329 21. Saltveit, ME, Morris LL. Overview on chilling injury of horticultural crops. In: Wang
330 CY(Ed.). Chilling Injury of Horticultural crops. CRC Press, Boca Raton, Florida.
331 1990; 3-15.
- 332 22. Ding, CK, Wang, CY, Gross, KC, Smith, DL. Reduction of chilling injury and
333 transcript accumulation of heat shock proteins in tomato fruit by methyl jasmonate
334 and methyl salicylate, Plant Science. 2001; 161: 1153-1159.
- 335 23. Moatshe, OG, Emongor, VE, Oagile, O. Chemical thinning of morula (*Sclerocarya*
336 *birrea* subspecies *caffra*) fruit using benzyladenine. Botswana Journal of Agriculture
337 and Applied Sciences. 2011; 7(2): 45-56.
- 338 24. Tucker GA, Grierson D. Fruit ripening. In: Davis D (Ed.). The Biochemistry of
339 Plants. Academic Press Inc., New York. 1987; 12: 265-319.
- 340 25. Prasanna V, Prabha TN, Tharanathan RN. Fruit ripening phenomena-An Overview.
341 Critical Reviews in Food Science and Nutrition. 2007; 47: 1-19.

- 342 26. Srinivasa PC, Baskaran R, Ramesh MN, Prashanth KVH, Tharanathan RN. Storage studies of
343 mango packed using biodegradable chitosan film. *European Food Research and Technology*.
344 2002; 215: 504-508.
- 345 27. Kudachikar VB, Kulkarani SG, Prakash MNK, Vasantha MS, Prasad BA, Ramana KVR.
346 Physico-chemical changes during maturity of mango (*Mangifera indica* L.) variety Neelum.
347 *Journal of Food Science and Technology*. 2001; 38: 540-542.
- 348 28. Kays, JS, Paull, RE. *Postharvest Biology*. Exon press. 2004; 568.
- 349 29. Hardenburg RE, Watada AE, Wang CY. *The Commercial Storage of Fruits,*
350 *Vegetables, and Florist and Nursery Stocks*. United States Department of Agriculture.
351 *Agriculture Handbook*. 1986; 66: 1-22.

REVIEWED

By atanda saburi at 3:06 pm, Dec 19, 2015

atandasa