CORDIAL AND MEAN LABELINGS
ON EXTENDED DUPLICATE GRAPH OF COMB GRAPH.

K. Sutha, K. Thirusangu, S. Bala

Department of Mathematics

S. I. V. E. T. College, Gowrivakkam, Chennai – 600 073.

E-mail: sutharavindran@yahoo.com; kthirusangu@gmail.com; balasankarasubbu@yahoo.com

Abstract

A graph labeling is a mapping that carries a set of graph elements onto a set of numbers called labels(usually the set of integers). In this paper we prove the existence of graph labeling such as cordial, total cordial, product cordial, total product cordial, primecordial,odd mean labeling and even mean labeling for extended duplicate graph of Comb graph by presenting algorithms .

AMS Subject Classification : 05C78

Keywords: Graph labeling, Comb , Duplicate graph.

1 . Introduction

Graph labeling has the origin from a seminal paper by Rosa in 1967 [7] . It is defined as an assignment of integers to the vertices or edges or both subject to certain conditions.The notion of cordial labeling was introduced by Cahit[5]. A function f : V $\rightarrow$ {0,1} is said to be cordial labeling if each edge *uv* receives the label |f(*u*) – f(*v*)| such that the number of vertices labeled with '0' and the number of vertices labeled with '1' differ by at most one, and the number of edges labeled with '0' and the number of edges labeled with '1' differ by at most one. Moreover, it admits total cordial labeling if the number of vertices and edges labeled with '0' and the number of vertices and edges labeled with '1'differ by at most one.

Andar et. al., [1] proved that a cordial labeling of G can be extended to a cordial labeling of the graph obtained from G by attaching 2m pendent edges at each vertex of G.

The concept of product, total product and prime cordial labeling was introduced by

Sundaram et. al,[8]. A function $f : V \rightarrow \{0,1\}$ such that each edge *uv* receives the label $f(u) \times f(v)$ is said to be product cordial labeling if the number of vertices labeled with '0' and the number of vertices labeled with '1' differ by at most one, and the number of edges labeled with '0' and the number of edges labeled with '1'differ by at most one. Moreover, it is said to be total product cordial labeling if the number of vertices and edges labeled with '0' and the number of vertices and edges labeled with '1' differ by at most one.

A prime cordial labeling of a graph G with vertex set V is a function $f : V \rightarrow \{1, 2, 3, ..., p\}$such that the induced function $f^* : E \rightarrow N$ is defined by $f^*(v_iv_j)$ is assigned the label 1, if $\gcd(f(v_i), f(v_j)) = 1$ and 0, if $\gcd(f(v_i), f(v_j)) > 1$, then the number of edges labeled with '0' and '1' differ by atmost 1. It was proved that the graphs C_n if and only if $n \geq 6$: P_n if and only if $n \neq 3$ or 5; bistars; dragons; crowns ; ladders are prime cordial.

Nirmala investigated the concept of odd and even mean labeling.A function *f* is called an odd mean labeling of a graph G if there exists an injective function $f : V \rightarrow \{1,3,5,\ldots\ldots,2q-1\}$ such that when each edge v_iv_j receives a label $(f(v_i)+f(v_j))/2$, then the resulting edge labels are distinct. A graph which admits an odd mean labeling is called an odd mean graph. A function f is called an even mean labeling of a graph G if there exists an injective function $f : V \rightarrow \{2,4,\ldots\ldots\ldots,2q\}$ such that when each edge v_iv_j receives a label $(f(v_i) + f(v_j))/2$ then the resulting edge labels are distinct. A graph which admits an even mean labeling is called an even mean graph.

Thirusangu et.al,[10] have introduced the concept of extended duplicate graph. A Duplicate graph of a path graph G(V,E) is $DG=(V_1,E_1)$ where the vertex set $V_1 = V \cup V'$ and $V \cap V' = \Phi$ and $f: V \rightarrow V'$ is bijective (for $v \in V$, we write $f(v) = v'$ for convenience) and the edge set E_1 of DG is defined as follows: The edge v_iv_j is in E if and only if both v_iv_j' and $v_i'v_j$ are edges in E_1. Clearly duplicate graph is disconnected. In order to make it as connected, add an edge v_iv_i' for any `i'. This graph is called the Extended Duplicate Graph of a path graph G.

It was shown that the extended duplicate graph of a path P_m admits cordial ,total cordial, product cordial and total product cordial labeling.

Singh et.al, [9] proved some results on Comb graph. Let P_{m+1} be a path graph.Comb graph is defined as $P_{m+1} \Theta (m+1)K_1$. It has 2m+2 vertices and 2m+1 edges.

In this paper, we prove that, the extended duplicate graph of Comb graph is cordial, total cordial, product cordial, total product cordial, prime cordial, odd mean and even mean labeling by presenting algorithms.

2. Main Result

In this section we present the structure of the extended duplicate graph of a comb graph and establish the existence of cordial, total cordial, product cordial, total product cordial, prime cordial, odd and even mean labeling for the extended duplicate graph of Comb graph by presenting algorithms.

Definition: (Structure of the extended duplicate graph of a comb graph)

Let G(V,E) be a comb graph. DG(comb)= (V_1,E_1) is the duplicate graph of comb graph with 4m+4 vertices and 4m+2 edges.

Denote the vertex set as $V_1= \{v_1,v_2,\ldots\ldots\ldots v_{2m+2},v_1',v_2'\ldots\ldots\ldots v_{2m+2}'\}$ and the edge set as $E_1= \{v_iv_{i+1}' \cup v_i'v_{i+1}$ for $1 \le i \le m) \cup (v_i'v_{m+i+1} \cup v_iv_{m+i+1}'$ for $1 \le i \le m+1)\}$. Clearly DG(comb) is disconnected. The extended duplicate graph of a comb EDG(Comb) is obtained from DG(comb) by adding the edges (i) v_1v_{m+1} and $v_{m+2}'v_{2m+2}'$ if $m \equiv 1 \pmod 2$ (ii) v_1v_{m+1}' and $v_{m+2}'v_{2m+2}$ if $m \equiv 0 \pmod 2$.

Thus the extended duplicate graph of comb graph has 4m+4 vertices and 4m+4 edges.

Algorithm2.1

Procedure (cordial labeling for extended duplicate graph of comb graph)

$V = \{ v_1 , v_2 \ldots\ldots v_{2m+2} , v_1', v_2'\ldots\ldots v_{2m+2}'\}$

// assignment of labels to vertices

If m = 2n

for i = 1 to m+1 do

$\{ \quad v_i , v_{m+i+1}' \leftarrow 0$

$\quad v_i' , v_{m+i+1} \leftarrow 1$

$\}$

end for

If m = 2n+1

for i = 1 to 2m+2 do

$$\{ v_i , v_i' \leftarrow \begin{cases} 0 & \text{if } i \equiv 0 \pmod 2 \\ 1 & \text{otherwise} \end{cases}$$

}

end for

end procedure

output : vertex labeled extended duplicate graph of comb graph.

Theorem 2.1

The extended duplicate graph of comb graph is cordial.

Proof :

From the construction of extended duplicate graph of comb graph , it is clear that EDG(comb) (V,E) graph has 4m+4 vertices and 4m+4 edges.

In order to label the vertices, define a function $f : V \rightarrow \{0, 1\}$ as given in algorithm 2.1. Clearly,

The number of vertices labeled with '0' = m+1 +m+1 =2m+2

The number of vertices labeled with '1' = m+1 +m+1 =2m+2

Thus the number of vertices labeled with '0' and '1' is differ by atmost '1'.

In order to get the edge labels, define the induced map $f^* : E \rightarrow N$ such that $f^*(v_iv_j) = [f(v_i) + f(v_j)](\text{mod } 2)$.The edge labels are obtained as follows:

For $1 \leq i \leq m$

$f^*(v_iv_{i+1}') = f^*(v_i'v_{i+1}) = 1$, $f^*(v_iv_{m+i+1}') = f^*(v_i'v_{m+i+1}) = 0$

If m = 2n

$f^*(v_1v_{m+1}') = f^*(v_{m+2}'v_{2m+2}) = 1$, $f^*(v_1v_{m+2}') = f^*(v_{m+1}'v_{2m+2}) = 0$

If m = 2n + 1

$f^*(v_1v_{m+1}) = f^*(v_{m+2}'v_{2m+2}') = 1$, $f^*(v_{m+1}v_{2m+2}') = f^*(v_{m+1}'v_{2m+2}) = 0$

Thus the number of edges labeled with '0' = m+m+2 = 2m+2

The number of edges labeled with '1' = m+m+2 = 2m+2

Hence the number of edges labeled with '0' and '1' differ by atmost one.

Therefore EDG (comb) graph is cordial.

Theorem 2.2

The extended duplicate graph of comb graph EDG (comb), $m \geq 2$ is total cordial.

Proof :

By theorem 2.1, the number of vertices labeled with '1' and '0' are same as 2m + 2. The number of edges labeled with '1' and '0' are same as 2m + 2. Thus the total number of vertices

and edges labeled with ‘1’ is 4m+4 and the total number of vertices and edges labeled with ‘0’ is 4m+4. Clearly the total number of vertices and edges labeled with ‘1’ and the total number of vertices and edges labeled with ‘0’ differ by atmost one.

Hence the extended duplicate graph of comb graph is total cordial.

Example 2.1

Cordial labeling of EDG (comb)for m=5 and m=6 are given in figure 1 and figure 2 respectively.



Figure:1 cordial labeling of EDG(comb) graph for m = 5



Figure : 2 cordial labeling of EDG(comb) graph for m = 6

Algorithm 2.2

Procedure :(product cordial labeling for extended duplicate graph of comb graph)

$V = \{ v_1 , v_2 . . . . . v_{2m+2} , v_1', v_2'. . . . . v_{2m+2}'\}$

// assignment of labels to vertices

v_1' , v_2 ← 0 ; v_1 , v_{m+2}' ← 1

for i = 3 to m+1 do

{ v_i ← 1

}

end for

for i = 2 to m+1 do

{ v_i' ← 1

}

end for

for i = 1 to m do

{ v_{m+i+1} ← 0

}

end for

for i = 2 to m do

{ v_{m+i+1}' ← 0

}

end for

if m = 2n

v_{2m+2} ← 1 : v_{2m+2}' ← 0

if m = 2n+1

v_{2m+2} ← 0 : v_{2m+2}' ← 1

end for

end procedure

output :vertex labeled extended duplicate graph of comb graph.

Theorem 2.3

The extended duplicate graph of comb graph EDG (comb), $m \geq 2$ is product cordial.

Proof :

Clearly, EDG (comb) (V,E) has 4m + 4 vertices and 4m + 4 edges. In order to label the vertices, define a function $f : V \rightarrow \{0 , 1\}$, using algorithm 2.2.

It is evident that the number of vertices labeled with '0' = 2+m-1+m+1 = 2m+2.

Number of vertices labeled with '1' = 2+m-1+m+1 = 2m+2.

Hence the number of vertices labeled with '0' and '1' differ by atmost one. In order to get the edge labels, define the induced map $f^* : E \rightarrow N$ such that $f^*[v_iv_j] = [f(v_i) \times f(v_j)]$.

Thus the edge labels are obtained as follows :

$f^*(v_1v_2') = f^*(v_1v_{m+2}') = 1$: $f^*(v_1'v_2) = f^*(v_2v_3') = 0$

For $1 \leq i \leq m$

$f^*(v_i'v_{m+i+1}) = 0$

For $2 \leq i \leq m$

$f^*(v_i'v_{i+1}) = 1$: $f^*(v_iv_{m+i+1}') = 0$

For $3 \leq i \leq m$; $f^*(v_iv_{i+1}') = 1$

If $m = 2n$

$f^*(v_{m+1}v_{2m+2}') = 0$, $f^*(v_1v_{m+1}') = f^*(v_{m+1}'v_{2m+2}) = f^*(v_{m+2}'v_{2m+2}) = 1$

If $m = 2n + 1$

$f^*(v_{m+1}'v_{2m+2}) = 0$, $f^*(v_1v_{m+1}) = f^*(v_{m+1}v_{2m+2}') = f^*(v_{m+2}'v_{2m+2}') = 1$

Thus the number of edges labeled with '0' = 2+m+m-1+1 = 2m+2.

Number of edges labeled with '1' = 2+m-1+m-2 + 3 = 2m+2.

Hence the number of edges labeled with '0' and '1' differ by atmost one.

Therefore EDG (comb **)** graph is product cordial.

Theorem 2.4 :

The extended duplicate graph of comb graph , EDG (comb) $m \geq 2$ is total product cordial.

Proof :

By theorem 2.3, the number of vertices labeled with '0' and '1' are same as 2m+ 2 . The number of edges labeled with '0' and '1' are same as 2m+2. Thus the total number of vertices and edges labeled with '0' is 4m + 4 and the total number of vertices and edges labeled

with ‘1’ is 4m + 4. Hence the total number of vertices and edges labeled with ‘0’ and ‘1’ differ by atmost one.

Therefore EDG (comb) graph is total product cordial.

Example 2.2

Product cordial labeling of EDG(comb) graph for m =5 and m=6 are given in figure 3 and figure 4 respectively .



Figure:3 product cordial labeling of EDG(comb) graph for m = 5



Figure:4 product cordial labeling of EDG(comb) graph for m = 6

Algorithm 2.3

Procedure :(prime cordial labeling for extended duplicate graph of comb graph)

$V = \{ v_1 , v_2 \ldots\ldots v_{2m+2} , v_1', v_2' \ldots\ldots v_{2m+2}'\}$

// assignment of labels to vertices

$v_1 \leftarrow 4m-2$; $v_{m+2}' \leftarrow 4m+2$

for i = 2 to m-1 do

$$\{ v_i \leftarrow \begin{cases} 2i & \text{if } i \equiv 0 (\text{mod } 2) \\ 2m+2i-2 & \text{otherwise} \end{cases}$$

}

end for

for i = 1 to m-1 do

$$\{ v_i' \leftarrow \begin{cases} 2i & \text{if } i \equiv 1 (\text{mod } 2) \\ 2m+2i-2 & \text{otherwise} \end{cases}$$

}

end for

for i = 2 to m do

$$\{ v_{m+i+1} \leftarrow \begin{cases} 2i-1 & \text{if } i \equiv 1 (\text{mod } 2) \\ 2m+2i-3 & \text{otherwise} \end{cases}$$

}

end for

for i = 2 to m do

$$\{ v_{m+i+1}' \leftarrow \begin{cases} 2i-1 & \text{if } i \equiv 0 (\text{mod } 2) \\ 2m+2i-3 & \text{otherwise} \end{cases}$$

}

end for

if m = 2n

$v_m \leftarrow 2m$, $v_m' \leftarrow 1$, $v_{m+2} \leftarrow 4m+3$, $v_{m+1} \leftarrow 4m+1$

$v_{m+1}' \leftarrow 4m$, $v_{2m+2} \leftarrow 4m+4$, $v_{2m+2}' \leftarrow 4m-1$

if m = 2n+1

$v_m \leftarrow 4m+3$, $v_m' \leftarrow 2m$, $v_{m+1} \leftarrow 4m$ $v_{m+1}' \leftarrow 4m+1$,

$v_{2m+2}' \leftarrow 4m+4$, $v_{2m+2} \leftarrow 4m-1$, $v_{m+2} \leftarrow 1$.

end procedure

output: vertex labeled extended duplicate graph of comb graph.

Theorem 2.5

The extended duplicate graph of comb graph, $m \geq 2$ admits prime Cordial labeling.

Proof :

The EDG (comb)(V,E) graph has 4m+4 vertices and 4m+4 edges.Label the vertices of EDG(comb) graph by defining a function $f : V \rightarrow \{1, 2, 3, 4, \ldots, 4m + 4\}$, as given in algorithm 2.3. Thus all the 4m + 4 vertices are labeled.

Define the induced map $f^* : E \rightarrow N$ such that

$$f^*(v_iv_j) = \begin{cases} 1 & \text{if g.c.d.}\left(f(v_i), f(v_j)\right) = 1 \\ 0 & \text{if g.c.d.}\left(f(v_i), f(v_j)\right) > 1 \end{cases}$$

By the above induced function, the edge labels are obtained as follows :

For $1 \leq i \leq m-2$

$f^*(v_iv_{i+1}') = f^*(v_i'v_{i+1}) = 0$

For $1 \leq i \leq m$; $f^*(v_i'v_{m+i+1}) = 1$

For $2 \leq i \leq m$; $f^*(v_iv_{m+i+1}') = 1$

If m = 2n

$f^*(v_1v_{m+1}') = f^*(v_1v_{m+2}') = f^*(v_{m+1}'v_{2m+2}) = f^*(v_{m+2}'v_{2m+2}) = f^*(v_{m1}'v_m)=f^*(v_mv_{m+1}')=0$

$f^*(v_{m-1}v_m') = f^*(v_m'v_{m+1}) = f^*(v_{m+1}v_{2m+2}') = 1$

If m = 2n + 1

$f^*(v_1v_{m+1})=f^*(v_1v_{m+2}') = f^*(v_{m+1}v_{2m+2}') = f^*(v_{m+2}'v_{2m+2}') = f^*(v_{m-1}v_m') = f^*(v_m'v_{m+1})=0$

$f^*(v_{m-1}'v_m) = f^*(v_mv_{m+1}') = f^*(v_{m+1}'v_{2m+2}) = 1$

Thus in both the above cases,

The number of edges labeled with '1' =m+m-1+3 =2m+2.

Number of edges labeled with '0' = m-2+m-2+6=2m+2.

Hence the number of edges labeled with '1' and '0' differ by atmost '1'

Therefore EDG (comb) graph is prime cordial .

Example 2.3

Prime cordial labeling of EDG(comb) graphs for m=5 and m=6 are given in figure 5 and figure 6 respectively.



Figure: 5 prime cordial labeling of EDG(comb) graph for m = 5



Figure: 6 prime cordial labeling of EDG(comb) graph for m = 6

Algorithm 2.4

Procedure :(odd mean labeling for extended duplicate graph of comb graph)

V = { v_1 , v_2 . . . . . v_{2m+2} , v_1', v_2'. . . . . v_{2m+2}'}

// assignment of labels to vertices

for i = 1 to m+1 do

$$\{ \quad v_i \leftarrow \begin{cases} 2i-1 & \text{if } i \equiv 0 \pmod 2 \\ 2m+2i+1 & \text{otherwise} \end{cases}$$

$$v_i' \leftarrow \begin{cases} 2i-1 & \text{if } i \equiv 1 \pmod 2 \\ 2m+2i+1 & \text{otherwise} \end{cases}$$

}

end for

for i = 2 to m+2 do

$$\{ \quad v_{m+i} \leftarrow \begin{cases} 4m+2i+1 & \text{if } i \equiv 0 \pmod 2 \\ 6m+2i+3 & \text{otherwise} \end{cases}$$

$$v_{m+i}' \leftarrow \begin{cases} 4m+2i+1 & \text{if } i \equiv 1 \pmod 2 \\ 6m+2i+3 & \text{otherwise} \end{cases}$$

}

end for

end procedure

output :vertex labeled extended duplicate graph of comb graph.

Theorem 2.6

The extended duplicate graph of EDG (comb) graph admits odd mean labeling.

Proof:

Clearly EDG (comb)(V,E) graph has p = 4m+4 vertices and q = 4m+4 edges.

To label the vertices , define a function $f : V \rightarrow \{1,3,5,\ldots\ldots\ldots 2q-1\}$ as given in algorithm 2.4.

Thus all the 4m+4 vertices are labeled.

To get the edge labels, define the induced map

$f^* : E \rightarrow \{1,2,3\ldots\ldots,2q-1\}$ such that$f^*(v_iv_j) = f(v_i)+f(v_j)/2$ for all $v_iv_j \in E$

By the induced function , the edge labels are obtained as follows:

For $1 \leq i \leq m$

$$f^*(v_iv_{i+1}') = \begin{cases} 2i & \text{if } i \equiv 0 \pmod 2 \\ 2m+2i+2 & \text{otherwise} \end{cases}$$

$$f^*(v_i'v_{i+1}) = \begin{cases} 2i & \text{if } i \equiv 1 \pmod 2 \\ 2m+2i+2 & \text{otherwise} \end{cases}$$

For $1 \leq i \leq m+1$

$$f^*(v_i v'_{m+i+1}) = \begin{cases} 2m+2i+1 & \text{if } i \equiv 0 \pmod 2 \\ 4m+2i+3 & \text{otherwise} \end{cases}$$

$$f^*(v'_i v_{m+i+1}) = \begin{cases} 2m+2i+1 & \text{if } i \equiv 1 \pmod 2 \\ 4m+2i+3 & \text{otherwise} \end{cases}$$

If m = 2n

$f^*(v_1 v'_{m+1}) = 2m+2$, $f^*(v'_{m+2} v_{2m+2}) = 6m+6$.

If m = 2n +1

$f^*(v_1 v_{m+1}) = 2m+2$, $f^*(v'_{m+2} v'_{2m+2}) = 6m+6$.

Thus all the labeled edges are distinct .

Hence EDG (comb) graph admits odd mean labeling .

Example 2.4

Odd mean labeling of EDG(comb) graphs for m=5 and m=6 are given in figure 7 and figure 8 respectively.



Figure : 7 odd mean labeling of EDG(comb) graph for m = 5



Figure : 8 odd mean labeling of EDG(comb) graph for m = 6

Algorithm 2.5

Procedure :(even mean labeling for extended duplicate graph of comb graph)

$V = \{ v_1 , v_2 \ldots\ldots v_{2m+2} , v_1', v_2' \ldots\ldots v_{2m+2}' \}$

// assignment of labels to vertices

for i = 1 to m+1 do

$$\{ \quad v_i \leftarrow \begin{cases} 2i & \text{if } i \equiv 0 (\text{mod } 2) \\ 2m+2i+2 & \text{otherwise} \end{cases}$$

$$v_i' \leftarrow \begin{cases} 2i & \text{if } i \equiv 1 (\text{mod } 2) \\ 2m+2i+2 & \text{otherwise} \end{cases}$$

}

end for

for i = 2 to m+2 do

$$\{ \quad v_{m+I} \leftarrow \begin{cases} 4m+2i+2 & \text{if } i \equiv 0 (\text{mod } 2) \\ 6m+2i+4 & \text{otherwise} \end{cases}$$

$$v_{m+i}' \leftarrow \begin{cases} 4m+2i+2 & \text{if } i \equiv 1 (\text{mod } 2) \\ 6m+2i+4 & \text{otherwise} \end{cases}$$

}

end for

end procedure

output :vertex labeled extended duplicate graph of comb graph.

Theorem 2.7

The extended duplicate graph of comb graph, EDG (comb)graph admits even mean labeling.

Proof:

Clearly EDG (comb)(V,E) graph has p = 4m+4 vertices and q = 6m+4 edges.

To label the vertices , define a function

$f : V \rightarrow \{2,4,6,\ldots\ldots\ldots, 2q\}$ as given in algorithm 2.5.

Thus 4m+4 vertices are labeled.

To get the edge labels, define, the induced map $f^* : E \rightarrow \{1,2,3\ldots\ldots\ldots,2q\}$ such that

$f^*(v_iv_j) = [f(v_i)+f(v_j)]/2$ for all $v_iv_j \in E$

By the induced function , the edge labels are obtained as follows:

For $1 \leq i \leq m$

$$f^*(v_iv_{i+1}') = \begin{cases} 2i+1 & \text{if } i \equiv 0 \pmod 2 \\ 2m+2i+3 & \text{otherwise} \end{cases}$$

$$f^*(v_i'v_{i+1}) = \begin{cases} 2i+1 & \text{if } i \equiv 1 \pmod 2 \\ 2m+2i+3 & \text{otherwise} \end{cases}$$

For $1 \leq i \leq m+1$

$$f^*(v_iv_{m+i+1}') = \begin{cases} 2m+2i+2 & \text{if } i \equiv 0 \pmod 2 \\ 4m+2i+4 & \text{otherwise} \end{cases}$$

$$f^*(v_i'v_{m+i+1}) = \begin{cases} 2m+2i+2 & \text{if } i \equiv 1 \pmod 2 \\ 4m+2i+4 & \text{otherwise} \end{cases}$$

If m = 2n

$f^*(v_1v_{m+1}') = 2m+3$; $f^*(v_{m+2}'v_{2m+2}) = 6m+7$.

If m = 2n +1

$f^*(v_1v_{m+1}) = 2m+3$; $f^*(v_{m+2}'v_{2m+2}') = 6m+7$.

Thus all the labeled edges are distinct .

Hence EDG (comb) graph admits even mean labeling .

Example 2.5

Even mean labeling of EDG(comb) graph for m = 5 and m = 6 are given in figure 9 and figure 10 respectively.



Figure : 9 even mean labeling of EDG(comb) graph for m = 5



Figure : 10 even mean labeling of EDG(comb) graph for m = 6

Conclusion:

In this paper we have presented algorithms and proved that the extended duplicate graph of Comb graph admits cordial, total cordial, product cordial,total product cordial,prime cordial, odd mean labeling and even mean labeling.

References:

1. Andar.M, Boxwala.S and Limaye.N. *New families of cordial graphs,* J. Combin.Math.Combin.comput.,53(2005) 117-154.
2. Bala.E, Thirusangu. K. *Some graph labelings in extended triplicate graph of a path P_n.* International Review of Applied Engineering Research (IRAER).,1(2011)81-92.
3. Baskar Babujee. J and Shobana. L, *prime cordial labeling,*Int.Review on Pure and Appl.Math.,5 (2009) 277-282.
4. Bondy J.A and Murthy U.S.R. *Graph theory with applications* (1976).
5. Cahit.I.Cordial graphs , *Weaker version of graceful and harmonious graphs*, Ars Combin,Vol.23(1987) pp.201-207
6. Gallian J.A. *A Dynamic survey of graph labeling. The Electronic Journal of combinatories.* 17(2014) # DS6.
7. Rosa. A. On certain valuations of a graph, Theory of graphs, International symposium, Rome, Gordon and Breach N.Y. at Dunod Paris, (1967) 349-355.
8. Sunadaram.M ,Ponraj. R and Somasundaram. S . *Prime cordial labeling of graphs,* J. Indian Acad.Math.,Vol.27(2005) pp.373-390 .
9. Singh G.S and Varkey . T.K.M, Some results on prime labeling of graphs, preprint.
10. Thirusangu.K, Selvam. B and Ulaganathan. P.P, *Cordial labeling in extended duplicate twig graphs,* International Journal of Computer , Mathematical Sciences and Applications,Vol.4,Nos 3-4 (2010) pp.319-328.