

## Weed control in direct seeded rice using new herbicide combination Under Indian Tropical Condition

### ABSTRACT

Direct seeded rice (DSR) is gaining momentum in India due to high demand of labour during peak season of transplanting and short period availability of water. Weed management is a major factor contributing a considerable share to the cost of production and deciding the final yield, especially in DSR as the crop and weeds emerge simultaneously due to which the crop suffers competition even from early stage of growth which in turn reduces the grain yield. Weeds are the main biological constraints to its success. Field experiments were conducted in the *rabi* season of 2013 and 2014 to study the new formulation of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE against weeds in DSR and their residual effect on succeeding crop. Results revealed that the post-emergence (POE) application of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> gave significantly lower total weed density (25.78 plants m<sup>-2</sup> in 2013 and 24.19 plants m<sup>-2</sup> in 2014), total weed biomass (24.89 g m<sup>-2</sup> in 2013 and 34.56 g m<sup>-2</sup> in 2014) and higher weed control efficiency (80.07% in 2013 and 81.68% in 2014) at 40 days after herbicide spray (DAHS). Application of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> as POE herbicide can keep the weed density and dry weight below the economic threshold level and increase the grain yield of 5676 and 6388 kg ha<sup>-1</sup> in DSR. Non-treated control accounted for lower grain yield which in turn reflected through higher weed index of 51.83 and 52.85%, respectively during both the years due to heavy competition of weeds for nutrient, space and light. Succeeding crop of green gram was sown immediately after the harvest of DSR was not affected by the residue of new formulation of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at all different doses. The results of this study are important for farmers growing DSR in making decisions regarding the application of POE herbicide combination, according to existing weed flora in the field.

Keywords: DSR, Herbicide combination, Weed density, Weed biomass, Weed control efficiency, Grain yield, Succeeding green gram.

### I. Introduction

Rice (*Oryza sativa* L.) is the leading cereal of the world and more than half of the human race for their daily sustenance (Chauhan and Johnson, 2011). Globally, actual rice yield losses due to pests have been estimated at 40%, of which weeds has the highest loss potential of 32%. The worldwide estimated loss in rice yield from weeds is around 10% of the total production (Oerke and Dehne, 2004). Though India has the largest rice growing area with 44.8 million hectares it stands second after China with respect to production 106.0 mt in 2013-14. Currently, India produces rice that is sufficient not only to meet the domestic demand but was largest exporter during 2012. However, the rapidly increasing population projected to be 1.6 billion by 2050 calls for stepping up the current production of 106 mt of milled rice to 140 mt at enhanced productivity of 3.5 t ha<sup>-1</sup>. Transplanting is the traditional system of rice cultivation and it is in vogue in many rice growing areas. Such a rice production system, however, requires a large amount of water during puddling and transplanting (Chauhan 2012a, Chauhan et al., 2012b). In order to check the declining water table, a new technique of direct seeding is now fast replacing traditional transplanting rice area with good drainage and irrigation facilities (Balasubramanian and Hill 2000). Weeds are main biological constraints to the production of direct seeded rice (Chauhan, 2012b; Chauhan and Johnson, 2010).

Direct seeded rice cultivation is subjected to greater weed competition than transplanted rice because both weeds and crop seeds emerge at the same time and compete with each other for germination resulting in less grain yield. Crop competitiveness is the ability of the crop to produce desirable yields in the presence of weeds (Zhao, 2006). In tropic, average rice yield losses from weeds is 35% (Oerke and Dehne, 2004). Sunil et al. (2010) as stated, season-long weed competition in DSR may cause yield reduction upto 80%. In DSR, weeds emerge simultaneously rice seedlings at the early growth stages when rice is highly susceptible to the weed competition (Khaliq and Matloob, 2011; Chauhan, 2012). Thus, an efficient and timely weed control is crucial for the success of DSR. Direct seeding can curtail water and labor inputs involved in rice production; nevertheless, its large-scale adoption is impeded by heavy weed infestation. However, for cultivation of DSR, weeds are a major hurdle as nearly all *rabi* season weeds depending upon seed bank in the field infest this crop. DSR is possible provided there is a good crop establishment as well as adequate weed control methods is available to keep the crop free from weeds (Rao and Nagamani, 2007). Efficient, cost-effective and timely weed

management options remain pivotal to making DSR profitable and commercially acceptable. Such a strategy should help improve yield and reduce production costs as well as minimize the negative effects of weeds on the quality of the produce. Timely and effective weed control has a positive correlation with good crop stand and high grain yield of DSR. Manual weeding although is effective and the most common practice of weed control in direct seeded rice; these have several limitations particularly during peak period which makes it further problematic. In hand weeding, it is difficult to differentiate and remove the grassy weeds especially *Echinochloa crusgalli* and *Echinochloa colonum* due to the phenotypical similarities between weeds and rice seedlings in the early stages. Herbicides are considered to be an alternative supplement to hand weeding. The use of herbicides offers selective control of weeds right from beginning, giving crop an advantage of good start and competitive superiority over weeds. Hence, chemical weed control in direct seeded rice has gained importance because of the intensity of weed problem, coupled with the lack of labour for weeding and high cost. Chemical weed control has expanded manifold in DSR (Chauhan and Opena, 2013a, b) and is likely to increase further with the increased adoption of direct seeding.

In India, the high cost and scarcity of labour and cost effective as well as timely control of weeds have increased use of herbicides for weed control in almost all crops (Rao et al., 2014). In order to control weeds, farmers use both pre and post emergence herbicides (Mahajan and Timsina, 2011). Both pre and post emergence herbicides, if properly used, are quite effective in suppressing weeds in DSR (Chauhan, 2012). To the best of our understanding, a very few studies in this line have been conducted in DSR grown in Western Zone of Tamil Nadu, India. Moreover, the rice herbicides presently used in Tamil Nadu are mainly pre-emergence therefore; weeds coming at later stages of crop growth are not controlled as effectively as the weeds at emergence stage. This situation warranted for initiating research efforts to evaluate and identify suitable post-emergence herbicides. But sometimes continuous use of a single herbicide (pretilachlor) may lead the buildup of herbicide resistance in weeds. Nevertheless, indiscriminate use of herbicides is driving agro-ecosystem toward declining species diversity and in many situations, is leading to herbicide resistance (Powles and Yu, 2010). Singh (2008) found that the continuous changes in weed community composition in just five years. Without any doubt, the development and availability of effective POE herbicides have encouraged farmers to try this

new method of crop establishment (DSR) in Tamil Nadu. Currently available rice herbicide have a narrow spectrum of activity and their efficacy is further limited when they are used alone (Singh, 2008; Chauhan, 2012). This rarely provides season long weed control (Khaliq et al., 2011a, b). Control of complex weed flora with a single POE application is really a difficult task for the DSR farmers (Mahajan et al., 2013). Therefore, the combined application of different herbicides with different mode of action is required for broad spectrum weed control in DSR and for delaying the development of herbicide resistance.

Hence, there is a need to focus attention on new herbicide combination to enhance the weed control efficiency, broadening the spectrum of weed control and saving the herbicide and labour requirements. Literatures suggest that the repeated use of the same herbicides encourages the problem of herbicide resistance in weeds (Kim, 1996). For a broad spectrum of weed control in DSR, applications of herbicides with different mode of action (chemistry) are needed. With changing scenario of weed management, farmers need new herbicides or new herbicide combination having with high efficacy, low phytotoxicity, there was no residual effect on succeeding crops and cost effective. Thus, it is essential to identify economic and effective herbicide combinations for managing complex weed flora in DSR. This study was conducted for general detailed information for managing a mixed population of grasses, sedges and broad leaved weeds in DSR effectively and economically with herbicide combination of newly available POE herbicides. The present work is intended to look out the broad spectrum weed control through new POE herbicide combination in *rabi* season DSR of Tamil Nadu.

## **2. Materials and Methods**

### *2.1. Experimental Site and Initial Soil Characteristics*

A field study was conducted for two years (*rabi* seasons of 2013 and 2014) at the research farm of Wetland Farm (Field No: N<sub>1</sub>), Tamil Nadu Agricultural University, Coimbatore, India. The experimental farm was located in Western Zone of Tamil Nadu is at 11°29'N latitude and 77°08'E longitude with an altitude of 256 m above MSL. The climate is semi arid, with an average of 674.2 mm distributed over 47 rainy days (mean of past 50 years). The maximum rainfall received during the cropping period was 70 mm. During the cropping period, the maximum and minimum temperature ranged from 35.7 to 27.0°C and 26.0 to 19.8°C, respectively. Relative humidity ranged from 61 to 95 per cent and 29 to 75 per cent during

forenoon and afternoon, respectively. The solar radiation received during the cropping period ranged from 224 to 462.6 cal/cm<sup>2</sup>/day and the sunshine hours ranged from 1.4 to 9.0 hrs/day. The evaporation prevailing during the cropping period ranged from 2.4 to 9.2 mm. The soil was clay loam in texture with low in available nitrogen (238 kg ha<sup>-1</sup>), medium in available phosphorus (16.8 kg ha<sup>-1</sup>) and high in available potassium (518 kg ha<sup>-1</sup>) with 0.5% organic matter with a pH of 7.4.

## *2.2. Experimental design and treatments*

The treatments in each year were arranged in a randomized complete block design with three replication. Twelve weed control treatments were included with different herbicide combination options for weed control in DSR. Herbicides included in the study were bispyribac-sodium, metamifop, almix, clincher and wetter (isoxadifen, a safener).

## *2.3. Experimental details, selection of cultivar and sowing*

In each year, rice (cv. ADT 43, a cultivar with the duration of 120 days) was seeded in the first week of September and the harvested in last week of December. Manually operated rice drum seeder developed by Tamil Nadu Agricultural University, Coimbatore was used for sowing the seeds. The seeder has two wheels at both the ends. It drops the seeds at 30 cm apart in continuous row. At a time, eight rows of rice seeds were sown. A seed rate of 40 kg ha<sup>-1</sup> was adopted. Before sowing the field was drained to keep it under saturated condition to facilitate easy sowing and uniform establishment of seedlings. A thin film of water was maintained at the time of sowing. For the next 8-15 days, irrigation and drainage of water were alternated to facilitate aeration, adequate moisture for germination of seed and establishments of seedlings. Thereafter, the plots were irrigated to 2 cm depth uniformly in all the treatments after the appearance of hair line cracks, upto panicle initiation stage. After panicle initiation, the plots were irrigated to 5 cm depth on disappearance of ponded water. Irrigation was stopped 15 days prior to harvest.

## *2.4. Treatment details*

New formulation of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE was applied as POE herbicide on 10 to 15 DAS. Bispyribac-sodium (Nominee gold) has been widely used for DSR with its excellent foliar efficacy against grasses, sedges and broad leaved weeds. Metamifop which was discovered by Dongbu Honnong Co., Ltd. is a novel grass

herbicide with excellent foliar efficacy against grasses and crop safety. Hand operated knapsack sprayer fitted with a flat fan type nozzle (WFN 40) was used for spraying the herbicides adopting a spray volume of 500 litres ha<sup>-1</sup> in DSR. The herbicides were sprayed by keeping a thin film of water in the field. The field was neither drained nor irrigated for 2 days after application of herbicides. The non-treated control plot was kept undisturbed for the entire cropping period.

**Table 1. Herbicide treatments used in the study**

| Tr. No          | Treatment details                                   | Dose g.a.i/ha             | Dose ml/gm/ha of Formulation | Time of Application |
|-----------------|---|---------------------------|------------------------------|---------------------|
| T <sub>1</sub>  | Bispyribac sodium 4% SE + metamifop 10% SE + Wetter | 42 g a.i. + 100 ml wetter | 300 ml +100 ml wetter        | 10-15 DAS           |
| T <sub>2</sub>  | Bispyribac sodium 4% SE + Metamifop 10% SE + Wetter | 56 g a.i. + 100 ml wetter | 400 ml +100 ml wetter        | 10-15 DAS           |
| T <sub>3</sub>  | Bispyribac sodium 4% SE + Metamifop 10% SE + Wetter | 70 g a.i. + 100 ml wetter | 500 ml +100 ml wetter        | 10-15 DAS           |
| T <sub>4</sub>  | Almix (Chlorimuron + Metsufuron 20% WP)             | 4 g a.i.                  | 20 g                         | 10-15 DAS           |
| T <sub>5</sub>  | Clincher (Cyhalofop Buthyl 10% EC)                  | 80 g a.i.                 | 800 ml                       | 10-15 DAS           |
| T <sub>6</sub>  | Bispyribac sodium 10% SC + Wetter                   | 20 g a.i. + 100 ml wetter | 200 ml + 100 ml wetter       | 10-15 DAS           |
| T <sub>7</sub>  | Metamifop 10% SE + Wetter                           | 50 g a.i. + 100 ml wetter | 500 ml +100 ml wetter        | 10-15 DAS           |
| T <sub>8</sub>  | Bispyribac sodium 4% SE + Metamifop 10% SE          | 70 g a.i.                 | 500 ml                       | 10-15 DAS           |
| T <sub>9</sub>  | Bispyribac sodium 10% SC                            | 20 g a.i.                 | 200 ml                       | 10-15 DAS           |
| T <sub>10</sub> | Metamifop 10% SE                                    | 50 g a.i.                 | 500 ml                       | 10-15 DAS           |
| T <sub>11</sub> | Hand weeding twice on 25 and 45 DAS                 | ---                       | ---                          | ---                 |
| T <sub>12</sub> | Non-treated control                                 | ---                       | ---                          | ---                 |

Abbreviation: DAS - Days after sowing.

## 2.5. Observation on weeds

### 2.5.1. Weed flora of the experimental field

To account for the general weed flora of the experimental field, species wise weeds observed in the treatment plots were recorded during the period of maximum appearance of 20 and 40 days after herbicide spray (DAHS). The weed flora of the experimental site was recorded species wise.

### 2.5.2. Weed density

The weed count was recorded species wise using 0.5 m x 0.5 m quadrant from four randomly fixed places in each plot and the weeds falling within the frames of the quadrant were counted and the mean values were expressed in number m<sup>-2</sup>. The density of grasses, sedges and broad leaved weeds and also the total weeds were recorded at 20 and 40 days after herbicide application (DAHS) and expressed in number m<sup>-2</sup>.

### 2.5.3. Weed dry weight

The weeds falling within the frames of the quadrant were collected, categorized into grasses, sedges and broadleaved weeds, first shade dried and later dried in hot-air oven at 80°C for 72 hrs. The dry weight of grasses, sedges and broadleaved weeds were recorded separately at 20 and 40 DAHS and expressed in g m<sup>-2</sup>.

### 2.5.4. Weed control efficiency

Weed control efficiency (WCE) was calculated as per the procedure given by Mani *et al.* (1973).

$$\text{WCE \%} = \frac{\text{WD}_c - \text{WD}_t}{\text{WD}_c} \times 100$$

Where,

WCE - weed control efficiency (%)

WD<sub>c</sub> - weed biomass (g m<sup>-2</sup>) in control plot

WD<sub>t</sub> - weed biomass (g m<sup>-2</sup>) in treated plot

### 2.5.5. Weed index

Weed index (WI) was calculated as per the method suggested by Gill and Vijaya Kumar (1969).

$$\text{WI} = \frac{X - Y}{X} \times 100$$

Where, X = yield (kg ha<sup>-1</sup>) from minimum weed competition plot

$Y$  = yield ( $\text{kg ha}^{-1}$ ) from the treatment plot for which WI is to be worked out.

## 2.6. Observation on crop

### 2.6.1. Grain yield

Grains from each net plot were cleaned, sun dried, weighed and adjusted to 14% moisture content and the grain yield was expressed in  $\text{kg ha}^{-1}$ .

## 2.7. Residual crop cultivation

To study the residual effect of herbicides applied to direct seeded rice, the succeeding crop of green gram (cv. Co 6) was raised without disturbing the layout of the previous experiment. After the harvest of rice crop, the follow up crop was dibbled in rice stubbles. A seed rate of  $20 \text{ kg ha}^{-1}$  was adopted for the green gram crop with a spacing of 30 cm x 10 cm.

## 2.8. Statistical analysis

The data collected for direct seeded rice was statistically analyzed following the procedure given by Gomez and Gomez (2010) for randomized block design. The data pertaining to weeds and germination were transformed to square root scale of  $\sqrt{(X+2)}$  and analyzed as suggested by Snedecor and Cochran (1967). Whenever significant difference existed, critical difference was constructed at five per cent probability level. Such of those treatments where the differences are not significant were denoted as NS.

## 3. Results and discussion

### 3.1. General weed flora of the experimental field

A critical analysis of relative proportion of grasses, sedges and broad leaved weeds to total weed population in non-treated control revealed that during the crop growth period, the population of sedges was higher than that of grasses and broad leaved weeds. Among the grasses, *Echinochloa crus-galli* (L.) Beauv., *Echinochloa colona* (L.) Link., *Dinebra retroflexa* (Vahl.) Panzer. and *Panicum repens* (L.) were the dominant species and major sedges were *Cyperus difformis* (L.), *Cyperus irria* (L.) and *Fimbristylis miliacea* (L.) Vahl. Among the broad leaved weeds *Marsilea quadrifoliata* (Linn.), *Ammania baccifera* (L.) and *Eclipta alba* (L.)



Hassk. were the dominant species. However, a species-wise result was given for the first five weeds only, as they were the predominant weeds in the experimental trial.

### 3.2. Effect on weeds

#### 3.2.1. Weed density and weed biomass

##### 3.2.1.1. *Echinochloa crus-galli*

*E. crus-galli* density in the non-treated control (20.36 and 29.45 plants  $m^{-2}$  in 2013 and 34.54 and 56.89 plants  $m^{-2}$  in 2014, respectively) recorded higher population of *E. crus-galli* at 20 and 40 DAHS (Table 2). During *rabi*, 2013 the lower density of *E. crus-galli* was observed by POE application of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.  $ha^{-1}$  + wetter at 100 ml  $ha^{-1}$  (2.30 and 6.54 plants  $m^{-2}$ ) and it was similar to the density of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.  $ha^{-1}$  without wetter (2.86 and 7.86 plants  $m^{-2}$ ) and bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i.  $ha^{-1}$  + wetter at 100 ml  $ha^{-1}$  (3.86 and 7.55 plants  $m^{-2}$ ). In this study, POE application of herbicide alone like almix at 4 g a.i.  $ha^{-1}$  (5.63 and 11.19 plants  $m^{-2}$ ) and clincher at 80 g a.i.  $ha^{-1}$  (7.21 and 12.77 plants  $m^{-2}$ ) were recorded poor control weed of *E. crus-galli* than that of herbicide combination of bispyribac sodium 4% SE + metamifop 10% SE + wetter at 100 ml  $ha^{-1}$  with three different doses. During *rabi* 2014, at 20 and 40 DAHS, POE application bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.  $ha^{-1}$  + wetter at 100 ml  $ha^{-1}$  (7.52 and 10.24 plants  $m^{-2}$ ) registered significantly lesser weed density of *E. crusgalli* which was comparable to bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.  $ha^{-1}$  without wetter (7.82 and 13.26 plants  $m^{-2}$ ) and bispyribac sodium 14% SE + metamifop 10% SE at 56 g a.i.  $ha^{-1}$  + wetter at 100 ml  $ha^{-1}$  (10.76 and 16.78 plants  $m^{-2}$ ). Our results confirmed that the POE application of new herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.  $ha^{-1}$  + wetter at 100 ml  $ha^{-1}$  obtained higher weed control during both the years due to vigorous growth of crop that did not allow later flush of *E. crus-galli*. While, this treatment was quite effective against *E. crus-galli* and *E. colona* thus facilitate the DSR in attaining vigorous growth at the initial stage than that in turn provided smothering effect at later stage of the crop. Mahajan and Chauhan (2013) in an earlier study revealed that the single application of azimsulfuron and bispyribac sodium did not control *D. aegyptium*.

##### 3.2.1.2. *Dinebra retroflexa*

The density of *D. retroflexa* in the non-treated control (9.56 and 14.23 plants m<sup>-2</sup> in 2013 and 8.24 and 9.45 plants m<sup>-2</sup> in 2014, respectively) registered higher population of *D. retroflexa* at 20 and 40 DAHS (Table 2). All the tested herbicide treatments reduced the density of *D. retroflexa* as compared to the non-treated control. During both the years of study, the lower density of *D. retroflexa* was observed in POE application of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (0.00 and 0.00 plants m<sup>-2</sup> in 2013 and 0.00 and 0.84 plants m<sup>-2</sup> in 2014, respectively) and it was similar to the density of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter (0.00 and 0.82 plants m<sup>-2</sup> in 2013 and 0.00 and 1.22 plants m<sup>-2</sup> in 2014, respectively) and bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (0.00 and 1.56 plants m<sup>-2</sup> in 2013 and 0.00 and 0.84 plants m<sup>-2</sup> in 2014, respectively). POE application of bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (2.36 and 3.82 plants m<sup>-2</sup> in 2013 and 1.86 and 2.42 plants m<sup>-2</sup> in 2014, respectively) showed effective in reducing its density of *D. retroflexa* as compared with the individual application of clincher at 80 g a.i. ha<sup>-1</sup> (3.22 and 4.64 plants m<sup>-2</sup> in 2013 and 3.02 and 5.73 plants m<sup>-2</sup> in 2014, respectively). Our findings proved that almost all the tested herbicide combination with different doses were effective for the control of *D. retroflexa*; however, the combined application of 4% SE + metamifop 10% SE was better when herbicides were applied as alone like almix, clincher, bispyribac sodium and metamifop. This study was revealed that relatively POE application of clincher and almix against lesser effective of *D. retroflexa* control as compared to POE application of bispyribac sodium and metamifop alone.

### 3.2.1.3. *Panicum repens*

The density of *P. repens* in the non-treated control (7.42 and 11.46 plants m<sup>-2</sup> in 2013 and 6.42 and 8.42 plants m<sup>-2</sup> in 2014, respectively) registered higher population of *D. retroflexa* at 20 and 40 DAHS (Table 2). POE application of metamifop 10% EC at 50 g a.i. ha<sup>-1</sup> (3.22 and 5.02 plants m<sup>-2</sup> in 2013 and 2.44 and 4.21 plants m<sup>-2</sup> in 2014, respectively) registered higher weed density of *P. repens* as compared to individual application of bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> alone. During both the years, POE application of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 42, 56 and 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> at all different doses recorded lower density of *P. repens* and also significantly superior to other

herbicidal treatments. The rice crop followed the herbicide combination bispyribac sodium 4% SE + metamifop 10% SE was almost weed free and did not allow the later flush of weed seedlings to grow due to vigorous growth of the crop. In general, POE application of clincher at 80 g a.i.ha<sup>-1</sup> showed lesser density of *P. repens* as compared to almix at 4 g a.i.ha<sup>-1</sup> at both the stages of observation.

#### 3.2.1.4. *Cyperus difformis*

*C. difformis* was one of the dominant sedge present in the experimental field. Different weed control treatments imposed to direct seeded rice significantly influenced the density of *C. difformis* at all the stages. During *rabi* 2013, application of POE herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (5.32 and 9.56 plants m<sup>-2</sup>) proved to be effective in controlling the density of sedge and recorded significantly lower density of *C. difformis* at 20 and 40 DAHS (Table 3). However, POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter (5.38 and 11.01 plants m<sup>-2</sup>) was comparable with application of bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (7.56 and 13.19 plants m<sup>-2</sup>) obtained lesser density of *C. difformis* at 20 and 40 DAHS. Individual application of bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> (12.50 and 19.54 plants m<sup>-2</sup>) and metamifop 10% EC at 50 g a.i. ha<sup>-1</sup> (18.16 and 24.98 plants m<sup>-2</sup>) were ineffective against sedge weed control when compared to herbicide combination. However, the combined application of bispyribac sodium 4% SE + metamifop 10% SE + wetter with all different doses were effectively controlled the sedges present in the experimental plots. The results also suggested a poor control of *C. difformis* by individual application of almix at 4 g a.i. ha<sup>-1</sup> and clincher at 80 g a.i. ha<sup>-1</sup> as compared to other herbicidal combination. POE of almix at 4 g a.i. ha<sup>-1</sup> (11.68 and 22.89 plants m<sup>-2</sup>) registered lower density of *C. difformis* when compared to at 80 g a.i. ha<sup>-1</sup> (14.56 and 20.19 plants m<sup>-2</sup>) at 20 and 40 DAHS. Higher density of *C. difformis* was invariably observed in non-treated control (39.40 and 52.46 plants m<sup>-2</sup>) at 20 and 40 DAHS. POE application of bispyribac sodium is effective mainly against *C. rotundus* was given by (Mahajan and Chauhan, 2013). During *rabi*, 2014, at 20 and 40 DAHS, bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (2.56 and 4.16 plants m<sup>-2</sup>) recorded least population of *C. difformis* among all the treatments used at all stages of observation followed by the treatment with application of bispyribac sodium 4% SE +

metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter (2.58 and 4.57 plants m<sup>-2</sup>). Based on the two year experimentation, density of *C. difformis* was lower in second season trial when compared to first season at all the stages of observation. Kumaran et al. (2015) revealed that early POE application of bispyribac sodium 10% SC at 40 g a.i. ha<sup>-1</sup> was more effective against *C. rotundus* as compared to pretilachlor S at 0.45 kg g a.i. ha<sup>-1</sup> followed by one hand weeding on 40 days after sowing (DAS).

#### 3.2.1.5. *Marsilea quadrifoliata*

The density of *M. quadrifoliata* in the non-treated control plot was 17.52 and 32.45 plants m<sup>-2</sup> in 2013 and 13.67 and 18.23 in 2014, respectively. All herbicide treatments reduced the density of *M. quadrifoliata* significantly as compared to the non-treated control (Table 3). The lower density of *M. quadrifoliata* was observed in POE application of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (1.15 and 2.98 plants m<sup>-2</sup> in 2013 and 2.37 and 5.24 in 2014, respectively). Individual application of herbicide almix was noticed lower density of *M. quadrifoliata* and it was closely followed by bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> and metamifop 10% EC at 50 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> during both the years. The combined application of bispyribac sodium 4% SE + metamifop 10% SE at 42, 56 and 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> at all different doses as POE application were better control of weeds when herbicides was applied as individual like almix, clincher, bispyribac sodium and metamifop. Our findings revealed that almost all the different doses of new herbicide combination were effective for the control of grasses and sedges when compared to broad leaved weeds. This information is very helpful for DSR farmers in Tamil Nadu to achieve broad spectrum weed control.

#### 3.2.2. Total weed density

Significant variation in total weed density was observed among the herbicidal weed control treatments. During both the years, lesser total weed density was observed with POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> and bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter and it was closely followed by application of bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (16.80, 17.09, 22.50 in 2013 and 13.90, 15.43, 18.44 in 2014, respectively). At 40 DAHS also similar results were recorded (Table 3). Bispyribac sodium is pyrimidinyl

carboxate group which inhibits the biosynthesis of amino acids. Metamifop is aryloxyphenoxy propionate group which inhibits the activity of acetyl coenzyme-A carboxylase (ACCase) leading to growth retardation of weeds. However, the combined application of both herbicides induces chlorosis selectively in weeds and insufficient chlorophyll production makes it difficult for thrive of weeds. The combined application of these herbicides was better than their individual application in reducing the weed density, weed biomass and enhancing the productivity of rice yield. Total weed density was higher in individual application as POE application of clincher at 80 g a.i. ha<sup>-1</sup> when compared to almix at 4 g a.i. ha<sup>-1</sup> and it was similar in the both years of study. Clincher is a systemic POE herbicide and it is aryloxyphenoxy propionate group. In the present study, POE applications of clincher (alone) effectively control grassy weeds than compared to sedges and broad leaved weeds in direct seeded rice. Total weed density in the non-treated control were 105.20 and 156.13 plants m<sup>-2</sup> in 2013; 85.93 and 1132.78 plants m<sup>-2</sup> in 2014, respectively at 20 and 40 DAHS. All the herbicide treatments recorded lower total weed density significantly as compared to the non-treated control. Earlier, Mahajan and Chauhan (2013) revealed that sequential applications of pre and post-emergence herbicides provided better weed control than the sole application pre or post-emergence herbicides in DSR.

### 3.2.3. Total weed biomass

With regard to the total weed biomass, significant variation was observed among the herbicidal weed management practices in DSR. During both the years, lower total weed biomass was observed in POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (8.92 and 24.89 g m<sup>-2</sup> in 2013 and 11.38 and 34.56 g m<sup>-2</sup> in 2014, respectively), bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter (9.54 and 31.42 g m<sup>-2</sup> in 2013 and 13.45 and 37.58 g m<sup>-2</sup> in 2014, respectively) and it was closely followed by application of bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (16.77 and 36.76 g m<sup>-2</sup> in 2013 and 18.56 and 52.62 g m<sup>-2</sup> in 2014, respectively), bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (21.56 and 40.97 g m<sup>-2</sup> in 2013 and 24.63 and 64.82 g m<sup>-2</sup> in 2014, respectively) and individual application of almix at 4 g a.i./ha (24.41 and 44.91 g m<sup>-2</sup> in 2013 and 28.44 and 65.89 g m<sup>-2</sup> in 2014, respectively) as POE herbicides at 20 and 40 DAHS (Table 4). In the present study, herbicides

differed in respect of their efficacy and bispyribac sodium emerged as promising one in averting both density and dry matter accumulation by weeds. The performance of this herbicide could be attributed to reasonable suppression of weeds and selectivity to rice crop as well. It is a member of pyrimidinyloxy benzoic chemical family and inhibits acetolactate synthase enzyme in susceptible plants thus retarding the synthesis of branch chain amino acids (Darren and Stephen, 2006). The effectiveness of bispyribac sodium as a post-emergence herbicide for DSR is also reported elsewhere (Mahajan et al., 2009; Khaliq et al., 2011b). At 20 and 40 DAHS, POE of bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> obtained lower weed biomass as compared with application of almix at 80 g a.i. ha<sup>-1</sup> (24.41 and 44.91 g m<sup>-2</sup> in 2013 and 28.44 and 65.89 g m<sup>-2</sup> in 2014, respectively) and clincher at 80 g a.i. ha<sup>-1</sup> (26.79 and 49.81 g m<sup>-2</sup> in 2013 and 30.44 and 63.24 g m<sup>-2</sup> in 2014, respectively). Total weed biomass in the non-treated control were 70.97 and 116.83 g m<sup>-2</sup> in 2013; 110.56 and 188.67 g m<sup>-2</sup> in 2014, respectively at 20 and 40 DAHS. All the herbicide treatments recorded lower total weed biomass significantly as compared to the non-treated control.

#### 3.2.4. Weed control efficiency

Adoption of herbicide combination of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> treatment exhibited lowest weed infestation with higher weed control efficiency than sole herbicide application in the present study. During both the years, it was observed that POE application of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> resulted the higher weed control efficiency of 87.43 and 80.07% in 2013 and 88.45 and 81.68%, in 2014, respectively and it was followed by application of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter (86.55 and 73.10% in 2013 and 86.35 and 80.08%, in 2014, respectively). In the present study at 40 DAHS, weed control efficiency with ranged from 47.89 to 66.06% (single herbicide application); 60.22 to 80.07% (new herbicide combination) in 2013, respectively and WCE ranged from 55.67 to 66.48% (single herbicide application); 63.14 to 81.68% (new herbicide combination) in 2014, respectively (Table 4).

### 3.3. Effect on crop

#### 3.3.1. Response of grain yield

Rice grain yield following all herbicide treatments ranged from 4276 to 5676 kg ha<sup>-1</sup> and 4658 to 6388 kg ha<sup>-1</sup>, while the non-treated control plots yield of 2734 and 3012 kg ha<sup>-1</sup> in 2013 and 2014, respectively (Table 4). Higher grain yield was recorded in the plots treated with the POE application of new formulation bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (5676 kg ha<sup>-1</sup> in 2013 and 6388 kg ha<sup>-1</sup> in 2014) and it was similar to the grain yield observed in the plots treated with the application of herbicide combination of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter (5488 kg ha<sup>-1</sup> in 2013 and 6232 kg ha<sup>-1</sup> in 2014), bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> (5442 kg ha<sup>-1</sup> in 2013 and 6076 kg ha<sup>-1</sup> in 2014) and hand weeding twice on 25 and 45 DAS (5256 kg ha<sup>-1</sup> in 2013 and 5908 kg ha<sup>-1</sup> in 2014). Bispyribac sodium 4% SE + metamifop 10% SE showed on par with hand weeding twice on 25 and 45 DAS for most of the yield parameters and grain yield. These treatments recorded lesser crop weed competition during the critical period of rice that was marked as more panicles per unit area, increased kernel number and kernel weight over non-treated control. Higher grain yield in response to efficient weed control are reported elsewhere (Mahajan et al., 2009; Khaliq et al., 2011a, b; Akbar et al., 2011). Our data showed effectiveness of manual weeding in limiting weed density and dry biomass merely owing to POE application of new herbicide combination as an effective tool for their weed management in direct seeded rice. Nonetheless, during later part of the growing season weeds were also suppressed by shading effect of rice in manually weeded plots due to quick and dense canopy closure (Baloch et al., 2005). In both the years, grain yield in the plots treated with already existing molecule of almix at 4 g a.i. ha<sup>-1</sup> (4948 kg ha<sup>-1</sup> in 2013 and 5792 kg ha<sup>-1</sup> in 2014) and clincher at 80 g a.i. ha<sup>-1</sup> (4404 kg ha<sup>-1</sup> in 2013 and 5248 kg ha<sup>-1</sup> in 2014) was similar, but lower than grain yield recorded in the bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter. Though, the combined application of bispyribac sodium 4% SE + metamifop 10% SE + wetter with all different doses were very effective, provide broad spectrum weed control and subsequently increasing the productivity of direct seeded rice in this study.

### 3.3.2. *Weed index*

The best treatment with the maximum yield was taken as the base to work out the weed index that gives the magnitude of yield reduction due to weed competition in other treatments. New herbicide combination of POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70

g a.i. ha<sup>-1</sup> at 100 ml ha<sup>-1</sup> registered maximum grain yield and therefore lower yield reduction of only 3.31% and 2.44% were recorded in bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter during both the years (Table 4). The yield reduction in the treatment of bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> and bispyribac sodium 10% SC at 20 g a.i. ha<sup>-1</sup> were found to be 4.12 and 8.97% in 2013 and 4.88 and 7.47% in 2014, respectively. The higher yield reduction of 24.67% occurred under metamifop 10% EC at 50 g a.i. ha<sup>-1</sup> in 2013 and bispyribac sodium 4% SE + metamifop 10% SE at 42 g a.i. ha<sup>-1</sup> + wetter of 27.08 in 2014. In non-treated control plots of 51.83 in 2013 and 52.85% in 2014, respectively. The higher weed index resulted in reduced vegetative growth and nutrient availability to the crop could be the reason attributed for yield reduction at lower doses of bispyribac sodium 4% SE + metamifop 10% SE combination and individual application of herbicide. This emphasise the importance of proper weed management for increasing dry matter production of rice with reduced weed indices, thereby increasing the crop growth and grain yield.

### 3.3.3. *Phytotoxicity rating in direct seeded rice*

Application of new molecule herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at all different doses did not have any phytotoxicity effect on direct seeded rice. The phytotoxicity effect has been rated as “none”.

### 3.4. *Carryover effect on succeeding green gram*

During both the years of study, the residual effect of herbicides applied to rice crop; succeeding crop of green gram (cv. CO 6) was raised without disturbing the previous layout of the experiment. After the harvest of rice crop, the follow up crop (green gram) was dibbled in rice stubbles. A seed rate of 20 kg ha<sup>-1</sup> was adopted with a spacing of 30 cm x 10 cm.

#### 3.4.1. *Effect on weeds*

During both years of study, at 40 days after sowing (DAS), POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> was found significantly superior in registering lower total weed density in comparison to the other treatments. POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> without wetter, bispyribac sodium 10% SC at 20 g a.i./ha and metamifop 10% EC at 50 g a.i. ha<sup>-1</sup>



+ wetter at 100 ml ha<sup>-1</sup> were found on par with each other (Table 5). Non-treated control registered higher total weed density even in succeeding green gram crop taken after harvest of rice.

### 3.4.2. *Effect on crop*

#### 3.4.2.1. *Germination*

Germination percentage of green gram indicated that there was no significant difference among the treatments (Table 5). It is also clear that there was no residual toxicity due to the POE application of herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70, 56 and 42 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> at all different doses on the germination of the succeeding crop during both the years of study.

#### 3.4.2.2. *Number of pods per plant*

Number of pods per plant of green gram showed no significant difference among the weed control treatments. The number of pods per plant in all the treatments was comparable to the observations in that of non-treated control during both the years of study. So, there was no residual toxicity due to new formulation of herbicide combination of POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70, 56 and 42 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> on the performance of the succeeding crop (Table 5).

#### 3.4.2.3. *Seed yield of green gram*

Yield of green gram raised as succeeding crop showed no distinct variation due to different doses of POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70, 56 and 42 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> in DSR during both the years (Table 5).

## 4. **Conclusions**

The individual application of POE herbicides like bispyribac sodium, metamifop, almix and clincher did not control the complex weed flora in DSR. The plots treated with herbicide combination of bispyribac sodium 4% SE + metamifop 10% SE at 70, 56 and 42 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> at all different doses obtained higher grain yield because of lower total weed density and weed biomass in these herbicide combination treated plots when compared to individual herbicide application. Herbicide combination of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> effectively control of *Echinochloa*

*crus-galli* (L.) Beauv., *Dinebra retroflexa* (Vahl.) Panzer and *Panicum repens* L., among grasses; *Cyperus difformis* L. and *Cyperus irria* L. among sedges; *Marsilea quadrifolia* Linn and *Ammania baccifera* L. among broad leaved weeds with higher weed control efficiency of 80% at critical period of crop growth stage in DSR. Our study thus demonstrated that POE herbicide combination of bispyribac sodium 4% SE + metamifop 10% SE + wetter are needed for maintaining good weed control and grain yield in DSR. Sometimes, if farmers missed the application of pre-emergence herbicide due to erratic rains or any other reasons, effective weed control and high yield can still be obtained with this new herbicide combination of bispyribac sodium 4% SE + metamifop 10% SE + wetter.

Carryover effect study results showed that new formulation of POE herbicide combination bispyribac sodium 4% SE + metamifop 10% SE at 70, 56 and 42 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> at all different doses applied in DSR was found to be safe on the succeeding green gram and this might be due to detoxification of herbicides in soil and do not adversely affect the growth and yield of the succeeding crop in terms of germination percentage, number of pods per plant and seed yield of the green gram. Hence, it is concluded that POE application bispyribac sodium 4% SE + metamifop 10% SE at 70, 56 and 42 g a.i. ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> at all different doses of combination can be safely applied for weed control in DSR without any residual toxicity. However, the impact of continuous and inappropriate application of bispyribac sodium 4% SE + metamifop 10% SE combination in clay loam soil needs to be investigated to assess its risk potential to non-target organisms. As a final point, POE application of bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha<sup>-1</sup> + wetter at 100 ml ha<sup>-1</sup> can keep the total weed density and weed biomass reasonably at lower level and enhance the productivity of DSR. Thus a synergistic composition of bispyribac sodium 4% SE + metamifop 10% SE, when applied to DSR, allow a reduction in the amount of herbicide needed, greater flexibility in timing of the application besides offering broad spectrum weed control.

**Table 2. Effect of treatments on weed density (No.m<sup>-2</sup>) at 20 and 40 DAHS in direct seeded rice**

| Herbicide treatments  | Weed density (No.m <sup>-2</sup> ) |                 |                           |                 |                       |                 |                               |                 |                           |                |                       |                |
|---|------------------------------------|-----------------|---------------------------|-----------------|-----------------------|-----------------|-------------------------------|-----------------|---------------------------|----------------|-----------------------|----------------|
|   | rabi, 2013                         |                 |                           |                 |                       |                 | rabi, 2014                    |                 |                           |                |                       |                |
|   | <i>Echinochloa crus-galli</i>      |                 | <i>Dinebra retroflexa</i> |                 | <i>Panicum repens</i> |                 | <i>Echinochloa crus-galli</i> |                 | <i>Dinebra Retroflexa</i> |                | <i>Panicum repens</i> |                |
|   | 20 DAHS                            | 40 DAHS         | 20 DAHS                   | 40 DAHS         | 20 DAHS               | 40 DAHS         | 20 DAHS                       | 40 DAHS         | 20 DAHS                   | 40 DAHS        | 1.41 (0.00)           | 40 DAHS        |
| T <sub>1</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 42 g a.i.ha <sup>-1</sup> + wetter | 2.73<br>(5.43)                     | 3.40<br>(9.56)  | 1.79<br>(1.21)            | 2.28<br>(3.22)  | 1.41<br>(0.00)        | 2.04<br>(2.16)  | 3.84<br>(12.74)               | 5.91<br>(32.89) | 1.41<br>(0.00)            | 1.79<br>(1.22) | 1.41<br>(0.00)        | 1.96<br>(1.86) |
| T <sub>2</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i.ha <sup>-1</sup> + wetter | 2.42<br>(3.86)                     | 3.09<br>(7.55)  | 1.41<br>(0.00)            | 1.89<br>(1.56)  | 1.41<br>(0.00)        | 1.79<br>(1.22)  | 3.57<br>(10.76)               | 4.33<br>(16.78) | 1.41<br>(0.00)            | 1.69<br>(0.84) | 1.29<br>(0.00)        | 1.79<br>(1.22) |
| T <sub>3</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha <sup>-1</sup> + wetter | 2.07<br>(2.30)                     | 2.92<br>(6.54)  | 1.41<br>(0.00)            | 1.41<br>(0.00)  | 1.41<br>(0.00)        | 1.41<br>(0.00)  | 3.09<br>(7.52)                | 3.50<br>(10.24) | 1.30<br>(0.00)            | 1.41<br>(0.00) | 1.69<br>(0.84)        | 1.40<br>(0.00) |
| T <sub>4</sub> - Almix (Chlorimuron + Metsufuron 20% WP) at 4 g a.i.ha <sup>-1</sup>              | 2.76<br>(5.63)                     | 3.63<br>(11.19) | 2.45<br>(4.01)            | 2.83<br>(6.02)  | 1.85<br>(1.43)        | 2.56<br>(4.56)  | 4.11<br>(14.88)               | 5.31<br>(26.23) | 2.40<br>(3.74)            | 2.33<br>(3.42) | 1.74<br>(1.02)        | 2.29<br>(3.24) |
| T <sub>5</sub> - Clincher (Cyhalofop Buthyl 10% EC) at 80 g a.i.ha <sup>-1</sup>                  | 3.03<br>(7.21)                     | 3.84<br>(12.77) | 2.28<br>(3.22)            | 2.58<br>(4.64)  | 1.79<br>(2.44)        | 2.42<br>(3.86)  | 4.45<br>(17.76)               | 5.50<br>(30.42) | 2.24<br>(3.02)            | 2.78<br>(5.73) | 1.41<br>(0.00)        | 1.89<br>(1.56) |
| T <sub>6</sub> - Bispyribac sodium 10% SC at 20 g a.i.ha <sup>-1</sup> + wetter                   | 2.56<br>(4.56)                     | 4.07<br>(14.56) | 1.69<br>(0.85)            | 1.81<br>(1.26)  | 1.41<br>(0.00)        | 1.57<br>(0.45)  | 4.74<br>(20.45)               | 5.18<br>(24.85) | 1.66<br>(0.77)            | 1.70<br>(0.89) | 1.70<br>(0.88)        | 1.41<br>(0.00) |
| T <sub>7</sub> - Metamifop 10% SE at 50 g a.i.ha <sup>-1</sup> + wetter                           | 2.71<br>(5.32)                     | 3.54<br>(10.54) | 2.28<br>(3.22)            | 2.77<br>(5.68)  | 1.77<br>(1.12)        | 2.04<br>(2.18)  | 3.40<br>(9.56)                | 4.06<br>(14.52) | 2.10<br>(2.41)            | 2.36<br>(3.58) | 1.41<br>(0.00)        | 1.79<br>(1.21) |
| T <sub>8</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha <sup>-1</sup>          | 2.20<br>(2.86)                     | 3.14<br>(7.86)  | 1.41<br>(0.00)            | 1.68<br>(0.82)  | 1.41<br>(0)           | 1.69<br>(0.86)  | 3.10<br>(7.82)                | 3.91<br>(13.26) | 1.41<br>(0.00)            | 1.79<br>(1.22) | 1.66<br>(0.74)        | 1.65<br>(0.72) |
| T <sub>9</sub> - Bispyribac sodium 10% SC at 20 g a.i.ha <sup>-1</sup>                            | 2.88<br>(6.32)                     | 3.68<br>(11.56) | 2.09<br>(2.36)            | 2.41<br>(3.82)  | 1.79<br>(1.22)        | 2.21<br>(2.89)  | 4.46<br>(17.85)               | 5.88<br>(32.56) | 1.96<br>(1.86)            | 2.10<br>(2.42) | 2.11<br>(2.44)        | 1.88<br>(1.54) |
| T <sub>10</sub> - Metamifop 10% SE at 50 g a.i.ha <sup>-1</sup>                                   | 3.38<br>(9.42)                     | 4.12<br>(14.98) | 2.50<br>(4.24)            | 2.77<br>(5.66)  | 2.28<br>(3.22)        | 2.65<br>(5.02)  | 3.70<br>(11.72)               | 4.42<br>(17.56) | 2.29<br>(3.24)            | 2.49<br>(4.22) | 1.41<br>(0.00)        | 2.49<br>(4.21) |
| T <sub>11</sub> - Hand weeding twice on 25 and 45 DAS   | 4.53<br>(18.52)                    | 3.09<br>(7.54)  | 3.14<br>(7.86)            | 2.33<br>(3.42)  | 2.96<br>(6.78)        | 1.89<br>(1.56)  | 5.71<br>(30.56)               | 4.53<br>(18.56) | 3.39<br>(9.52)            | 1.89<br>(1.56) | 2.80<br>(5.86)        | 1.80<br>(1.24) |
| T <sub>12</sub> - Unsprayed control   | 4.73<br>(20.36)                    | 5.61<br>(29.45) | 3.40<br>(9.56)            | 4.03<br>(14.23) | 2.49<br>(7.42)        | 3.67<br>(11.46) | 6.04<br>(34.54)               | 7.67<br>(56.89) | 3.20<br>(8.24)            | 3.38<br>(9.45) | 2.90<br>(6.42)        | 3.23<br>(8.42) |
| <b>SEd</b>  | <b>0.34</b>                        | <b>0.45</b>     | <b>0.13</b>               | <b>0.24</b>     | <b>0.09</b>           | <b>0.19</b>     | <b>0.31</b>                   | <b>0.40</b>     | <b>0.11</b>               | <b>0.20</b>    | <b>0.07</b>           | <b>0.15</b>    |
| <b>CD (P=0.05)</b>  | <b>0.71</b>                        | <b>0.92</b>     | <b>0.25</b>               | <b>0.48</b>     | <b>0.18</b>           | <b>0.40</b>     | <b>0.63</b>                   | <b>0.82</b>     | <b>0.21</b>               | <b>0.42</b>    | <b>0.15</b>           | <b>0.31</b>    |

Figures in parenthesis are original values; Data subjected to square root transformation; DAHS: Days after herbicide spray

**Table 3. Effect of treatments on weed density and total weed density (No.m<sup>-2</sup>) at 20 and 40 DAHS in direct seeded rice**

| Herbicide treatments  | Weed density and total weed density (No./m <sup>2</sup> ) |                 |                               |                 |                    |                   |                          |                 |                               |                 |                    |                   |
|---|---|-----------------|-------------------------------|-----------------|--------------------|-------------------|--------------------------|-----------------|-------------------------------|-----------------|--------------------|-------------------|
|   | rabi, 2013  |                 |                               |                 |                    |                   | rabi, 2014               |                 |                               |                 |                    |                   |
|   | <i>Cyperus Difformis</i>                                  |                 | <i>Marsilea quadrifoliata</i> |                 | Total weed Density |                   | <i>Cyperus Difformis</i> |                 | <i>Marsilea quadrifoliata</i> |                 | Total weed Density |                   |
|   | 20 DAHS   | 40 DAHS         | 20 DAHS                       | 40 DAHS         | 20 DAHS            | 40 DAHS           | 20 DAHS                  | 40 DAHS         | 20 DAHS                       | 40 DAHS         | 20 DAHS            | 40 DAHS           |
| T <sub>1</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 42 g a.i.ha <sup>-1</sup> + wetter | 3.40<br>(9.54)  | 4.08<br>(14.65) | 2.32<br>(5.36)                | 2.75<br>(7.54)  | 5.83<br>(32.03)    | 6.87<br>(45.17)   | 1.95<br>(1.80)           | 3.41<br>(9.66)  | 2.89<br>(6.33)                | 3.59<br>(10.89) | 5.07<br>(23.68)    | 7.86<br>(59.80)   |
| T <sub>2</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i.ha <sup>-1</sup> + wetter | 3.09<br>(7.56)  | 3.27<br>(13.19) | 1.20<br>(1.45)                | 2.05<br>(4.21)  | 4.95<br>(22.50)    | 5.98<br>(33.81)   | 1.94<br>(1.77)           | 3.19<br>(8.20)  | 2.16<br>(2.67)                | 3.40<br>(9.56)  | 4.52<br>(18.44)    | 6.73<br>(43.35)   |
| T <sub>3</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha <sup>-1</sup> + wetter | 2.71<br>(5.32)  | 3.40<br>(9.56)  | 1.07<br>(1.15)                | 1.73<br>(2.98)  | 4.43<br>(16.80)    | 5.27<br>(25.78)   | 2.14<br>(2.56)           | 2.48<br>(4.16)  | 2.09<br>(2.37)                | 2.69<br>(5.24)  | 3.99<br>(13.90)    | 5.12<br>(24.19)   |
| T <sub>4</sub> - Almix (Chlorimuron + Metsufuron 20% WP) at 4 g a.i.ha <sup>-1</sup>              | 3.70<br>(11.68)   | 4.99<br>(22.89) | 3.25<br>(10.57)               | 2.76<br>(7.63)  | 6.75<br>(43.61)    | 7.79<br>(58.76)   | 3.67<br>(11.47)          | 4.46<br>(17.90) | 2.58<br>(4.65)                | 3.35<br>(9.21)  | 6.01<br>(34.13)    | 8.07<br>(63.14)   |
| T <sub>5</sub> - Clincher (Cyhalofop Buthyl 10% EC) at 80 g a.i.ha <sup>-1</sup>                  | 4.07<br>(14.56)   | 3.42<br>(20.19) | 3.52<br>(12.36)               | 3.03<br>(9.21)  | 7.51<br>(54.47)    | 6.99<br>(46.88)   | 3.28<br>(8.77)           | 4.15<br>(15.20) | 2.87<br>(6.23)                | 3.99<br>(13.89) | 6.04<br>(34.54)    | 8.43<br>(69.06)   |
| T <sub>6</sub> - Bispyribac sodium 10% SC at 20 g a.i.ha <sup>-1</sup> + wetter                   | 3.24<br>(8.47)  | 4.30<br>(16.45) | 2.89<br>(8.34)                | 3.40<br>(11.56) | 5.84<br>(32.12)    | 7.50<br>(54.26)   | 2.65<br>(5.03)           | 3.67<br>(11.46) | 2.80<br>(5.86)                | 3.23<br>(8.42)  | 5.81<br>(31.70)    | 6.86<br>(45.05)   |
| T <sub>7</sub> - Metamifop 10% SE at 50 g a.i.ha <sup>-1</sup> + wetter                           | 4.75<br>(20.56)   | 3.52<br>(26.19) | 1.86<br>(3.45)                | 2.64<br>(6.98)  | 6.81<br>(44.41)    | 6.79<br>(44.08)   | 4.10<br>(14.77)          | 4.82<br>(21.20) | 2.58<br>(4.67)                | 4.18<br>(15.46) | 6.09<br>(35.13)    | 7.44<br>(53.29)   |
| T <sub>8</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha <sup>-1</sup>          | 2.72<br>(5.38)  | 3.22<br>(11.01) | 1.11<br>(1.24)                | 2.07<br>(4.30)  | 4.37<br>(17.09)    | 5.39<br>(27.02)   | 2.19<br>(2.58)           | 2.56<br>(4.57)  | 2.11<br>(2.46)                | 3.00<br>(7.02)  | 4.17<br>(15.43)    | 5.76<br>(31.23)   |
| T <sub>9</sub> - Bispyribac sodium 10% SC at 20 g a.i.ha <sup>-1</sup>                            | 3.81<br>(12.5)  | 4.64<br>(19.54) | 3.57<br>(12.78)               | 2.83<br>(8.00)  | 6.70<br>(42.91)    | 7.29<br>(51.15)   | 3.18<br>(8.12)           | 4.07<br>(14.55) | 2.71<br>(5.34)                | 3.45<br>(9.90)  | 5.95<br>(33.40)    | 8.17<br>(64.69)   |
| T <sub>10</sub> - Metamifop 10% SE at 50 g a.i.ha <sup>-1</sup>                                   | 4.49<br>(18.16)   | 5.19<br>(24.98) | 2.75<br>(7.56)                | 3.40<br>(11.57) | 7.43<br>(53.20)    | 8.55<br>(71.14)   | 3.94<br>(13.56)          | 4.96<br>(22.56) | 3.28<br>(8.78)                | 4.10<br>(14.85) | 6.49<br>(40.09)    | 8.03<br>(62.44)   |
| T <sub>11</sub> - Hand weeding twice on 25 and 45 DAS   | 5.71<br>(30.56)   | 4.07<br>(14.56) | 4.07<br>(14.56)               | 2.96<br>(6.78)  | 9.22<br>(82.93)    | 6.37<br>(38.54)   | 4.63<br>(19.48)          | 3.54<br>(10.56) | 4.08<br>(14.62)               | 3.06<br>(7.34)  | 9.21<br>(82.90)    | 6.84<br>(44.75)   |
| T <sub>12</sub> - Unsprayed control   | 6.43<br>(39.4)  | 7.38<br>(52.46) | 4.42<br>(17.52)               | 5.24<br>(32.45) | 10.35<br>(105.02)  | 12.57<br>(156.13) | 4.49<br>(18.14)          | 5.62<br>(29.54) | 3.96<br>(13.67)               | 4.50<br>(18.23) | 8.77<br>(85.93)    | 11.61<br>(132.78) |
| <b>SEd</b>  | <b>0.52</b>   | <b>0.62</b>     | <b>0.23</b>                   | <b>0.38</b>     | <b>0.65</b>        | <b>0.62</b>       | <b>0.29</b>              | <b>0.40</b>     | <b>0.26</b>                   | <b>0.31</b>     | <b>0.50</b>        | <b>0.60</b>       |
| <b>CD (P=0.05)</b>  | <b>1.07</b>   | <b>1.27</b>     | <b>0.48</b>                   | <b>0.79</b>     | <b>1.38</b>        | <b>1.28</b>       | <b>0.61</b>              | <b>0.81</b>     | <b>0.52</b>                   | <b>0.64</b>     | <b>1.02</b>        | <b>1.23</b>       |

Figures in parenthesis are original values; Data subjected to square root transformation; DAHS: Days after herbicide spray

**Table 4. Total weed dry weight, weed control efficiency, grain yield and weed index as influenced by different weed management practices in direct seeded rice**

| Herbicide treatments  | Total weed dry weight (g/m <sup>2</sup> ), WCE (%), grain yield (kg/ha) & Weed Index (WI) |                 |         |         |             |            |   |                 |         |         |             |            |
|---|---|-----------------|---------|---------|-------------|------------|---|-----------------|---------|---------|-------------|------------|
|   | rabi, 2013  |                 |         |         |             |            | rabi, 2014                                |                 |         |         |             |            |
|   | Total weed dry weight (g/m <sup>2</sup> )   |                 | WCE (%) |         | Grain yield | Weed Index | Total weed dry weight (g/m <sup>2</sup> ) |                 | WCE (%) |         | Grain yield | Weed Index |
|   | 20 DAHS   | 40 DAHS         | 20 DAHS | 40 DAHS |             |            | 20 DAHS                                   | 40 DAHS         | 20 DAHS | 40 DAHS |             |            |
| T <sub>1</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 42 g a.i.ha <sup>-1</sup> + wetter | 5.49<br>(23.18)   | 6.76<br>(47.68) | 67.33   | 60.22   | 4286        | 24.49      | 5.40<br>(27.11)                           | 8.34<br>(69.54) | 72.49   | 63.14   | 4658        | 27.08      |
| T <sub>2</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i.ha <sup>-1</sup> + wetter | 4.10<br>(16.77)   | 5.90<br>(36.76) | 76.37   | 69.73   | 4978        | 12.30      | 4.53<br>(18.56)                           | 7.39<br>(52.62) | 81.17   | 72.11   | 5722        | 10.43      |
| T <sub>3</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha <sup>-1</sup> + wetter | 2.69<br>(8.92)  | 4.78<br>(24.89) | 87.43   | 80.07   | 5676        | 0.00       | 3.66<br>(11.38)                           | 6.05<br>(34.56) | 88.45   | 81.68   | 6388        | 0.00       |
| T <sub>4</sub> - Almix (Chlorimuron + Metsufuron 20% WP) at 4 g a.i.ha <sup>-1</sup>              | 5.24<br>(24.41)   | 6.55<br>(44.91) | 65.60   | 62.63   | 4948        | 12.83      | 5.52<br>(28.44)                           | 8.12<br>(65.89) | 71.14   | 65.08   | 5792        | 9.33       |
| T <sub>5</sub> - Clincher (Cyhalofop Buthyl 10% EC) at 80 g a.i.ha <sup>-1</sup>                  | 5.27<br>(26.79)   | 6.91<br>(49.81) | 62.25   | 58.36   | 4404        | 22.41      | 5.70<br>(30.44)                           | 7.95<br>(63.24) | 69.12   | 66.48   | 5248        | 17.85      |
| T <sub>6</sub> - Bispyribac sodium 10% SC at 20 g a.i.ha <sup>-1</sup> + wetter                   | 4.64<br>(21.56)   | 6.24<br>(40.97) | 69.62   | 66.06   | 5442        | 4.12       | 5.16<br>(24.63)                           | 8.05<br>(64.82) | 75.01   | 65.64   | 6076        | 4.88       |
| T <sub>7</sub> - Metamifop 10% SE at 50 g a.i.ha <sup>-1</sup> + wetter                           | 5.39<br>(26.03)   | 7.56<br>(59.16) | 63.32   | 50.22   | 5004        | 11.84      | 5.49<br>(28.19)                           | 8.52<br>(72.61) | 71.40   | 61.51   | 5748        | 10.02      |
| T <sub>8</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha <sup>-1</sup>          | 2.92<br>(9.54)  | 5.68<br>(31.42) | 86.55   | 73.10   | 5488        | 3.31       | 3.93<br>(13.45)                           | 6.29<br>(37.58) | 86.35   | 80.08   | 6232        | 2.44       |
| T <sub>9</sub> - Bispyribac sodium 10% SC at 20 g a.i.ha <sup>-1</sup>                            | 5.28<br>(27.86)   | 7.08<br>(52.10) | 60.74   | 59.00   | 5167        | 8.97       | 5.87<br>(32.51)                           | 8.49<br>(72.15) | 67.02   | 61.76   | 5911        | 7.47       |
| T <sub>10</sub> - Metamifop 10% SE at 50 g a.i.ha <sup>-1</sup>                                   | 5.45<br>(29.70)   | 7.74<br>(61.84) | 58.15   | 47.89   | 4276        | 24.67      | 6.18<br>(36.19)                           | 9.15<br>(83.64) | 63.28   | 55.67   | 4968        | 22.23      |

|   |                 |                   |       |       |            |       |                   |                   |      |       |            |       |
|---|-----------------|-------------------|-------|-------|------------|-------|-------------------|-------------------|------|-------|------------|-------|
| T <sub>11</sub> - Hand weeding twice on 25 and 45 DAS | 7.39<br>(52.55) | 6.15<br>(35.84)   | 25.95 | 69.32 | 5256       | 7.40  | 10.33<br>(104.63) | 7.20<br>(49.87)   | 5.36 | 73.56 | 5908       | 7.51  |
| T <sub>12</sub> - Unsprayed control                   | 8.42<br>(70.97) | 10.72<br>(116.83) | -     | -     | 2734       | 51.83 | 10.03<br>(110.56) | 13.81<br>(188.67) | -    | -     | 3012       | 52.85 |
| <b>SEd</b>  | <b>0.58</b>     | <b>0.88</b>       | -     | -     | <b>352</b> | -     | <b>0.61</b>       | <b>0.87</b>       | -    | -     | <b>309</b> | -     |
| <b>CD (P=0.05)</b>                                    | <b>1.21</b>     | <b>1.79</b>       | -     | -     | <b>688</b> | -     | <b>1.23</b>       | <b>1.76</b>       | -    | -     | <b>623</b> | -     |

Figures in parenthesis are original values; Data subjected to square root transformation; DAHS: Days after herbicide spray

**Table 5. Effect of treatments on weed density, germination percentage, number of pods plant<sup>-1</sup> and seed yield of succeeding green gram**

| Herbicide treatments  | Succeeding green gram crop                         |                    |                                       |   |  |                    |                                       |                          |
|---|--|--------------------|---------------------------------------|---|--|--------------------|---------------------------------------|--------------------------|
|   | <i>rabi, 2013</i>                                  |                    |                                       |   | <i>rabi, 2014</i>                                  |                    |                                       |                          |
|   | Weed density<br>(No./m <sup>2</sup> )<br>at 40 DAS | Germination<br>(%) | Number of<br>pods plant <sup>-1</sup> | Seed<br>yield<br>(kg ha <sup>-1</sup> ) | Weed density<br>(No./m <sup>2</sup> )<br>at 40 DAS | Germination<br>(%) | Number of<br>pods plant <sup>-1</sup> | Seed<br>yield<br>(kg/ha) |
| T <sub>1</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 42 g a.i.ha <sup>-1</sup> + wetter | 6.51<br>(40.32)                                    | 82.99              | 21.67                                 | 622                                     | 7.77<br>(58.44)                                    | 87.56              | 24.89                                 | 660                      |
| T <sub>2</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 56 g a.i.ha <sup>-1</sup> + wetter | 6.14<br>(35.65)                                    | 85.55              | 24.33                                 | 655                                     | 7.32<br>(51.62)                                    | 89.31              | 27.62                                 | 694                      |
| T <sub>3</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha <sup>-1</sup> + wetter | 4.75<br>(20.58)                                    | 82.32              | 21.00                                 | 667                                     | 6.29<br>(37.54)                                    | 90.56              | 30.24                                 | 672                      |
| T <sub>4</sub> - Almix (Chlorimuron + Metsufuron 20% WP) at 4 g a.i.ha <sup>-1</sup>              | 6.17<br>(36.05)                                    | 82.55              | 24.00                                 | 602                                     | 7.44<br>(53.33)                                    | 87.41              | 27.14                                 | 652                      |
| T <sub>5</sub> - Clincher (Cyhalofop Buthyl 10% EC) at 80 g a.i.ha <sup>-1</sup>                  | 6.06<br>(34.78)                                    | 84.99              | 23.67                                 | 615                                     | 8.08<br>(63.24)                                    | 89.85              | 28.32                                 | 643                      |
| T <sub>6</sub> - Bispyribac sodium 10% SC at 20 g a.i.ha <sup>-1</sup> + wetter                   | 6.32<br>(37.88)                                    | 85.52              | 23.33                                 | 620                                     | 7.36<br>(52.13)                                    | 90.38              | 27.98                                 | 647                      |
| T <sub>7</sub> - Metamifop 10% SE at 50 g a.i.ha <sup>-1</sup> + wetter                           | 5.50<br>(28.30)                                    | 81.99              | 24.00                                 | 567                                     | 8.01<br>(61.98)                                    | 89.85              | 28.65                                 | 623                      |
| T <sub>8</sub> - Bispyribac sodium 4% SE + metamifop 10% SE at 70 g a.i.ha <sup>-1</sup>          | 5.14<br>(24.42)                                    | 84.45              | 23.67                                 | 630                                     | 6.71<br>(43.08)                                    | 89.31              | 29.87                                 | 668                      |
| T <sub>9</sub> - Bispyribac sodium 10% SC at 20 g a.i.ha <sup>-1</sup>                            | 5.21<br>(25.18)                                    | 87.94              | 24.26                                 | 653                                     | 7.24<br>(50.37)                                    | 89.46              | 27.56                                 | 667                      |

|   |                 |       |             |           |                 |       |             |           |
|---|-----------------|-------|-------------|-----------|-----------------|-------|-------------|-----------|
| T <sub>10</sub> - Metamifop 10% SE at 50 g a.i.ha <sup>-1</sup> | 6.02<br>(34.24) | 84.45 | 24.38       | 649       | 8.32<br>(67.21) | 90.41 | 29.76       | 684       |
| T <sub>11</sub> - Hand weeding twice on 25 and 45 DAS           | 7.70<br>(57.26) | 84.33 | 24.27       | 644       | 9.42<br>(86.81) | 88.56 | 26.54       | 672       |
| T <sub>12</sub> - Unsprayed control                             | 8.02<br>(62.3)  | 84.99 | 24.00       | 586       | 9.67<br>(91.47) | 89.85 | 28.65       | 528       |
| <b>SEd</b>  | <b>0.45</b>     | -     | <b>0.85</b> | <b>62</b> | <b>0.51</b>     | -     | <b>2.17</b> | <b>71</b> |
| <b>CD (P=0.05)</b>  | <b>0.92</b>     | -     | <b>NS</b>   | <b>NS</b> | <b>1.03</b>     | -     | <b>NS</b>   | <b>NS</b> |

Figures in parenthesis are original values;      Data subjected to square root transformation;      DAS: Days after sowing

**References**

- Akbar, N., Ehsanullah, K., Jabran., Ali, M.A., 2011. Weed management improves yield and quality of direct seeded rice. *Aus. J. Crop Sci.* 5 (6), 688-694.
- Balasubramanian, V., Hill, J., 2000. Direct wet seeding of rice in Asia: Emerging issues and strategic research needs for the 21 century. Annual Workshop of the Directorate of Rice Res., Hyderabad, Andhra Pradesh.
- Baloch, M.S., Hassan, G., Morimoto, T., 2005. Weeding techniques in transplanted and wet seeded rice in Pakistan. *Weed Biolog. Mang.* 5, 190-196.
- Chauhan, B.S., 2012. Weed ecology and weed management strategies for dryseeded rice in Asia. *Weed Technol.* 26, 1-13.
- Chauhan, B.S., 2012a. Weed Management in Direct-Seeded Rice Systems. International Rice Research Institute, Los Banos, Philippines. p. 20.
- Chauhan, B.S., 2012b. Weed ecology and weed management strategies for dry-seeded rice in Asia *Weed Technol.* 26, 1-13.
- Chauhan, B.S., Johnson, D.E., 2011. Row spacing and weed control timing affect yield of aerobic rice. *Field Crops Res.* 121, 226-231.
- Chauhan, B.S., Opena, J., 2013a. Implications of plant geometry and weed control options in designing a low-seeding seed-drill for dry-seeded rice systems. *Field Crops Res.* 144, 225-231.
- Chauhan, B.S., Opena, J., 2013b. Weed management and grain yield of rice sown at low seeding rates in mechanized dry-seeded systems. *Field Crops Res.* 141: 9-15.
- Darren, W.L., Stephen, E.H., 2006. Foliar and root absorption and translocation of bispyribacsodium in cool-season turf grass. *Weed Technol.* 20 (4), 1015-1022.
- Gill, G.S., Vijaya Kumar, K., 1969. Weed Index - A new method of reporting weed control trials. *Indian J. Agron.*, 14, 96-98.
- Gomez, K.A., Gomez, A.A., 2010. Statistical procedures for Agricultural Research. Wiley India Pvt. Ltd., New Delhi, India.
- Khaliq, A., Matloob, A., 2011. Weed-crop competition period in three fine rice cultivars under direct-seeded rice culture. *Pak. J. Weed Sci. Res.* 17, 229-243.



- Khaliq, A., Matloob, A., Shafique, H.M., Farooq, M., Wahid, A., 2011b. Evaluating sequential application of pre and post emergence herbicides in dry seeded finer rice. *Pakistan J. Weed Sci. Res.* 17, 111-123.
- Khaliq, A., Riaz, Y., Matloob, A., 2011a. Bioeconomic assessment of chemical and nonchemical weed management strategies in dry seeded fine rice (*Oryza sativa* L.). *J. Plant Breeding Crop Sci.* 3 (12), 302-310.
- Kim, K.U., 1996. Ecological forces influencing weed competition and herbicide resistance. In: Naylor, R. (Ed.), *Herbicides in Asian Rice: Transitions in Weed Management*. Stamford University, California and IRRI, Philippines, pp. 129-142.
- Kumaran, S.T., Kathiresan, G., Murali Arthanari, P., Chinnusamy, C., Sanjivkumar, V., 2015. Efficacy of new herbicide (bispribac sodium 10% SC) against different weed flora, nutrient uptake in rice and their residual effects on succeeding crop of green gram under zero tillage. *J. Appl. and Natural Sci.* 7 (1), 279-285.
- Mahajan, G., Chauhan, B.S., 2013. Herbicide options for weed control in dry-seeded aromatic rice in India. *Weed Technol.* 27, 682-689.
- Mahajan, G., Chauhan, B.S., 2013. The role of cultivars in managing weeds in dryseeded rice production systems. *Crop Prot.* 49, 52-57.
- Mahajan, G., Chauhan, B.S., Johnson, D.E., 2009. Weed management in aerobic rice in Northwestern Indo-Gangetic Plains. *J. Crop Improv.* 23 (4), 366-382.
- Mahajan, G., Timsina, J., 2011. Effect of nitrogen rates and weed control methods on weeds abundance and yield of direct seeded rice. *Arch. Agron. Soil Sci.* 57, 239-250.
- Mani, V.S., Mala, M.L., Gautam, K.C., Bhagavandas., 1973. Weed killing chemicals in potato cultivation. *Indian Fmg.*, 23 (1), 17-18.
- Oerke, E.C., Dehne, H.W., 2004. Safeguarding production losses in major crops and the role of crop protection. *Crop Prot.* 23, 275-285.
- Powles, S.B., Yu, Q., 2010. Evolution in action: plants resistant to herbicides. *Ann. Rev. Plant Biol.* 61, 317-347.
- Rao, A.N., Nagamani. A., 2007. Available technologies and future research challenges for managing weeds in dryseeded rice in India. In: *Proc. 21<sup>st</sup> Asian Pacific Weed Sci. Soc.* Colombo, Sri Lanka on 2-6 October.

- Rao, A.N., Wani, S.P., Ladha, J.K., 2014. Weed management research in India an analysis of the past and outlook for future. In: Souvenir (1989-2014). Directorate of Weed Research, Jabalpur, India, DWR Publication No.18, pp. 1-26.
- Singh, G., 2008. Integrated weed management in direct-seeded rice. In Y. Singh, V.P. Singh, B. Chauhan, A. Orr, A.M. Mortimer, D.E. Johnson and B. Hardy eds., Direct Seeding of Rice and Weed Management in the Irrigated Rice-Wheat Cropping System of the Indo-Gangetic Plains. International Rice Research Institute, Los Banos, Philippines: Directorate of Experiment Station, G.B. Pant University of Agriculture and Technology, Pantnagar. 161-175.
- Snedecor, G.W., Cochran, W.G., 1967. Statistical Methods. Oxford and IBH Publ. Co., New Delhi. p. 593.
- Sunil, C.M., Shekara, B.G., Kalyanmurthy, K.N., Shankaralingapa, B.C., 2010. Growth and yield of aerobic rice as influenced by integrated weed management practices. Ind J Weed Sci. 42 (3&4), 180-183.
- Zhao, D., 2006. Weed competitiveness and yielding of aerobic rice genotypes. PhD thesis, Wageningen University, The Netherlands.