

Characterization of *Bacillus cereus* symbiotic to hemi-parasitic plant *Santalum album* L.

Short title: Characterization of *Bacillus cereus* associated with *Santalum album* Linn.

Siuli Batabyal¹, Priyanka Mukhopadhyay², Soumendranath Chatterjee^{2*}, Jagatpati Tah¹ and Nimai Chandra Saha³

¹ Cyto-Genetics and Molecular Biology Laboratory, Department of Botany, The University of Burdwan, Burdwan 713104, West Bengal, India

² Microbiology and Parasitology Research Laboratory, Department of Zoology, The University of Burdwan 713104, West Bengal, India

³ Aquatic Toxicology and Fisheries Research Laboratory, Department of Zoology, The University of Burdwan 713104, West Bengal, India

Abstract:

Aims:

Santalum album L., known as sandalwood plant (white sandal) belongs to the family Santalaceae, is characteristically a hemi-parasite that requires host plant in the early stages for the better growth and development. Besides its extreme economic importance, significant work has not been done to reveal the relationship of beneficial microorganisms with these plants for their better growth and development. Present investigation is an attempt to isolate and characterize the rhizospheric soil bacteria of *Santalum album* L. occurring in some areas of Bankura district of West Bengal, India.

Methodology:

The microbial colonies in the soils were estimated as colony forming units (cfu/g dr.soil) from plates prepared by different medium. Phenotypic, biochemical and molecular characters of the bacteria were studied following standard methods. The physico-chemical parameters, and

microbial population was determined on the rhizospheric soil of the hemiparasitic sandalwood plant *Santalum album* Linn. occurring at four locations of Bankura district in West Bengal, India.

Results:

The population diversity of cultivable heterotrophic, Gram negative, nitrifying, phosphate solubilizing, starch hydrolyzing, spore forming bacteria were higher at Hirbandh with higher organic carbon level than other three locations. Bacterial population was comparatively lower in Basudevpur due to lower water holding capacity. One spore forming bacterium (SW1) was isolated from Hirbandh soil. The isolate (SW1) was characterized by phenotypic properties, scanning electron microscopy, biochemical properties, analysis of fatty acid methyl esters and 16S rRNA gene sequence and identified as *Bacillus cereus* (KT626448) which branched with *Bacillus cereus* BSFN12r (KM405329) with 100% bootstrap support.

Conclusion:

Present investigation is an attempt to isolate and characterize the rhizospheric soil bacteria from economically important plant *Santalum album* L. Further studies may find out the positive role of the symbiotic association of *Bacillus cereus* (SW1) with the root of *S. album* as a key factor for the better growth and development of this economically important plant occurring in Bankura district, West Bengal, India.

Key words: *Santalum album* L., Rhizospheric bacteria, *Bacillus cereus* (SW1), Scanning Electron Microscopy, 16S rRNA gene sequence

Introduction

Santalum album L., commonly known as sandalwood plant (white sandal) belongs to the fam. *Santalaceae* comprising of more than 15 species and their variants. It is characteristically a hemiparasite and requires host plant in the early stages for the better growth and development. Rao (1911) reports that sandal seedlings were incapable to grow beyond one year without haustoria, and confirmed its selectivity for host as although almost all plants in its surroundings may be

attacked but the better growth was observed in association with *Pongamia pinnata*, *Albizia lebbbeck*, *Tectona grandis* etc (Rao 1911). Though the plant grows naturally in wide agroclimatic conditions like warm desert of Australia, dry and monsoon climate of India, Vanuatu, eastern Indonesia, subtropical climate of Hawaii and New Caledonia where receives almost uniform rainfall but in India the plant is mostly restricted in southern part only. Presently, there are a few patches in one or two districts of West Bengal with white sandal plants, but it can be introduced in many more areas of West Bengal because it can adapt to various soils although prefers light to medium and well drained soil (Merlin *et al.* 2006; Rao *et al.* 2011). Fragrant wood and essential oil obtained from sandal are used for the preparation of incense, perfumes, carving and medicine. In spite of its extreme economic importance, significant work has not been done to reveal the relationship of beneficial microorganisms with this plant for their better growth and development. Present study is an attempt to isolate and characterize the rhizospheric soil bacteria of *Santalum album* L. occurring at some areas of Bankura district of West Bengal, India in relation to its possible role on the growth of the plant.

Materials and methods

Site of soil collection

The rhizospheric soil samples were collected from four different places viz., Hirbundh, Basudevpur, Bagaldhara and Maitybundh of Bankura district of West Bengal, India.

Isolation of the bacteria from soil

Hundred grams of soil samples from the rhizosphere of the plant were collected from different areas of Bankura district of West Bengal, India. Soil samples were mixed thoroughly and the soils were put separately in sterile polythene bags, sealed with rubber bands and analyzed in the laboratory. Viable aerobic bacterial population were assessed from the plates prepared from the soil suspensions after incubating the plates at 30 ± 0 °C for required days (3-21 d) in a BOD incubator. The microbial colonies in the soils were enumerated as colony forming units (cfu/g dr. soil) from plates prepared with 10 µl soil suspension of (10^{-2}) dilution mixed with 100 ml of different medium. Soil suspension was heated at $60^{\circ}\text{C} \pm 0.1^{\circ}\text{C}$ for 30 minutes for enrichment culture of the spore formers. Gram negative bacterial population were determined in Nutrient Agar (NA) medium (g/l: peptone 5, beef extract 3, agar 2, pH 7.0) and crystal violet solution (peptone 5 g/l, beef extract 3 g/l, lactose 10 g/l, crystal violet 0.0033 g/l, agar 15 g/l, pH $6.8 \pm$

0.1) was added to the medium before plating (Pelczar *et al.* 1957; Lacey. 1997; Dangar *et al.* 2010; Chatterjee *et al.* 2012 ; Chatterjee *et al.* 2015). To determine gram-negative bacterial population, crystal violet (0.01 g/l) was added to the medium before plating. The nitrifying bacterial population were assessed on Winogradsky's medium containing (NH₄)₂SO₄ (1.0 g/l) and the colonies were visualized (pink colour) by flooding the plates with sulphanillic acid reagent (Pelczar *et al.* 1957). Nitrifying bacterial colonies were recorded from 5-30 d (5 d intervals) but other colonies were counted after 3 day of incubation. The inorganic phosphate solubilizing bacteria were assessed from the halo zone formation around the bacterial colonies on the insoluble phosphate [Ca₃ (PO₄)₂] containing medium Pelczar and coworkers (1957). The asymbiotic nitrogen fixing bacterial populations were determined on the nitrogen free medium (Pelczar 1957).

Characterization of the bacterial isolate

The predominant bacterial colonies isolated from the medium were purified and characterized. The cultural and morphological characters viz. shape, size, elevation, margin, color, opacity and consistency of the colonies were recorded. Phenotypic and staining properties of the bacteria were studied. Antibiotic sensitivity tests were done with standard antibiotic discs (Brown 2007). The bacterial isolate was identified on the basis of biochemical properties, FAME analysis (MIS,MIDI,Sharlock®USA) and 16S rRNA gene sequence analysis (Janssen 1994). The spore forming bacterial isolates were observed under scanning electron microscope. The smear preparation of bacterial suspension was done on a cover glass, air dried and heat fixed over a flame for one to two seconds followed by 2.5% glutaraldehyde (aqueous) for 45 min. Slides were then dehydrated passing through 50%, 70%, 90% and finally with absolute alcohol for 5 min. each. Then the gold coated suspensions were observed under scanning electron microscope (HITACHI S-530). For the fatty acid methyl ester analysis (FAME), whole cell fatty acids were converted to methyl ester and analysed by gas chromatography. The fatty acid methyl ester composition of bacterial isolates was compared to Sherlock library of known bacterial strains in order to find a closest match.

16S rRNA gene sequence analysis

Pure cultured colony of bacterial isolate SW1 was picked up with a sterilized toothpick, suspended in 0.5 ml of sterilized saline in a 1.5 ml centrifuge tube and centrifuged at 10,000 rpm for 10 min. After removal of supernatant, the pellet was suspended in 0.5 ml of Insta Gene Matrix (Bio-Rad, USA), incubated at 56°C for 30 min and then heated 100°C for 10 min. After heating, supernatant was used for PCR. The PCR reaction was prepared with 1µl of template DNA in 20 µl of PCR reaction solution using primers and amplified for 35 cycles at 94°C for 45 sec., 55°C for 60 sec. and 72°C for 60 sec. which produced about 1,400 bp DNA fragment. dNTPs from PCR products were purified by using Montage PCR clean up kit (Millipore) and the purified PCR products were sequenced using the forward and reverse primers.

Results and discussion

Population dynamics of microorganisms in rhizospheric soils of *S. album* showed that the total aerobic heterotrophic bacteria ranged from 2.9×10^7 to 3.3×10^7 cfu/g, the nitrifying bacterial were also higher ranging from 5.2×10^6 cfu/g in Basudevpur to 5.7×10^6 cfu/g in Hirbandh, phosphate solubilising bacteria ranged from 7.9×10^2 cfu/g (Basudevpur) to 8.3×10^2 cfu/g (Hirbandh), starch hydrolyzing bacteria ranged from 8.0×10^4 to 10.0×10^4 cfu/g and spore forming bacteria ranged from 7.5×10^4 to 8.0×10^4 cfu/g (Table 1).

The organic carbon level varied from 0.37 to 0.51%, nitrogen, phosphorus, potassium and soil pH were also recorded (Table 2). The soil physico-chemical parameters, as well as, the soil types are important factors which influence the soil microbial community. Rhizospheric soil is a hot spot of bacterial diversity and harbours those bacterial strains that may have some impact on soil functional status, as well as, the growth of the plant.

The bacterial isolate (SW1) was found in all rhizospheric soil samples of *S. album* of all locations throughout the year. The bacterial colony was off-white with smooth margin and vegetative cell was more than 1 µm and rod shaped. Under SEM study only spores were visualized but no crystal was detected (Fig.1). The isolate was positive for the tests of catalase, methyl red- Vogues-Proskauer, starch, casein and gelatin hydrolysis and negative for citrate utilization test (Table 3). The isolate reduced nitrate to nitrite which reflects its role nitrogen metabolism in the rhizospheric soil (Table 3). The strain was sensitive to nalidixic acid (30 µg/disc), doxycycline (30 µg/disc), bacitracin (10 µg/disc) and tetracycline (30 µg/disc), and

resistant to amoxycillin (10 µg/disc), ampicillin (10 µg/disc), polymyxin-B (50 µg/disc) and nystatin (100 µg/disc; Table 3). On the basis of morpho-physiological and biochemical properties, the bacterium SW1 was identified as *Bacillus* sp. Through the FAME analysis, 16:1w7c alcohol, 17 isow10c fatty acid supports that the organism belongs to the genus *Bacillus* and the 17:1 iso w5c, 12:0 iso fatty acid supports that the organism belongs to the species *Bacillus cereus* (Fig. 2). Phylogenetic affiliation of the bacterial isolate reveals that *Bacillus* sp. SW1 (KT626448) branched with *Bacillus cereus* BSFN12r (KM405329) with 100% bootstrap support (Fig. 3) which confirmed the identity of SW1 as *Bacillus cereus* (Logan *et al.* 2009). The nucleotide base composition of 16 rRNA gene sequence of the bacterial isolate SW1 was determined which revealed that the AT and GC content were 47.16% and 52.84%, respectively (Table 3).

It has already been reported that different strains of *Bacillus* act as plant growth promoters for *Saccharum officinerum* sugarcane (Nakade *et al.* 2013) and *Triticum aestivum* (Rawat *et al.* 2011) belonging to the family Poaceae. *B. cereus* has been proved to be a growth promoting rhizobacteria of some plants viz.. *Brassica juncea* (Aziz *et al.* 2012), *Arabidopsis thaliana* (Niu *et al.* 2011), *Sophora alopecuroides* (Zhao 2011) and *Allium ascalonicum* (Aziz *et al.* 2012), belonging to the families Brassicaceae, Fabaceae and Lilliaaceae respectively. Different strains of *B. cereus* have been established as facultative mosquito pathogens (Krattiger, 1997; Wirth *et al.* 2004; Teng *et al.* 2005, Chatterjee *et al.* 2008) which can colonize in mosquito larval guts in relation to the control of *Aedes aegypti* and *A. subpictus* larvae. But its growth promoting function on sandalwood plant has not known to date. Present study clearly established the strong association of *B. cereus* SW1 with sandalwood plant *Santalum album* L.

Conclusion:

Present study is an attempt to isolate and characterize the rhizospheric soil bacteria from economically important plant *Santalum album* L. It has already been reported that different strains of *Bacillus* act as plant growth promoters for some economically important plants. Further studies may elucidate the positive role of the symbiotic association of *Bacillus cereus* (SW1) with the root of *S. album* as a key factor for the better growth and development of this economically important plant occurring in Bankura district, West Bengal, India.

173

174 **Table 1. Population density (cfu/g dry soil) of different microbial groups in the rhizosphere**
 175 **of *Santalum album* L occurring at different localities of Bankura district**

Sl.no.	Types of organisms	Hirbandh	Basudevpur	Bagaldhara	Maity bandh
1.	Aerobic heterotrophic bacteria ($\times 10^7$)	3.3 \pm 0.001	2.9 \pm 0.001	3.1 \pm 0.001	3.0 \pm 0.001
2.	Gram (-) bacteria ($\times 10^6$)	3.9 \pm 0.001	3.5 \pm 0.002	3.7 \pm 0.001	3.3 \pm 0.001
3.	Nitrifying bacteria ($\times 10^6$)	5.7 \pm 0.003	5.5 \pm 0.013	5.2 \pm 0.002	5.5 \pm 0.013
4.	Phosphate solubilising bacteria ($\times 10^2$)	8.3 \pm 0.014	7.9 \pm 0.018	8.1 \pm 0.011	8.0 \pm 0.020
5.	Starch hydrolyzing bacteria ($\times 10^4$)	10 \pm 0.02	8 \pm 0.02	9.1 \pm 0.018	8.5 \pm 0.021
6.	Spore forming bacteria ($\times 10^4$)	8 \pm 0.012	7.5 \pm 0.015	7.8 \pm 0.008	7.6 \pm 0.018
7.	Fungi ($\times 10^4$)	4.66 \pm 0.002	4.1 \pm 0.003	4.5 \pm 0.001	4.2 \pm 0.011

176 Results are mean of three replication \pm SE, cfu: colony-forming unit

177 **Table 2. Physicochemical properties of rhizospheric soil**

Sl. no.	Area	N (kg/Acre)	P (kg/Acre)	K (kg/Acre)	pH	Organic carbon(%)	Texture
1	Hirbundh	94.67 \pm 0.89	30 \pm 0.03	6.9 \pm 0.013	6.5 \pm 0.019	0.44 \pm 0.001	Lateritic, hard rocky
2	Basudevpur	72.43 \pm 0.40	25 \pm 0.01	6.35 \pm 0.011	6.2 \pm 0.001	0.37 \pm 0.001	Lateritic, hard rocky
3	Bagaldhara	82 \pm 0.71	50 \pm 0.06	7.5 \pm 0.02	6.4 \pm 0.017	0.51 \pm 0.002	Lateritic, hard rocky
4	Maity bundh	76 \pm 0.46	40 \pm 0.04	7.1 \pm 0.017	6.3 \pm 0.013	0.47 \pm 0.001	Lateritic, hard rocky

178 Results are mean of three replication \pm SE

179

180 **Table 3. Biochemical properties of the bacterial isolate (SW1)**

Name of the tests	Observations/ Result
Colony characters	Off white, spherical, elevated
Bacterial properties	rod shaped, >1 µm, motile
Biochemical properties	
Citrate utilization	- ve
Nitrate reduction	+ ve
Catalase	+ ve
Methyl red	+ ve
Voges-Proskauer	+ ve
Antibiotic sensitivity tests:	
amoxycillin (10 µg/disc)	R
ampicillin (10 µg/disc)	R
polymyxin-B(50 µg/disc)	R
pystatin (100 µg/disc)	R
nalidixic acid (30 µg/disc)	S
doxycycline (30 µg/disc)	S
bacitracin (10 µg/disc)	S
tetracycline (30 µg/disc)	S

Where, S = sensitive; R = Resistant





Plate 1

Plate 2

Fig. 1: Scanning electron micrograph of *Bacillus cereus* SW1 (Plate 1 and 2)




Fig 2. FAME analysis of the bacterial isolate SW1




Fig. 3 : A, C, G ,T content (Mol%) of 16S rDNA gene of bacterial isolate SW1





Fig. 4: Restriction map of 16S rDNA gene sequence of bacterial isolate SW1



200

201 **Fig. 5 : Neighbor-joining tree based on 16S rRNA genes sequences of *Bacillus cereus***
 202 **SW1 (KT626448) strain along with few other 16S rRNA genes retrieved from NCBI.**

References:

Aziz AFZ, Halimi MS, Khairuddin AR, Osumanu HA. Variable responses on early development of shallot (*Allium ascalonicum*) and mustard (*Brassica juncea*) plants to *Bacillus cereus* inoculation. Malaysian J. of Microbiol. 2012;8(1) : 47-50.

Brown AE .Benson`s Microbiological Applications. Laboratory Manual in General Microbiology, Short Version. 10th Edition. The McGraw Hill companies. p. 450; 2007.

Chatterjee SN, Das D, Dangar TK . Efficacy of *Bacillus sphaericus* against *Aedes* (Stegomyia) *aegypti* , *Mansonia indiana* (Edward) and *Culex vishnui* . Entomon .2008;33(3): 181-187.

Chatterjee S N, Chatteraj KK, Banerjee P, De MK, Majumder A .Population dynamics and diversity of soil microbes in some areas of Jalpaiguri district of West Bengal, India. Asian journal of Microbial. Biotech. Env. Sc. 2012; Vol.15, no.(2):2013:287-290.

Chatterjee S N, Mukhopadhyay P and Dangar TK. Population dynamics, diversity and characterization of soil bacteria in some south-eastern regions of the Sundarbans, West Bengal, India. Int J Pharm Bio Sci. 2015; 6(2): (B) 353 – 361.

Dangar T K, Babu Y K, Das J .Population dynamics of soil microbes and diversity of *Bacillus thuringiensis* in agricultural and botanic garden soils of India, African J. Biotech. 2010;9 :496-501.

Janssen K. Current protocols in molecular biology, Vol. I. New York: Greene Publ Assoc Inc and John Wiley Sons Inc; 1994.

Krattiger F .Insect resistant crops: A case study of *Bacillus thuringiensis* (BT) and its transfer to developing countries. ISAAA Briefs. 1997; 2: p. 42.

Lacey LA. Manual of techniques in insect pathology. New York: Academic Press; 1997.

Logan NA, Berge O, Bishop AH, Bussse HJ, de Vos P, Fritz D, Heyndrickx M, Kampfer P, Rabinovitch L, Salkinoja-Salonen MS, Seldin L, Ventosa A. Proposed minimal standards for describing new taxa of aerobic, endospore forming bacteria. *Int J Syst Evol Microbiol.*2009;59, 2114-2121.

Niu DD, Liu HX, Jiang CH, Wang YP, Wang QY, Jin HL, Guo JH. The Plant Growth-Promoting Rhizobacterium *Bacillus cereus* AR156 Induces Systemic resistance in *Arabidopsis thaliana* by Simultaneously Activating Salicylate- and Jasmonate/Ethylene-Dependent Signaling Pathways. *Molecular Plant-Microbe Interactions, The American Phytopathol. Soc.*2011;24(5): 533–542.

Merlin, MD, Thomson L, and Elevitch CR. *Santalum ellipticum*, *S. freycinetianum*, *S. haleakalae*, and *S. paniculatum* (Hawaiian sandalwood) Santalaceae (sandalwood family). Permanent Agriculture Resources Holualoa, Hawaii. 21 pages;2006.

Nakade Dhanraj B. Bacterial Diversity in Sugarcane (*Saccharum Officinarum*) Rhizosphere of Saline Soil. *Int. Research J. Biol. Sciences* .2013; 2(2) :0-64.

Pelczar MJ, Bard RC, Burnett GW, Conn HJ, Demoss RD, Euans EE, Weiss FA, Jennison MW, Meckee AP, Riker AJ, Warren J, Weeks OB. *Manual of Microbiological Methods*. McGraw Hill Book Company, Inc., *New York* :315; 1957.

Rao Rama. Host plant of the sandal tree, *Indian Forest records*. 1911; 2(4):159-207.

Rao MN, Soneji JR and Sudarshana P. *Santalum* Wild Crop Relatives: Genomic and Breeding Resources, *Forest Trees* Kole C (ed.), 2011; p. 131-144.

Rawat S, Izhari A, Khan A. Bacterial Diversity in Wheat Rhizosphere and their Characterization. *Advances in Applied Science Res.*2011; (2): 351-356.

Teng HJ, Lu LC, Wu YL, Fang JG .Evaluation of various control agents against mosquito larvae in rice paddies in Taiwan. J. Vect. Ecol. 2005;30: 126-132

Wirth MC, Delecluse A, Walton WE .Laboratory selection for resistance to *Bacillus thuringiensis* subsp. *jegathesan* or a component toxin, Cry 11B, in *Culex quinquefasciatus* (Diptera:Culicidae). J. Med. Entomol. 2004; 41(3): 435-441.

Zhao L. Identification and characterization of the endophytic plant growth promoter *Bacillus Cereus* strain mq23 isolated from *Sophora Alopecuroides* root nodules. Braz. J. of Microbiol. 2011;vol.42 no.2 .