

# Hypergeometric Representation of Modified Beta Stancu Operators

**Prerna Maheshwari**

Department of Mathematics  
SRM University, NCR Campus,  
Modinagar (U.P), India  
E-mail: *mprerna\_anand@yahoo.com*

**Rupa Sharma**

Research Scholar, Mewar University  
Chittorgarh (Rajasthan), India  
E-mail: *vsrsys@gmail.com*

**Abstract:** In the present paper, we extend the results of Gupta and Ahmad [8] by applying hypergeometric series. We study local and global properties, Voronovskaja type asymptotic formula and error estimation for modified Beta operators. We also derive some other characteristics of these operators.

**Key words:** Hypergeometric series, Voronovskaja type asymptotic formula, Modified-Beta Stancu operators, Estimation of error, Direct theorem

**AMS Subject Classification:** 41A25, 41A35.

## 1. Introduction

To approximate integrable function on  $[0, \infty)$ , the modified Beta operators given by Gupta and Ahmad [8] are defined as

$$B_n(f, x) = \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) f(t) dt, \quad x \in [0, \infty) \quad (1.1)$$

where

$$b_{n,k}(x) = \frac{(n+k)!}{k!(n-1)!} \frac{x^k}{(1+x)^{n+k+1}}$$

and

$$p_{n,k}(t) = \frac{(n+k-1)!}{k!(n-1)!} \frac{t^k}{(1+t)^{n+k}}.$$

It was observed in [8], that by taking the modification of Beta operators defined in (1.1), we can have better approximation. Some approximation properties for these operators and other similar type of operators were studied in Agarwal and Singh [2], Abel et al. [1], V. Gupta [5], [6], [7], Zeng et al. [19], Maheshwari [14], [15], Maheshwari-Sharma [16] and Gupta et al. [10], [11] etc. Ismail-Simeonov [13] studied certain Beta type operators in the hypergeometric form and such notations can be studied in [4]. **Some important aspects of approximation theory can be studied in the book written by Gupta - Agarwal [12].** Using Pochhammer symbol  $(n)_k$  defined as  $(n)_k = n(n+1)(n+2)(n+3)\dots(n+k-1)$ , we can write operators (1.1) as

$$\begin{aligned} B_n(f, x) &= \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} \frac{(n)(n+1)_k}{k!} \frac{x^k(1+x)^{-1}}{(1+x)^{n+k}} \int_0^{\infty} \frac{(n)_k}{k!} \frac{t^k}{(1+t)^{n+k}} f(t) dt \\ &= (n-1) \int_0^{\infty} \frac{f(t)(1+x)^{-1}}{[(1+x)(1+t)]^n} \sum_{k=0}^{\infty} \frac{(n)_k(n+1)_k (xt)^k}{(k!)^2 [(1+x)(1+t)]^k} dt. \end{aligned}$$

Using the hypergeometric series  ${}_2F_1(a, b; c; x) = \sum_{k=0}^{\infty} \frac{(a)_k(b)_k}{(c)_k k!} x^k$  and the equality  $(1)_k = k!$ , we have

$$B_n(f, x) = (n-1) \int_0^{\infty} \frac{f(t)(1+x)^{-1}}{[(1+x)(1+t)]^n} {}_2F_1\left(n, n+1; 1; \frac{xt}{(1+x)(1+t)}\right) dt.$$

Now using  ${}_2F_1(a, b; c; x) = {}_2F_1(b, a; c; x)$  and applying Pfaff-Kummer transformation,

$${}_2F_1(a, b; c; x) = (1-x)^{-a} {}_2F_1(a, c-b; c; \frac{x}{x-1}),$$

we have

$$B_n(f, x) = (n-1) \int_0^{\infty} \frac{f(t)(1+x)^{-1}}{(1+x+t)^n} {}_2F_1(n, -n; 1; \frac{-xt}{1+x+t}) dt. \quad (1.2)$$

which is another form of the operator (1.1) in terms of hypergeometric functions.

The Stancu type generalization of Bernstein operators was given in [18].

For  $0 \leq \alpha \leq \beta$ , we propose the Stancu type generalization of **modified-Beta operators**,

$$B_{n,\alpha,\beta}(f, x) = (n-1) \int_0^\infty f\left(\frac{nt+\alpha}{n+\beta}\right) \frac{(1+x)^{-1}}{(1+x+t)^n} {}_2F_1\left(n, -n; 1; \frac{-xt}{1+x+t}\right) dt. \quad (1.3)$$

As a special case if  $\alpha = \beta = 0$ , the operators (1.3) reduce to modified Beta operators defined in (1.1). Let us consider

$$C_\gamma[0, \infty) = \{f \in C[0, \infty) : f(t) = O(t^\gamma), \gamma > 0\}.$$

The operators  $B_{n,\alpha,\beta}(f, x)$  are well defined for  $f \in C_\gamma[0, \infty)$ . In the year 2009, Maheshwari and Gupta [17] studied rate of approximation and weighted approximation of modified Beta operators. Gupta et al. [9] introduced some direct results in simultaneous approximation for Baskakov-Durrmeyer-Stancu operators and Finta [3] obtained a new type of Baskakov-Durrmeyer operators by taking the weight function of Beta operators. In the present paper, we establish moments of modified Beta Stancu operators using the technique of hypergeometric series. Next, we study Voronovskaja type **asymptotic** formula and an estimation of error in simultaneous approximation for the MBS operators.

## 2. Moment estimation and auxiliary results

In this section, we establish certain lemmas which will be useful for the proof of our main theorems.

**Lemma 1.** For  $n \in \mathbb{N}$  and  $r > -1$ , we have

$$B_n(t^r, x) = (n-1) \frac{\Gamma(n-r-1)\Gamma(r+1)}{\Gamma(n)} (1+x)^r {}_2F_1\left(-n, -r; 1; \frac{x}{1+x}\right). \quad (2.1)$$

Moreover

$$B_n(t^r, x) = \frac{(n-r-2)!(n+r)!}{n!(n-2)!} x^r + r^2 \frac{(n-r-2)!(n+r-1)!}{n!(n-2)!} x^{r-1} + O(n^{-2}) \quad (2.2)$$

**Proof:** Taking  $f(t) = t^r$  and  $t = (1+x)u$ , by using Pfaff-Kummer transformation in (1.2), we have

$$\begin{aligned} B_n(t^r, x) &= (n-1) \int_0^\infty \frac{(1+x)^r u^r}{[(1+x)(1+u)]^n} \sum_{k=0}^\infty \frac{(n)_k (-n)_k}{(k!)^2} \frac{(-x(1+x)u)^k}{[(1+x)(1+u)]^k} du \\ &= (n-1) \sum_{k=0}^\infty \frac{(n)_k (-n)_k}{(k!)^2} (-x)^k (1+x)^{r-n} \int_0^\infty \frac{u^{r+k}}{(1+u)^{n+k}} du \\ &= (n-1) \sum_{k=0}^\infty \frac{(n)_k (-n)_k}{(k!)^2} (-x)^k (1+x)^{r-n} B(r+k+1, n-r-1) \\ &= (n-1) \sum_{k=0}^\infty \frac{(n)_k (-n)_k}{(k!)^2} (-x)^k (1+x)^{r-n} \frac{\Gamma(r+k+1)\Gamma(n-r-1)}{\Gamma(n+k)}. \end{aligned}$$

Again taking  $\Gamma(n+k+1) = \Gamma(n+1)(n+1)_k$ , we get

$$\begin{aligned} B_n(t^r, x) &= (n-1) \sum_{k=0}^\infty \frac{(n)_k (-n)_k}{(k!)^2} (-x)^k (1+x)^{r-n} \frac{\Gamma(r+1)(r+1)_k \Gamma(n-r-1)}{\Gamma(n)(n)_k} \\ &= (n-1)(1+x)^{r-n} \frac{\Gamma(r+1)\Gamma(n-r-1)}{\Gamma(n)} \sum_{k=0}^\infty \frac{(-n)_k (r+1)_k}{(k!)^2} (-x)^k \\ &= (n-1)(1+x)^{r-n} \frac{\Gamma(r+1)\Gamma(n-r-1)}{\Gamma(n)} {}_2F_1(-n, r+1; 1; -x). \end{aligned}$$

Using Pfaff-Kummer transformation, we get

$$B_n(t^r, x) = (n-1)(1+x)^r \frac{\Gamma(r+1)\Gamma(n-r-1)}{\Gamma(n)} {}_2F_1\left(-n, -r; 1; \frac{x}{x-1}\right).$$

$$\begin{aligned}
B_n(t^r, x) &= (n-1)(1+x)^r \frac{r!(n-r-2)!}{(n-1)!} {}_2F_1\left(-r, 1+n; 1; \frac{x}{x-1}\right) \\
&= \frac{r!(n-r-2)!}{(n-2)!} \sum_{k=0}^{\infty} \frac{(-r)_k (n+1)_k}{(k!)^2} (-x)^k \\
&= \frac{(r!)^2 (n-r-2)!}{n!(n-2)!} \sum_{k=0}^{\infty} \frac{(n+k)!}{(r-k)!(k!)^2} (x)^k \\
B_n(t^r, x) &= \frac{(n-r-2)!(n+r)!}{n!(n-2)!} x^r + r^2 \frac{(n-r-2)!(n+r-1)!}{n!(n-2)!} x^{r-1} + O(n^{-2})
\end{aligned}$$

**Lemma 2.** For  $0 \leq \alpha \leq \beta$ , we have

$$\begin{aligned}
B_{n,\alpha,\beta}(t^r, x) &= x^r \frac{n^r}{(n+\beta)^r} \frac{(n+r)!(n-r-2)!}{n!(n-2)!} \\
&+ x^{r-1} \left\{ r^2 \frac{n^r}{(n+\beta)^r} \frac{(n+r-1)!(n-r-2)!}{n!(n-2)!} + r\alpha \frac{n^{r-1}}{(n+\beta)^r} \frac{(n+r-1)!(n-r-1)!}{n!(n-2)!} \right\} \\
&+ x^{r-2} \left\{ r(r-1)^2 \alpha \frac{n^{r-1}}{(n+\beta)^r} \frac{(n+r-2)!(n-r-1)!}{n!(n-2)!} + \frac{r(r-1)\alpha^2}{2} \frac{n^{r-2}}{(n+\beta)^r} \frac{(n+r-2)!(n-r)!}{n!(n-2)!} \right\} \\
&+ O(n^{-2}).
\end{aligned}$$

**Proof:** By using Binomial theorem, the relation between operators (1.2) and (1.3) can be defined as

$$\begin{aligned}
B_{n,\alpha,\beta}(t^r, x) &= (n-1) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left( \frac{nt+\alpha}{n+\beta} \right)^r dt \\
&= (n-1) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \sum_{j=0}^r \binom{r}{j} \frac{(nt)^j \alpha^{r-j}}{(n+\beta)^r} dt \\
&= \sum_{j=0}^r \binom{r}{j} \frac{n^j \alpha^{r-j}}{(n+\beta)^r} B_n(t^j, x).
\end{aligned}$$

Applying this on (2.2), we get the required result.

**Lemma 3.** [8] For  $m \in N \cup \{0\}$ , if

$$T_{n,m}(x) = \frac{1}{n} \sum_{k=0}^{\infty} b_{n,k}(x) \left( \frac{k}{n+1} - x \right)^m.$$

then there exists the following recurrence relation

$$(n+1)T_{n,m+1}(x) = x(1+x)[T'_{n,m}(x) + mT_{n,m-1}(x)].$$

Consequently,

- (1)  $T_{n,m}(x)$  are polynomials in  $x$  of degree at most  $m$ .
- (2)  $T_{n,m}(x) = O(n^{-[(m+1)/2]})$ , where  $[\alpha]$  denotes the integral part of  $\alpha$ .

**Lemma 4.** We define the central moments as

$$T_{n,m}(x) = B_{n,\alpha,\beta}((t-x)^m, x) = \left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left( \frac{nt+\alpha}{n+\beta} - x \right)^m dt, \quad m \in N \setminus \{0\},$$

We have the following recurrence relation, for  $n > m+1$ .

$$\begin{aligned}
(n-m-2) \left( \frac{n+\beta}{n} \right) T_{n,m+1}(x) &= x(1+x)[T'_{n,m}(x) + mT_{n,m-1}(x)] \\
&+ \left[ (m+nx+1+x) + \left( \frac{n+\beta}{n} \right) \left( \frac{\alpha}{n+\beta} - x \right) (n-2m-2) \right] T_{n,m}(x) \\
&- \left( \frac{\alpha}{n+\beta} - x \right) \left[ 1 - \left( \frac{\alpha}{n+\beta} - x \right) \left( \frac{n+\beta}{n} \right) \right] mT_{n,m-1}(x).
\end{aligned}$$

We can obtain first three moments as  $T_{n,0}(x) = 1$ ,  $T_{n,1}(x) = \frac{x[3n+\beta(2-n)+n+\alpha(n-2)]}{(n+\beta)(n-2)}$ ,

$$\begin{aligned} T_{n,2}(x) &= \frac{x^2[2n^2(n+7) - n^2\beta(7-n) + n\beta(n\beta - 5\beta + 18 + 6\beta^2)]}{(n+\beta)^2(n-2)(n-3)} \\ &+ \frac{x[2n^2(n+5) + 2n^2\alpha(5-2n) - 2n^2\beta(1+\alpha) + 2n(5\alpha\beta + 3\beta - 9\alpha) - 12\alpha\beta]}{(n+\beta)^2(n-2)(n-3)} \\ &+ \frac{2n^2 + n^2\alpha(\alpha+2) - n\alpha(5\alpha+6) + 6\alpha^2}{(n+\beta)^2(n-2)(n-3)}. \end{aligned}$$

From the recurrence relation, it is easily verified that for all  $x \in [0, \infty)$ , we have

$$T_{n,m}(x) = O(n^{-[(m+1)/2]}).$$

**Proof:**  $T_{n,0}(x) = 1$  (by the definition of the operators (1.3)). For the proof of other moments we follow the recurrence relation. Now we prove the recurrence relation as follows.

Using the identities

$$\begin{aligned} x(1+x)b'_{n,k}(x) &= [k - (n+1)x]b_{n,k}(x) \\ t(1+t)p'_{n,k}(t) &= (k - nt)p_{n,k}(t), \end{aligned}$$

We have

$$x(1+x)T'_{n,m}(x) = \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} [k - (n+1)x]b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^m dt - mx(1+x)T_{n,m-1}(x).$$

Thus

$$\begin{aligned} &x(1+x)[T'_{n,m}(x) + mT_{n,m-1}(x)] \\ &= \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} (k - nx)b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^m dt \\ &\quad - \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} xb_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^m dt \\ &= \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} [(k - nt) + n(t - x)]p_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^m dt - xT_{n,m}(x) \\ &= \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} t(1+t)p'_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^m dt \\ &\quad + \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} n(t - x)p_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^m dt - xT_{n,m}(x). \end{aligned}$$

Now writing

$$t = \left(\frac{n+\beta}{n}\right) \left[\frac{nt+\alpha}{n+\beta} - x - \left(\frac{\alpha}{n+\beta} - x\right)\right],$$

We have

$$\begin{aligned} x(1+x)[T'_{n,m}(x) + mT_{n,m-1}(x)] &= \left(\frac{n+\beta}{n}\right) \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p'_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^{m+1} dt \\ &\quad - \left(\frac{n+\beta}{n}\right) \left(\frac{\alpha}{n+\beta} - x\right) \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p'_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^m dt \\ &\quad + \left(\frac{n+\beta}{n}\right)^2 \left[\left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p'_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^{m+2} dt \right. \\ &\quad \left. + \left(\frac{\alpha}{n+\beta} - x\right)^2 \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p'_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^m dt \right. \\ &\quad \left. - 2\left(\frac{\alpha}{n+\beta} - x\right) \left(\frac{n-1}{n}\right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p'_{n,k}(t) \left(\frac{nt+\alpha}{n+\beta} - x\right)^{m+1} dt \right] \\ &\quad + n\left(\frac{n+\beta}{n}\right) T_{n,m+1}(x) - n\left(\frac{\alpha}{n+\beta} - x\right) \left(\frac{n+\beta}{n}\right) T_{n,m}(x) \\ &\quad - nxT_{n,m}(x) - xT_{n,m}(x). \end{aligned}$$

Integrating by parts and by simple computation, we get

$$\begin{aligned} (n-m-2)\frac{n+\beta}{n}T_{n,m+1}(x) &= x(1+x)[T'_{n,m}(x) + mT_{n,m-1}(x)] \\ &+ \left[ (m+nx+1+x) + \frac{n+\beta}{n} \left( \frac{\alpha}{n+\beta} - x \right) (n-2m-2) \right] T_{n,m}(x) \\ &- \left( \frac{\alpha}{n+\beta} - x \right) \left[ 1 - \left( \frac{\alpha}{n+\beta} - x \right) \left( \frac{n+\beta}{n} \right) \right] mT_{n,m-1}(x). \end{aligned}$$

**Lemma 5.** [10] *There exist the polynomials  $q_{i,j,r}(x)$  on  $[0, \infty)$ , independent of  $n$  and  $k$  such that*

$$(x(1+x))^r \frac{d^r}{dx^r} b_{n,k}(x) = \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^i (k - (n+1)x)^j q_{i,j,r}(x) b_{n,k}(x).$$

### 3. Direct Estimates

In this section, we propose some direct results including asymptotic formula and an error estimation in simultaneous approximation.

**Theorem 1.** *Let  $f \in C_\gamma[0, \infty)$  be bounded on every finite sub-interval  $[0, \infty)$  admitting the derivative of order  $(r+2)$  at a fixed  $x \in [0, \infty)$ . Let  $f(t) = O(t^\gamma)$  as  $t \rightarrow \infty$  for some  $\gamma > 0$ , then we have*

$$\lim_{n \rightarrow \infty} n \left( B_{n,\alpha,\beta}^{(r)}(f, x) - f^{(r)}(x) \right) = r(r+2-\beta)f^{(r)}(x) + [(1+r+\alpha) + x(2r+3-\beta)]f^{(r+1)}(x) + x(1+x)f^{(r+2)}(x),$$

**Proof:** By Taylor's expansion of  $f$ , we have

$$f(t) = \sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} (t-x)^i + \varepsilon(t, x)(t-x)^{r+2},$$

where  $\varepsilon(t, x) \rightarrow 0$  as  $t \rightarrow x$  and  $\varepsilon(t, x) = O((t-x)^\delta)$  as  $t \rightarrow \infty$  for some  $\delta > 0$ , therefore we can write,

$$n[B_{n,\alpha,\beta}^{(r)}(f, x) - f^{(r)}(x)] = n \left[ \sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} B_{n,\alpha,\beta}^{(r)}((t-x)^i, x) - f^{(r)}(x) \right] + nB_{n,\alpha,\beta}^{(r)}(\varepsilon(t, x)(t-x)^{r+2}, x) =: E_1 + E_2$$

From Lemma 2, we have

$$\begin{aligned} E_1 &= n \sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} \sum_{j=r}^i \binom{i}{j} (-x)^{i-j} B_{n,\alpha,\beta}^{(r)}(t^j, x) - n f^{(r)}(x) = \frac{f^{(r)}(x)}{r!} n \left( B_{n,\alpha,\beta}^{(r)}(t^r, x) - r! \right) \\ &+ \frac{f^{(r+1)}(x)}{(r+1)!} n \left\{ (r+1)(-x) B_{n,\alpha,\beta}^{(r)}(t^r, x) + B_{n,\alpha,\beta}^{(r)}(t^{r+1}, x) \right\} \\ &+ \frac{f^{(r+2)}(x)}{(r+2)!} n \left\{ \frac{(r+2)(r+1)}{2} x^2 B_{n,\alpha,\beta}^{(r)}(t^r, x) + (r+2)(-x) B_{n,\alpha,\beta}^{(r)}(t^{r+1}, x) + B_{n,\alpha,\beta}^{(r)}(t^{r+2}, x) \right\} \\ &= n \left[ \frac{n^r(n+r)!(n-r-2)!}{(n+\beta)^r n!(n-2)!} - 1 \right] f^{(r)}(x) + n \frac{f^{(r+1)}(x)}{(r+1)!} \left\{ (r+1)(-x) \frac{n^r(n+r)!(n-r-2)!}{(n+\beta)^r n!(n-2)!} r! \right. \\ &+ \frac{n^{r+1}(n+r+1)!(n-r-3)!}{(n+\beta)^{r+1} n!(n-2)!} (r+1)! x + \frac{(r+1)^2 n^{r+1}(n+r)!(n-r-3)!}{(n+\beta)^{r+1} n!(n-2)!} r! \\ &+ (r+1)\alpha \frac{n^r(n+r)!(n-r-2)!}{(n+\beta)^{r+1} n!(n-2)!} r! \left. \right\} + n \frac{f^{(r+2)}(x)}{(r+2)!} \left\{ \frac{(r+2)(r+1)}{2} x^2 \frac{n^r(n+r)!(n-r-2)!}{(n+\beta)^r n!(n-2)!} r! \right. \\ &- (r+2)x \left( \frac{n^{r+1}(n+r+1)!(n-r-3)!}{(n+\beta)^{r+1} n!(n-2)!} (r+1)! x + \frac{(r+1)^2 n^{r+1}(n+r)!(n-r-3)!}{(n+\beta)^{r+1} n!(n-2)!} r! \right. \\ &+ (r+1)\alpha \frac{n^r(n+r)!(n-r-3)!}{(n+\beta)^{r+1} n!(n-2)!} r! \left. \right\} + \frac{n^{r+2}(n+r+2)!(n-r-4)!}{(n+\beta)^{r+2} n!(n-2)!} \frac{(r+2)!}{2} x^2 \\ &+ \frac{(r+2)^2 n^{r+2}(n+r+1)!(n-r-4)!}{(n+\beta)^{r+2} n!(n-2)!} (r+1)! x + (r+2)\alpha \frac{n^{r+1}(n+r+1)!(n-r-3)!}{(n+\beta)^{r+2} n!(n-2)!} (r+1)! x \\ &+ \frac{(r+2)(r+1)^2 n^{r+1}\alpha(n+r)!(n-r-3)!}{(n+\beta)^{r+2} n!(n-2)!} r! + \frac{(r+2)(r+1)\alpha^2 n^r(n+r)!(n-r-2)!}{2(n+\beta)^{r+2} n!(n-2)!} r! \left. \right\} + O(n^{-2}). \end{aligned}$$

The coefficients of  $f^{(r)}(x)$ ,  $f^{(r+1)}(x)$ ,  $f^{(r+2)}(x)$  in the above expression are respectively  $r(r+2-\beta)$ ,  $(1+r+\alpha)+x(2r+3-\beta)$  and  $x(1+x)$ . Taking the limits as  $n \rightarrow \infty$  and by using induction hypothesis on  $r$ . To complete the proof of the theorem, it will be enough to show that  $E_2 \rightarrow 0$  as  $n \rightarrow \infty$ . To estimate  $E_2$  using Lemma 5, we have

$$|E_2| \leq (n-1) \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^i \frac{|q_{i,j,r}(x)|}{x^r(1+x)^r} \sum_{k=0}^{\infty} b_{n,k}(x) |k - (n+1)x|^j \int_0^{\infty} p_{n,k}(t) |\varepsilon(t, x)| \left| \frac{nt+\alpha}{n+\beta} - x \right|^{r+2} dt.$$

For a given  $\varepsilon > 0$ , there exist  $\delta > 0$  such that  $|\varepsilon(t, x)| < \varepsilon$  whenever  $|t - x| < \delta$ . For  $|t - x| \geq \delta$ , since  $\varepsilon(t, x) \rightarrow 0$  as  $t \rightarrow x$ , further if  $\lambda$  is any integer  $\lambda > \max(\gamma, r+2)$ , then we find a constant  $K > 0$  such that  $|\varepsilon(t, x)| \left| \frac{nt+\alpha}{n+\beta} - x \right|^{r+2} \leq K \left| \frac{nt+\alpha}{n+\beta} - x \right|^\lambda$ . Thus

$$|E_2| \leq (n-1)A_1 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^{i+1} \sum_{k=0}^{\infty} b_{n,k}(x) |k - (n+1)x|^j \left\{ \int_{|t-x| < \delta} \varepsilon p_{n,k}(t) \left| \frac{nt+\alpha}{n+\beta} - x \right|^{r+2} dt \right. \\ \left. + \int_{|t-x| \geq \delta} K p_{n,k}(t) \left| \frac{nt+\alpha}{n+\beta} - x \right|^\lambda dt \right\} =: E_3 + E_4.$$

Now applying Schwarz inequality for the integration and summation, we have

$$|E_3| \leq (n-1)\varepsilon A_1 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^{i+1} \sum_{k=0}^{\infty} b_{n,k}(x) |k - (n+1)x|^j \\ \times \left( \int_0^{\infty} p_{n,k}(t) dt \right)^{\frac{1}{2}} \left( \int_0^{\infty} p_{n,k}(t) \left( \frac{nt+\alpha}{n+\beta} - x \right)^{2r+4} dt \right)^{\frac{1}{2}} \\ \leq (n-1)\varepsilon A_1 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^{i+1} \left( \sum_{k=0}^{\infty} b_{n,k}(x) |k - (n+1)x|^{2j} \right)^{\frac{1}{2}} \\ \times \left( \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left( \frac{nt+\alpha}{n+\beta} - x \right)^{2r+4} dt \right)^{\frac{1}{2}}.$$

Using Lemma 3 and Lemma 4, we get

$$|E_3| \leq (n-1)\varepsilon A_1 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^{i+1} \cdot O(n^{j/2}) \cdot O(n^{-(r+2)/2}) \leq \varepsilon O(1).$$

For arbitrary chosen  $\varepsilon$ , we follow that  $E_3 = o(1)$ . Now using Schwarz inequality for the integration and summation, Lemma 3 and Lemma 4, we get

$$|E_4| \leq (n-1)A_2 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^{i+1} \sum_{k=0}^{\infty} b_{n,k}(x) |k - (n+1)x|^j \int_{|t-x| \geq \delta} p_{n,k}(t) \left| \frac{nt+\alpha}{n+\beta} - x \right|^\lambda dt \\ \leq (n-1)A_2 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^{i+1} \left( \sum_{k=0}^{\infty} b_{n,k}(x) (k - (n+1)x)^{2j} \right)^{1/2} \\ \times \left( \sum_{k=1}^{\infty} b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left( \frac{nt+\alpha}{n+\beta} - x \right)^{2\lambda} dt \right)^{1/2} \\ = (n-1) \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^{i+1} \cdot O(n^{j/2}) \cdot O(n^{-\lambda/2}) = O(n^{(r+2-\lambda)/2}) = o(1).$$

Hence we get  $E_2 \rightarrow 0$  as  $n \rightarrow \infty$ . Combining the estimates of  $E_1$  and  $E_2$ , we get the required results. This completes the proof of theorem.

**Theorem 2.** Let  $f \in C_\gamma[0, \infty)$  for some  $\gamma > 0$  and  $r \leq m \leq r+2$ . If  $f^{(m)}$  exist and is continuous on  $(a - \eta, b + \eta) \subset [0, \infty)$ ,  $\eta > 0$ , then for  $n$  sufficiently large

$$\left\| B_{n,\alpha,\beta}^{(r)}(f, x) - f^{(r)}(x) \right\|_{c[a,b]} \leq A_1 n^{-1} \sum_{i=r}^m \left\| f^{(i)} \right\|_{c[a,b]} + A_2 n^{-1/2} \omega(f^{(m)}, n^{-1/2}) + O(n^{-2}),$$

where  $A_1, A_2$  are constants which do not depend on  $f$  and  $n$  and  $\omega(f, \delta)$  is the modulus of continuity of  $f$  on  $(a - \eta, b + \eta)$  and  $\|\cdot\|_{c[a,b]}$  being the sup-norm on  $[a, b]$ .

**Proof:** Using Taylor's expansion of  $f$ , we have

$$f(t) = \sum_{i=0}^m \frac{f^{(i)}(x)}{i!} (t-x)^i + \frac{f^{(m)}(\xi) - f^{(m)}(x)}{m!} (t-x)^m \chi(t) + h(t, x)(1 - \chi(t)),$$

where  $\xi$  lies between  $t$  and  $x$  and  $\chi(t)$  is the characteristic function on the interval  $(a - \eta, b + \eta)$ .  
Now

$$\begin{aligned} B_{n,\alpha,\beta}^{(r)}(f, x) - f^{(r)}(x) &= \left\{ \sum_{i=0}^m \frac{f^{(i)}(x)}{i!} B_{n,\alpha,\beta}^{(r)}((t-x)^i) - f^{(r)}(x) \right\} \\ &\quad + B_{n,\alpha,\beta}^{(r)} \left( \frac{f^{(m)}(\xi) - f^{(m)}(x)}{m!} (t-x)^m \chi(t), x \right) \\ &\quad + B_{n,\alpha,\beta}^{(r)}(h(t, x)(1 - \chi(t)), x) =: J_1 + J_2 + J_3. \end{aligned}$$

By using Lemma 2, we have

$$\begin{aligned} J_1 &= \sum_{i=0}^m \frac{f^{(i)}(x)}{i!} \sum_{j=0}^i \binom{i}{j} (-x)^{i-j} \frac{d^r}{dx^r} \left[ x^j \frac{n^j}{(n+\beta)^j} \frac{(n+j)!(n-j-2)!}{n!(n-2)!} \right. \\ &\quad \left. + x^{j-1} (j^2 \frac{n^j}{(n+\beta)^j} \frac{(n+j-1)!(n-j-2)!}{n!(n-2)!} \right. \\ &\quad \left. + j\alpha \frac{n^{j-1}}{(n+\beta)^j} \frac{(n+j-1)!(n-j-1)!}{n!(n-2)!} \right) \\ &\quad \left. + x^{j-2} \{ j(j-1)^2 \alpha \frac{n^{j-1}}{(n+\beta)^j} \frac{(n+j-2)!(n-j-1)!}{n!(n-2)!} \right. \\ &\quad \left. + \frac{j(j-1)\alpha^2}{2} \frac{n^{j-2}}{(n+\beta)^j} \frac{(n+j-2)!(n-j)!}{n!(n-2)!} \} + O(n^{-2}) \right] - f^{(r)}(x). \end{aligned}$$

Therefore,  $\|J_1\|_{c[a,b]} \leq A_1 n^{-1} \sum_{i=r}^m \|f^{(i)}\|_{c[a,b]} + O(n^{-2})$ , uniformly on  $[a, b]$ .

Next, we estimate  $J_2$  as

$$\begin{aligned} \|J_2\| &\leq \left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} |b_{n,k}^r(x)| \int_0^{\infty} p_{n,k}(t) \left\{ \left| \frac{f^{(m)}(\xi) - f^{(m)}(x)}{m!} \right| \left| \frac{nt+\alpha}{n+\beta} - x \right|^m \chi(t) \right\} dt \\ &\leq \frac{\omega(f^{(m)}, \delta)}{m!} \left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} |b_{n,k}^r(x)| \int_0^{\infty} p_{n,k}(t) \left( 1 + \frac{\left| \frac{nt+\alpha}{n+\beta} - x \right|^m}{\delta} \right) \left| \frac{nt+\alpha}{n+\beta} - x \right|^m dt \\ &\leq \frac{\omega(f^{(m)}, \delta)}{m!} \left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} |b_{n,k}^r(x)| \int_0^{\infty} p_{n,k}(t) \left( \left| \frac{nt+\alpha}{n+\beta} - x \right|^m + \delta^{-1} \left| \frac{nt+\alpha}{n+\beta} - x \right|^{m+1} \right) dt. \end{aligned}$$

Using Schwarz inequality for integration and summation, we get

$$\begin{aligned} &\left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} b_{n,k}(x) |k - (n+1)x|^j \int_0^{\infty} p_{n,k}(t) \left| \frac{nt+\alpha}{n+\beta} - x \right|^m dt \\ &\leq \left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} b_{n,k}(x) |k - (n+1)x|^j \left( \int_0^{\infty} p_{n,k}(t) dt \right)^{1/2} \left( \int_0^{\infty} p_{n,k}(t) \left( \frac{nt+\alpha}{n+\beta} - x \right)^{2m} dt \right)^{1/2} \\ &\leq (n+1)^j \left( \frac{1}{n} \sum_{k=0}^{\infty} b_{n,k}(x) \left| \frac{k}{n+1} - x \right|^{2j} \right)^{1/2} \left( \left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left( \frac{nt+\alpha}{n+\beta} - x \right)^{2m} dt \right)^{1/2} \\ &= O(n^{j/2}). O(n^{-m/2}) = O(n^{(j-m)/2}), \end{aligned} \tag{3.1}$$

uniformly on  $[a, b]$ .

Therefore by Lemma 5 and (3.1), we get

$$\begin{aligned}
& \left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} |b_{n,k}(x)| \int_0^{\infty} p_{n,k}(t) \left| \frac{nt+\alpha}{n+\beta} - x \right|^m dt \\
& \leq \left( \frac{n-1}{n} \right) \sum_{k=0}^{\infty} \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^i |k - (n+1)x|^j \frac{|q_{i,j,r}(x)|}{x^r(1+x)^r} b_{n,k}(x) \int_0^{\infty} p_{n,k}(t) \left| \frac{nt+\alpha}{n+\beta} - x \right|^m dt \\
& \leq \left( \sup_{\substack{2i+j \leq r \\ i,j \geq 0}} \frac{|q_{i,j,r}(x)|}{x^r(1+x)^r} \right) \left( \frac{n-1}{n} \right) \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^i \left( \sum_{k=0}^{\infty} b_{n,k}(x) |k - (n+1)x|^j \int_0^{\infty} p_{n,k}(t) \left| \frac{nt+\alpha}{n+\beta} - x \right|^m dt \right) \\
& = M \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^i O(n^{(j-m)/2}) = O(n^{(r-m)/2}). \tag{3.2}
\end{aligned}$$

uniformly on  $[a, b]$ , where

$$M = \sup_{\substack{2i+j \leq r \\ i,j \geq 0}} \sup_{x \in [a,b]} \frac{|q_{i,j,r}(x)|}{x^r(1+x)^r}.$$

Choosing  $\delta = n^{-1/2}$  and applying (3.2), we obtain.

$$\begin{aligned}
\|J_2\|_{c[a,b]} & \leq \frac{\omega(f^{(m)}, n^{-1/2})}{m!} \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^i [O(n^{(j-m)/2}) + n^{-1/2} O(n^{(r-m-1)/2})] + O(n^{-m}) \\
& \leq M_2 n^{-(r-m)/2} \omega(f^{(m)}, n^{-1/2}).
\end{aligned}$$

For  $t \in [0, \infty) \setminus (a - \eta, b + \eta)$ , we can choose  $\delta$  such that  $|t - x| \geq \delta$  for all  $x \in [a, b]$ . Thus by Lemma 5, we get

$$|J_3| \leq M \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^i b_{n,k}(x) |k - (n+1)x|^j \int_0^{\infty} p_{n,k}(t) |h(t, x)| dt.$$

For  $|t - x| \geq \delta$ , we can find a constant  $K$  such that  $|h(t, x)| \leq K \left| \frac{nt+\alpha}{n+\beta} - x \right|^\beta$ , where  $\beta$  is an integer  $\geq \max(\gamma, m)$ . Hence applying the Schwarz inequality for both integration and summation, Lemma 3 and Lemma 4, it is easy to show that  $J_3 = O(n^{-s})$  for any  $s > 0$ , uniformly on  $[a, b]$ . Combining the estimates of  $J_1, J_2, J_3$ , we get the required result.

## REFERENCES

- [1] U. Abel, M. Ivan, Z. Li, *Local approximation by generalized Baskakov-Durrmeyer operators*, Numer. Funct. Anal. Optim. 28 (3-4) (2007), 245-264.
- [2] P. N. Agarwal, K. K. Singh, *Higher order approximation by iterates of modified Beta operators*, Thai J. of Math. 10 (3) (2012), 643-650.
- [3] Z. Finta, *On converse approximation theorems*, J. Math. Anal. Appl. 312 (1) (2005), 159-180.
- [4] G. Gasper, M. Rahman, *Basic hypergeometric series*, Encyclopedia of Applied and Computational Mathematics, Cambridge University Press, Cambridge, UK, 1990.
- [5] V. Gupta, *Rate of convergence for generalized Baskakov Beta operators*, South East Asian Bull Math. 33 (1) (2009), 57-64.
- [6] V. Gupta, *Some approximation properties for modified Baskakov type operators*, Georgian Math. J. 12 (2) (2005), 217-228.
- [7] V. Gupta, *Rate of approximation by new sequence of linear positive operators*, Comput. Math. Appl. 45 (12) (2003), 1895-1904.
- [8] V. Gupta, A. Ahmad, *Simultaneous approximation by modified Beta operators*, Istanbul Univ. Fen Fak. Mat.Der. 54 (1995), 11-22.
- [9] V. Gupta, D. K. Verma, P. N. Agrawal, *Simultaneous approximation by certain Baskakov-Durrmeyer Stancu operators*, J. of the Egyptain Mathe. Soc. 20 (3) (2012), 183-187.
- [10] V. Gupta, P. Maheshwari, V. K. Jain, *Lp inverse theorem for modified Beta operators*, Int. J. Math. and Mathematical Sciences 20 (2003), 1295-1303.
- [11] V. Gupta, P. Maheshwari, V. K. Jain, *On the linear combinations of modified Beta operators*, J. Pure and Appl. Math. Sci. 48 (2003), 63-75.
- [12] V. Gupta, R. P. Agarwal, *Convergence estimate in approximation theory*, XIII (2014), 361 Springer.
- [13] M. Ismail, P. Simeonov, *On a family of positive linear integral operators*, notions of positivity and the geometry of polynomials, Trends in Mathematics, Springer, Basel AG (2011), 259-274.
- [14] P. Maheshwari, *Saturation theorem for the combinations of modified Beta operators in Lp-spaces*, Applied Mathematics E-Notes 7 (2007), 32-37.
- [15] P. Maheshwari, *An inverse result in simultaneous approximation by modified Beta operators*, Georgian Math. J. 13(2) (2006), 297-306.
- [16] P. Maheshwari, D. Sharma, *Approximation by q Baskakov-Beta-stancu operators*, Rend. Circ Mat Palermo 61 (2012), 297-305.

- [17] P. Maheshwari, V. Gupta, *On rate of approximation by modified Beta operators*, Int. J. of Math. and Mathematical Sciences 205649 (2009), 8 pages.
- [18] D. D. Stancu, *Approximation of function by mean of a new generalized Bernstein operators*, Calcolo 20 (1983), 211-229.
- [19] X. M. Zeng, U. Abel, M. Ivan, *Approximation by a Kantorovich variant of the Bleimann, Butzer and Hann operators*, Math. Inequal. Appl. 11 (2) (2008), 317-325.