Combined Solitary Wave Solutions in Higher-order Effects Optical Fibers

Jean Roger Bogning

University of Bamenda, Higher Teachers Training College Bambili, Department of Physics, Pobox 39 Bamenda, Cameroon

.

ABSTRACT

We construct in this manuscript, the combined solitary wave solutions of nonlinear Schrödinger equation that governs the dynamics of propagation of waves in optical fibers with higher-order effects. We base our survey on the sum of two analytic shapes of the solitary waves of bright and dark type to form a resulting solitary wave to determine.

1. INTRODUCTION

The physical systems are generally governed by the partial differential equations that are in most cases nonlinear. When these partial differential equations are linear, they obey the principles of superposition and especially of the uniqueness of the solution. When these partial differential equations are nonlinear, the principle of linear combination of the solutions is not applicable and the approach in the resolution varies according to the type of equation and especially of the type of solutions that one wants to get. Thus, it becomes difficult to speak of standard method of resolution or unique solution. It is exactly for this reason that the multiple methods and different approaches developed by many authors exist [1-16] with the aim of solving the considered equation completely or to find the approached solutions with a minimal error margin. Beyond all these methods of integration of the nonlinear partial differential equations, the research of the forced solutions occupies an important place. To adapt and to exploit the natural phenomena which surround us in order to improve our daily work. Among the searched solutions, the solitary wave probably occupies a very important place because of its futuristic applications. This fact is translated in many articles written by many researchers [17-24]. In this dynamics, we construct some solitary wave solutions of the nonlinear partial differential equation that govern the propagation of waves in the greatly nonlinear effects optical fibers and govern by the equation [25, 26]

$$U_z - i\alpha_1 U_{tt} - i\alpha_2 |U|^2 U - \alpha_3 U_{ttt} - \alpha_4 \left(|U|^2 U\right)_t - \alpha_5 U \left(|U|^2\right)_t = 0, \tag{1}$$

where U is the slowly varying envelope of the electric field, the indexes z and t denote the spatial and temporal partial derivatives, and α_1, α_2, α_3, α_4 and α_5 are real constants related to the group-velocity dispersion, the self-phase modulation, the third-order dispersion, the self-steepening, and the delayed nonlinear response effect, respectively. The solutions that we construct effectively are of type combined solitary wave; that means resulting from a combination of the bright solitary

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr

wave and dark solitary wave. Otherwise to say the solutions that we want to construct are of the shapes

$$U_{lj}(z,t)=\sum_{l,j} a_{lj}\frac{\sinh^{j}(\alpha t)}{\cosh^{l}(\alpha t)}\exp\left[-i(kz-\omega t)\right], \tag{2}$$

where a_{lj}, α are constants, k the spatial frequency of the wave and ω the angular frequency of the wave to be determined as a function of the parameters $\alpha_{\chi}\,(\chi=1,2,\ldots,5)$ of equation (1), $l=1,2,\ldots$; $j=0,1$ and $i^2=-1$. If we define for example from Eq. (2), two solutions as

$$U_1(z,t)=\left(\frac{a}{\cosh(\alpha t)}+b\frac{\sinh(\alpha t)}{\cosh(\alpha t)}\right)\exp\left[-i(kz-\omega t)\right], \tag{3}$$

and

$$U_2(z,t)=\left(\frac{c}{\cosh^2(\alpha t)}+d\frac{\sinh(\alpha t)}{\cosh^2(\alpha t)}\right)\exp\left[-i(kz-\omega t)\right], \tag{4}$$

where the constants a, b, c and d are to be determined, we see that U_1 is constituted by $U_{11}=\left(a/\cosh(\alpha t)\right)\exp\left[-i(kz-\omega t)\right]$ which is a bright solitary wave and $U_{12}=\left(b\sinh(\alpha t)/\cosh(\alpha t)\right)\exp\left[-i(kz-\omega t)\right]$ which is a dark solitary wave. On the other hand U_2 is constituted by $U_{21}=\left(c/\cosh^2(\alpha t)\right)\exp\left[-i(kz-\omega t)\right]$ which is a bright solitary wave and $U_{22}=\left(d\sinh(\alpha t)/\cosh^2(\alpha t)\right)\exp\left[-i(kz-\omega t)\right]$ which is a dark solitary wave.



Fig. 1. Curves of $|U_{11}|$, $|U_{12}|$, $|U_{21}|$ and $|U_{22}|$ as a function of time; (a):the red curve indicates the profile of $|U_{11}|$ for $a=5$ and $\alpha=1$; the green curve the profile of $|U_{12}|$ for $b=5$ and $\alpha=1$; (b): the red curve indicates the profile of $|U_{21}|$ for $c=5$ and $\alpha=1$; the green curve gives the profile of $|U_{22}|$ for $d=5$ and $\alpha=1$;

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr



Fig. 2. Curves of $|U_1|$ and $|U_2|$ as a function of time; (a):the red curve indicates the profile of $|U_1|$ for $a=6$, $b=2$ and $\alpha=1$; the green curve the profile of $|U_1|$ for $a=2$, $b=6$ and $\alpha=1$; (b): the red curve indicates the profile of $|U_2|$ for $c=6$, $d=2$ and $\alpha=1$; the green curve gives the profile of $|U_1|$ for $c=2$, $d=6$ and $\alpha=1$;

The curves of the Figure1 show the profiles of single solitary waves and the curves of the Figure 2 show the profiles of the combined solitary waves. In the two cases the red curves indicate the bright solitary waves and the green curves indicate the dark solitary waves.

Thus, the combined solutions U_1 and U_2 can be bright or dark type according to the choice of the coefficients a, b, c and d. When $a \succ b$, U_1 is a bright soliton. It becomes a dark soliton when $a \prec b$. For U_2, we have the same findings depending on $c \succ d$ or $c \prec d$. The above diagrams illustrate these different analyses. So in a general manner when we have a combined solitary wave formed of an association of two solitary waves of the type pulse and kink, the profile of the solitary wave gotten is either the one of a pulse or a dark. All depends on the choice of the parameters. In some cases the gotten profiles can stretch toward a bright or toward a dark without take the complete shape of the bright or the dark. In the setting of this survey, what interests us is the case where the combined solitary wave is formed of an association of a pulse and a kink and why? Because the sum of the solitary waves of the same nature gives a resultant solitary wave of this considered nature.

These types of solutions as we want to look for can have numerous applications in physics. It is besides this principle of analysis that motivates the construction of the solutions as we propose in this work. We organize the manuscript in the following manner:

Before constructing some solutions under the shape proposed in Eq. (2), we look for in section 2 of the possible solutions of the shape $\left(a\sinh^m(\alpha t)/\cosh^n(\alpha t)\right)\exp\left[-i(kz-\omega t)\right]$. In section 3, we construct the solutions of the type given by Eq.(3). Section 4, proposes the solutions of the type given by Eq. (4). Finally, section 5 concludes the work.

2. METHOD OF RESOLUTION

The principle consist of searching globally for the solutions of Eq.(1) in the shape

$$v=\sum_{i,j} a_{ij}\left(\cosh(\alpha x)\right)^j/\left(\sinh(\alpha x)\right)^i, \tag{5}$$

where $i=0,1,\ldots;\ j=0,1,\ldots$; α and a_{ij} are the coefficients to determine. When we introduce the ansatz (5) in Eq.(1) we obtain with the help of adequate transformations [7, 8], an equation of the form

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr

$$\begin{aligned}&\sum_{i,j,n} F\left(a_{ij}\right)/\cosh^{n}\left(\alpha x\right)+\sum_{i,j,m} G\left(a_{ij}\right)\sinh\left(\alpha x\right)/\cosh^{m}\left(\alpha x\right)\\&+\sum_{i,j,k} H\left(a_{ij}\right)\cosh^{k}\left(\alpha x\right)+\sum_{i,j,l} T\left(a_{ij}\right)\cosh^{l}\left(\alpha x\right)\sinh\left(\alpha x\right)+\sum_{i,j} W\left(a_{ij}\right)=0\end{aligned}. \tag{6}$$

$F\left(a_{ij}\right)$, $G\left(a_{ij}\right)$, $H\left(a_{ij}\right)$, $T\left(a_{ij}\right)$ and $W\left(a_{ij}\right)$ are linear functions of the coefficients a_{ij}. From Eq.(6), we obtain the series of equations of constants a_{ij} to solve. Notably the equations such as follows:

Term in $1/\cosh^{n}\left(\alpha x\right)$,

$$\sum_{i,j} F\left(a_{ij}\right)=0, \tag{7}$$

Term in $\sinh\left(\alpha x\right)/\cosh^{m}\left(\alpha x\right)$,

$$\sum_{i,j} G\left(a_{ij}\right)=0, \tag{8}$$

Term in $\cosh^{k}\left(\alpha x\right)$,

$$\sum_{i,j} H\left(a_{ij}\right)=0, \tag{9}$$

Term in $\cosh^{l}\left(\alpha x\right)\sinh\left(\alpha x\right)$,

$$\sum_{i,j} T\left(a_{ij}\right)=0, \tag{10}$$

Term in $\left(\sinh\left(\alpha x\right)\right)^{0}/\left(\cosh\left(\alpha x\right)\right)^{0}$,

$$\sum_{i,j} W\left(a_{ij}\right)=0. \tag{11}$$

Finally, the resolution of the above equations (7), …,(11) permits to obtain the constants a_{ij}.

3. RESULTS AND DISCUSSION

3.1 Single solitary wave solution

In this section we look for the single solitary waves that are the solutions of the Eq.(1). Thus, we suppose that

$$U\left(z,t\right)=a\frac{\sinh^{m}\left(\alpha t\right)}{\cosh^{n}\left(\alpha t\right)}\exp\left[-i\left(kz-\omega t\right)\right], \tag{12}$$

is the solution of Eq.(1) where a, α and k are constants to be determined, n and m are the whole numbers to be determined and $i^{2}=-1$. Then inserting Eq.(12) in Eq.(1) yields

$$\sum_{n,m} F_{nm}\left(a,k,\alpha,n,m\right)\frac{\sinh^{m}\left(\alpha t\right)}{\cosh^{n}\left(\alpha t\right)}=0, \tag{13}$$

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr

where $F_{nm}(a,k,\alpha,n,m)$ are the function of the constants a, k, α, n and m. The non trivial solutions are only gotten for $m=0$ and $n=1$. Thus, the coefficients of the terms in $1/\cosh^3(\alpha t)$, $1/\cosh(\alpha t)$, $\sinh(\alpha t)/\cosh^4(\alpha t)$ and $\sinh(\alpha t)/\cosh^2(\alpha t)$ give respectively

$$\alpha^2(2\omega\alpha_3-2\alpha_1)-(\alpha_2+\omega\alpha_4)|a|^2=0, \tag{14}$$

$$i\left[-k+\alpha_1(1+\omega^2)\alpha^2-2\alpha^2\omega\alpha_3\right]-\alpha_3\omega^2=0, \tag{15}$$

$$6\alpha_3\alpha+(3\alpha_4+2\alpha_5)|a|^2=0, \tag{16}$$

and

$$2\alpha_1\omega+(2\alpha\omega^2+1+\omega^2)\alpha^2\alpha_3=0 \tag{17}$$

The resolution of equations (14) and (16) gives

$$\alpha=\frac{-6(\alpha_3\alpha_2+\omega\alpha_3\alpha_4)}{6\omega\alpha_3\alpha_4-4\omega\alpha_3\alpha_5-6\alpha_1\alpha_4-4\alpha_1\alpha_5}, \tag{18}$$

and

$$|a|=\frac{\alpha^2(2\omega\alpha_3-2\alpha_1)}{\alpha_2+\omega\alpha_4}, \tag{19}$$

with $6\omega\alpha_3\alpha_4-4\omega\alpha_3\alpha_5-6\alpha_1\alpha_4-4\alpha_1\alpha_5\neq 0$ and $\alpha_2+\omega\alpha_4\neq 0$. On the other hand Eq.(15) gives

$$k=\alpha_1(1+\omega^2)\alpha^2-2\alpha^2\omega\alpha_3+i\alpha_3\omega^2, \tag{20}$$

where α is given by Eq.(18). Similarly Eq.(17) is written

$$(2\alpha^3+\alpha^2\alpha_3)\omega^2-2\alpha_1\omega+\alpha^2\alpha_3=0. \tag{21}$$

In the case where the condition $\alpha_1^2-\alpha^2\alpha_3(2\alpha^3+\alpha^2\alpha_3)\geq 0$ is verified, we obtain from Eq.(21)

$$\omega=\frac{\alpha_1\pm\sqrt{\alpha_1^2-\alpha^2\alpha_3(2\alpha^3+\alpha^2\alpha_3)}}{2\alpha^3+\alpha^2\alpha_3}. \tag{22}$$

Taking into account the equations (18), (19), (20) and (22) in Eq.(12), we obtain the solution

$$U(z,t)=\pm\alpha\sqrt{\frac{2\omega\alpha_3-2\alpha_1}{\alpha_2+\omega\alpha_4}}\,\mathrm{sec}\,h(\alpha t)\exp(\alpha_3\omega^2 z)\exp\left[-i(\Omega z-\omega t)\right], \tag{23}$$

with $\Omega=\alpha_1(1+\omega^2)\alpha^2-2\alpha^2\omega\alpha_3$ such that α and ω are given respectively by equations (18) and (22).

3.2 Solution of the form

$$U_1(z,t)=\left(a/\cosh(\alpha t)+b\sinh(\alpha t)/\cosh(\alpha t)\right)\exp-i(kz-\omega t)$$

The substitution of Eq.(3) in Eq.(1) leads to

$$\sum_{q,p}\left(H_{qp}(a,b)+iG_{qp}(a,b)\right)\frac{\sinh^p(\alpha t)}{\cosh^q(\alpha t)}=0,\ q=1,2,\ldots,5;\quad p=0,1 \tag{24}$$

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr

where $H_{qp}(a,b)$ and $G_{qp}(a,b)$ are the functions of a and b, $i^2=-1$. From Eq.(24), we obtain the series of equations of constants a and b

$$\sum_{q,p} H_{qp}(a,b)=0\,, \tag{25}$$

and

$$\sum_{q,p} G_{qp}(a,b)=0\,. \tag{26}$$

Some particular solutions of a and b can be obtain by summing equations (25) and (26). Then Eq.(25) and Eq.(26) lead to

$$\sum_{q,p}\left(H_{qp}(a,b)+G_{qp}(a,b)\right)=0\,. \tag{27}$$

Thus, from Eq.(27), we have

Term in $1/\cosh^4(\alpha t)$,

$$a^2\alpha(\alpha_5+9\alpha_4)-\alpha(\alpha_5+3\alpha_4)b^2+3\alpha_2 ab=0\,, \tag{28}$$

Term in $1/\cosh^3(\alpha t)$,

$$b^2-a^2=-\frac{\alpha^2}{\alpha_2}(\alpha_3\omega+2\alpha_1)\,, \tag{29}$$

Term in $1/\cosh^2(\alpha t)$,

$$\alpha_4\alpha\left(2b^2-6a^2\right)-\left(3\alpha_2\omega^2+2\alpha_1\omega\right)\alpha-2\alpha_2 a=0\,, \tag{30}$$

Term in $1/\cosh(\alpha t)$,

$$\alpha_5\alpha\left(4a^2b-2b^3\right)-\alpha_2 ab^2+\left(a\omega^2-3\alpha_3\omega\alpha^2+\alpha_1\omega^2-\alpha_1\alpha^2-k\right)a=0\,. \tag{31}$$

Another equations which derive from the term in $\sinh\alpha t/\cosh^l\alpha t$, $l=1,2,\ldots,5$ can allow to establish the constraint relations between the parameters of the physics system governs by Eq.(1). Combining equations (29) and (30), we obtain

$$4\alpha_2\alpha_4\alpha a^2+2\alpha_2^2 a+2\alpha_4\alpha^3+\alpha_2\left(3\alpha_2\omega^2+2\alpha_1\omega\right)\alpha=0\,. \tag{32}$$

Equation (32) is a quadratic equation in a. It resolution gives

$$a=\frac{-\alpha_2^2\pm\sqrt{\Delta'}}{4\alpha_2\alpha_4}\,, \tag{33}$$

with $\Delta'=\alpha_2^4-4\alpha_2\alpha_4\alpha^2\left(2\alpha_4\alpha^2+3\alpha_2^2\omega^2+2\alpha_1\alpha_2\omega\right)$ such that $\Delta'\geq 0$. Inserting Eq.(33) in Eq.(29), we obtain

$$b=\pm\sqrt{\frac{1}{4\alpha_2\alpha_4}\left[\left(-\alpha_2^2\pm\sqrt{\Delta'}\right)^2-16\alpha_2\alpha_4^2\left(\alpha_3\omega+2\alpha_1\right)\right]}\,. \tag{34}$$

Taking into account Eq.(33) and Eq.(34) into Eq.(3), we obtain

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr

$$U_1(z,t)=\left[\begin{array}{l}\left(\dfrac{-\alpha_2^2\pm\sqrt{\Delta'}}{4\alpha_2\alpha_4}\right)\sec h(\alpha t)\\ \pm\sqrt{\dfrac{1}{4\alpha_2\alpha_4}\left[\left(-\alpha_2^2\pm\sqrt{\Delta'}\right)^2-16\alpha_2\alpha_4^2(\alpha_3\omega+2\alpha_1)\right]}\tanh(\alpha t)\end{array}\right]\exp\left[-i(kz-\omega t)\right]. \quad (35)$$

3.1 Solution of the form

$$U_2(z,t)=\left(c/\cosh^2(\alpha t)+d\sinh(\alpha t)/\cosh^2(\alpha t)\right)\exp\left[-i(kz-\omega t)\right]$$

Substituting ansatz (4) in Eq.(1), we obtain the range equation

$$\sum_{\mu,\gamma}R_{\mu\gamma}(a,b,\alpha,\omega,k)\frac{\sinh^{\gamma}(\alpha t)}{\cosh^{\mu}(\alpha t)}+i\sum_{\mu,\gamma}T_{\mu\gamma}(a,b,\alpha,\omega,k)\frac{\sinh^{\gamma}(\alpha t)}{\cosh^{\mu}(\alpha t)}=0, \quad (36)$$

where $R_{\mu\gamma}(a,b,\alpha,\omega,k)$ and $T_{\mu\gamma}(a,b,\alpha,\omega,k)$ are functions of constants to determine; $i^2=-1$.

The real part and the imaginary part of Eq.(36) lead to the set of equations according to the Bogning-Djeumen Tchaho-Kofané method (BDKm) [7, 8]

$$\sum_{\mu,\gamma}R_{\mu,\gamma}(a,b,\alpha,\omega,k)=0, \quad (37)$$

and

$$\sum_{\mu,\gamma}T_{\mu\gamma}(a,b,\alpha,\omega,k)=0. \quad (38)$$

The combination of equations (37) and (38) which derive from the same terms in $\sinh^{\gamma}(\alpha t)/\cosh^{\mu}(\alpha t)$ ($\gamma=0$) leads to the following equations:

Term in $1/\cosh^7(\alpha t)$,

$$4\alpha\left(c^2d-d^3\right)(\alpha_4+\alpha_5)=0, \quad (39)$$

Term in $1/\cosh^6(\alpha t)$,

$$8c^2d\alpha(\alpha_4+\alpha_5)+c\alpha_2\left(c^2-3d^2\right)=0, \quad (40)$$

Term in $1/\cosh^5(\alpha t)$,

$$2(\alpha_4+\alpha\alpha_5)c^2-(2\alpha_4+\alpha\alpha_4+3\alpha\alpha_5)d^2+12\alpha_1\alpha^2=0, \quad (41)$$

Term in $1/\cosh^4(\alpha t)$,

$$2\alpha^2c^2d(\alpha_4+\alpha_5)-6\alpha_1\alpha^2c+3\alpha_2d^2c+6\alpha_3\alpha^2\omega c=0, \quad (42)$$

Term in $1/\cosh^3(\alpha t)$,

$$4\alpha_1\alpha\omega d-\alpha_3\left(20\alpha^3d-6\alpha\omega^2d\right)+2\alpha(\alpha_4+\alpha_5)d^3=0, \quad (43)$$

Term in $1/\cosh(\alpha t)$,

$$\alpha_1\omega\alpha d-\alpha_3\left(-\alpha^3+2\alpha\omega^2\right)d=0. \quad (44)$$

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr

The resolution of Eq.(39) supposes three possibilities. The case where $d\left(c^2-d^2\right)=0$ and $\alpha_4+\alpha_5=0$, the case where $d\left(c^2-d^2\right)=0$ and $\alpha_4+\alpha_5\neq 0$ and the case where $d\left(c^2-d^2\right)\neq 0$ and $\alpha_4+\alpha_5=0$. It is important to mention that we are looking for the non trivial solutions of Eq.(1). So a and b must be different from zero. Only the case $d\neq 0$, $c\neq d$ and $\alpha_4+\alpha_5=0$ permits to obtain the non trivial solutions. Thus, taking into account Eq.(39) into Eq.(40) yields

$$c^2=3d^2. \tag{45}$$

Introducing Eq.(45) in Eq.(41) gives

$$d^2=\frac{12\alpha_1\alpha^2}{\alpha\alpha_4-4\alpha_4-3\alpha\alpha_5}, \tag{46}$$

and

$$c^2=\frac{36\alpha_1\alpha^2}{\alpha\alpha_4-4\alpha_4-3\alpha\alpha_5}, \tag{47}$$

with $\alpha\alpha_4-4\alpha_4-3\alpha\alpha_5\neq 0$. Taking into account equations (46) and (47) into equations (41) and (42) allows to obtain α as a function of $\alpha_i\left(i=1,2,...,5\right)$ and consequently the relation which can help to determine ω as a function of a_i such as

$$\begin{aligned}&20\alpha_3\left(24\alpha_3\alpha_4\omega-24\alpha_1\alpha_4-108\alpha_1\right)^2\\&=\left(4\alpha_1\omega+6\alpha_3\omega^2\right)\left(6\alpha_1\alpha_4-18\alpha_1\alpha_5-6\alpha_3\alpha_4\omega+18\alpha_3\alpha_4\omega\right)^2.\end{aligned} \tag{41}$$

On the other hand introducing the obtained expressions of α and ω in other equations as (37 and (38)), we obtain the expression of k as a function of α_i. Then taking into account equations (39) and (40) in Eq.(4), we obtain

$$U_2\left(z,t\right)=\begin{bmatrix}\pm 6\alpha\sqrt{\dfrac{\alpha_1}{\alpha\alpha_4-4\alpha_4-3\alpha\alpha_5}}\sec h^2\left(\alpha t\right)\\ \pm\alpha\sqrt{\dfrac{12\alpha_1}{\alpha\alpha_4-4\alpha_4-3\alpha\alpha_5}}\sinh\alpha t\sec h^2\left(\alpha t\right)\end{bmatrix}\exp\left[-i\left(kz-\omega t\right)\right]. \tag{42}$$

4. CONCLUSION

The aim of this work was to construct some forced solutions of type lone wave. For this reason we carried our choice on the combined solitary wave solutions that means a solitary wave that result from the association of a solitary wave of type pulse and kink. Otherwise to say a solitary wave which can take according to the choice of the parameters, a shape pulse or kink. In the setting of BDKm used in this work, when one has several equations of the constants, one limits itself to the resolution of the equations of the first range susceptible to produce the solutions that come closer to best of the exact solution. In case of the solution given by the equation (46), we concentrated our

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr

analysis to the range of equations in $1/\cosh^l \alpha t$ with $l = 1, 2, ..., 7$. What spur our interest in the construction of such solutions is that these solutions will have important applications in physics or in engineering of telecommunication. The survey made in this manuscript can be spread to the cases where the solution combined solitary wave is an association of three solitary waves, four and more.

ACKNOWLEDGEMENTS

I acknowledge support from the ministry of Higher Education of Cameroon through its program of support to the research, which enabled us to carry out this work. I also thank Mr Kenfack Christopher Nguimatio for some corrections after the reading of the work.

REFERENCES

[1] Fan , E. (200). Extended tanh-function method and its applications to nonlinear equations. Physics Letter A 277, 212-218.

[2] E. J. Parkes , E. J. and Duffy, B. R. (1996). An automated tanh-function method for finding solitary wave solutions to nonlinear evolution equations. Comput. Phys. Commun, 98 288-300.

[3] A. H. Khater, A. H., Malfiet, W., Callebaut D. K. and Kamel E. S. (2002). The tanh method, a simple transformation and exact analytical solutions for nonlinear reaction-diffusion equations. Chaos, Solitons & Fractals, 14, 513-522.

[4] Kaup , D. J. and Lakoba, T. I. (1996). Variational method: How it can generate false instabilities, J. Math. Phys, 37 (7), 3442-3462.

[5] A. M. Wazwaz, A. M. (2002). The tanh method for traveling wave solutions of nonlinear equations, Appl. Math. Comput, 133, 213-227.

[6] Wazwaz, A. M. (2001). A computational approach to soliton solutions of the Kadomtsev-Petviashvili equation, Appl. Math. Comput. 123, 205-217.

[7] Bogning,J. R., Djeumen Tchaho , C. T. and Kofané, T. C. (2012) Construction of the soliton solutions of the Ginzburg-Landau equations by the new Bogning-Djeumen Tchaho-Kofané method, Phys. Scr, 85, 025013-025018.

[8] Bogning, J. R., Djeumen Tchaho, C. T. and Kofané, T. C. (2012). Generalization of the Bogning-Djeumen Tchaho-Kofané method for the construction of the solitary waves and the survey of the instabilities, Far East J. Dyn. Sys, **20**(2), 101-119.

[9] Djeumen Tchaho, C. T., Bogning, J. R. and Kofané, T. C. (2012). Modulated Soliton solution of the Modified Kuramoto-Sivashinsky's equation. American Journal of Computational and Applied Mathematics, 2(5), 218- 224.

[10] Djeumen Tchaho, C. T., Bogning , J. R. and Kofane, T. C. (2011). Multi-Soliton solutions of the modified Kuramoto-Sivashinsky's equation by the BDK method, Far East J. Dyn. Sys, 15(2), 83-98.

[11] Bogning, J. R. and Kofane, T. C. (2006). Analytical Solutions of discrete nonlinear Schrödinger equation in arrays of optical fibers, Chaos, Solitons &Fractals, 28, 148-153.

[12] Bogning, J. R. (2013). Pulse soliton Solutions of the modified KdV and Born- Infeld Equations, International Journal of Modern Nonlinear Theory and Applications, 2, 135-140.

[13] Kamruzzaman Khan and Ali Akbar, M. (2013). Exact solutions of the nonlinear generalized shallow water wave equation. World Applied Sciences Journal, 27(12), 1581-1587

[14] Kamruzzaman Khan, Ali Akbar, M. and Harun-Or-Roshid, (2014). Exact traveling wave solutions of nonlinear evolution equation via enhanced (G'/G)-expansion method, British Journal of Mathemetics and Computer Science, 4(10), 1318-1334.

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr

[15]Kamruzzaman Khan and Ali Akbar, M. (2013). Application of exp($\phi(\xi)$)-expansion method to find the exact solutions of modified Benjamin-Bona-Mahony equation. World Applied Sciences Journal. 24(10), 1373-1377.

[16] Zakharenko, A. A. (2010). Slow acoustic waves with the anti-plane polarization in layered systems. International Journal of Modern Physics B. 24(5), 515-536.

[17] Bogning, J. R., Djeumen Tchaho, C. T. and Kofané, T. C. (2013). Solitary wave solutions of the modified Sasa- Satsuma nonlinear partial differential equation, American Journal of Computational and Applied Mathematics, 3(2), 97-107.

[18] Bogning, J. R. (2013). Analytical soliton Solutions and wave solutions of discrete Nonlinear Cubic-quintique Ginzburg-Landau equations in array of dissipative optical systems. American Journal of Computational and Applied Mathematics, 3(2), 97-105.

[19] Hirota , R. (1973). Exact N-Soliton Solution of Nonlinear Self-Dual Network Equations, J. Math. Phys.14, 805-809.

[20] Ablowitz , M. J. and larkson , P. A. (1991). Soliton, nonlinear evolution equations and inverse scattering, New York Cambridge University Press.

[21] Guch et al. (1990). Soliton theory and its application. Zhejiang: Zhejiang Science and Technology press.

[22] Wadati, M. (1975). Wave propagation in nonlinear lattice I. J. Phys. Soc. JPn. 38, 673-680.

[23] Wadati, M. (1975). Wave propagation in nonlinear lattice II. J. Phys. Soc. JPn. 38, 681-686.

[24] Yajima, T. and Wadati, M. (1990). Solution and its property of unstable nonlinear Schrödinger equation. J. Phys. Soc. JPn.59, 41- 47.

[25] Shuqing Li , Lu Li, Zhonghao Li and Guosheng Zhou. (2004). Properties of soliton solutions on the cw background in optical fibers with higher-order effects. J. Opt. Soc. Am. B, 21(12), 2089-2094.

[26] Nakkeeran, K. (2002). Bright and dark optical solitons in fiber media with higher-order effects. Chaos Solitons & Fractals, 13, 673-679.

E-mail address: rbogning@yahoo.com / jrbogning@yahoo.fr