

*Method article***On Finding Geodesic Equation of Two Parameters
Inverse Gaussian Distribution****ABSTRACT**

The class of Inverse Gaussian distributions is quite commonly used as a life-time model in reliability studies. The books by Chhikara and Folks [1] and Seshadri[2] present extensive discussions on classical inference for the parameters of Inverse Gaussian distribution. However, in this paper, we switch our attention to find its geodesic equation. We applied two different algorithms to solve some partial differential equations, where these equations originated from the Inverse Gaussian distribution. As expected , the two algorithms yield the same result.

MATHEMATICAL SUBJECT CLASSIFICATION : 62E99

KEY WORDS AND PHASES: Darboux Theory, Differential Geometry, Geodesic Equation, Inverse Gaussian Distribution, Triply Partial Differential Equation,

INTRODUCTION

The inverse Gaussian distribution is particularly important to probabilists and physicists due to its relation to

Brownian motion. Balakrishnan N. and Chen W.W.S.[3] have completed the tremendous task of computing the means, variances and covariances of order statistics for all sample sizes up to twenty five and for many choices of the shape parameter. The major reason for taking up this study is to make use of the tabulated values of means, variances and covariances of order statistics in order to derive the best linear unbiased estimators of the location and scale parameters based on complete as well as Type-II censored samples. Aside from the theoretical issues, the distribution has much to offer to the practicing statistician in many important areas of application. For example, the pioneering work of Tweedie[4] used the inverse Gaussian in a clinical trial study of the effect of a drug on the first passage time taken for a jejunal biopsy capsule, on leaving the stomach, to travel from the pylorus through the duodenum and into the jejunum. Working in collaboration with statisticians in the Clinical Cancer Research Institute in Liverpool, Tweedie studied the distribution of survival times in a series of patients who had been treated for cancer, and fitted the inverse Gaussian with considerable success. He also found that the antilognormal and Weibull were poor fits. In this paper, instead of studying all previously stated topics of interest, we switch to a new direction, by finding the geodesic equation of an inverse Gaussian distribution. We used two different algorithms to approach this purpose, and found that both algorithms reach the same result. In section 2, we list the fundamental tensor we need for later use. In section 3, we present two algorithms to show the process of deriving the required geodesic equation. In section 4, we repeat some more detail procedures on how the fundamental tensor has been derived. Six Christoffel symbols have been collected in the appendix. There are many related books

published in this area of study, including Kass R.E. and Vos P.W.[5]and Amari S-I[6].

2. LIST THE FUNDAMENTAL TENSOR

The standard or canonical form of the two-parameter Inverse Gaussian distribution has the probability density function

given by,

$$f(y,u,v) = \left(\frac{v}{2\pi y^3}\right)^{\frac{1}{2}} \exp\left(-\frac{v}{2u^2 y}(y-u)^2\right) \quad y > 0$$

$$\ln f(y,u,v) = \frac{1}{2}(\ln v - \ln(2\pi y^3)) - \frac{v}{2u^2 y}(y-u)^2$$

It is known that, in this form of the distribution, u is the mean and $\frac{u^3}{v}$ is the variance, Furthermore, $\frac{u}{v}$ is the square of the coefficient of variation. From the equation above, we derive the metric tensor components for this distribution as follows,

$$E = -E\left(\frac{\partial^2 \ln f(x)}{\partial u^2}\right) = \frac{v}{u^3}, \quad F = -E\left(\frac{\partial^2 \ln f(x)}{\partial v \partial u}\right) = 0,$$

$$G = -E\left(\frac{\partial^2 \ln f(x)}{\partial v^2}\right) = \frac{1}{2v^2}$$

Using the results above, we can easily derive the required basic tensor metric and Christoffel symbols as follows:

$$E_u = \frac{-3v}{u^4}, \quad E_v = \frac{1}{u^3}, \quad F_u = 0, \quad F_v = 0, \quad G_u = 0,$$

$$G_v = \frac{-1}{v^3}, \quad EG = \frac{1}{2u^3 v}$$

$$\begin{aligned} \Gamma_{11}^1 &= \frac{E_u}{2E} = \frac{-3}{2u}, \quad \Gamma_{12}^2 = \frac{G_u}{2G} = 0, \quad \Gamma_{11}^2 = \frac{-E_v}{2G} = \frac{-v^2}{u^3}, \\ \Gamma_{22}^1 &= \frac{-G_u}{2E} = 0, \quad \Gamma_{12}^1 = \frac{E_v}{2E} = \frac{1}{2v}, \quad \Gamma_{22}^2 = \frac{G_v}{2G} = \frac{-1}{v} \end{aligned}$$

3. The Geodesic Equation

One method to find the geodesic equation of the Inverse Gaussian distribution is by solving a triply of partial differential equations given in the appendix. (see Struik, D.J. or Grey, A [7][8]) We seek its solution as below. To avoid confusion, we will only index those formulas that we will use later, and we will ignore the others:

$$\frac{d^2u}{ds^2} - \frac{3}{2u} \left(\frac{du}{ds} \right)^2 + \frac{1}{v} \frac{dudv}{ds} = 0, \quad (1)$$

$$\frac{d^2v}{ds^2} - \frac{v^2}{u^3} \left(\frac{du}{ds} \right)^2 - \frac{1}{v} \left(\frac{dv}{ds} \right)^2 = 0, \quad (2)$$

And the first fundamental form of distance function is given by

$$\begin{aligned} ds^2 &= Edu^2 + 2Fdudv + Gdv^2 \\ &= \frac{v}{u^3} du^2 + \frac{1}{2v^2} dv^2 \end{aligned} \quad (3)$$

It needs only two out of the three equations above to find the geodesic equation. We will choose equations (1) and (3). To simplify the notation, we let

$$\begin{aligned} \frac{du}{ds} &= p, \quad \frac{d^2u}{ds^2} = \frac{dp}{ds} \quad \text{then} \\ \frac{dp}{ds} - \frac{3}{2u} p^2 + \frac{p}{v} \frac{dv}{ds} &= 0. \end{aligned} \quad (4)$$

100 Dividing the equation (4) by p , and integrating on both sides with
101 respect to p , we get

$$\begin{aligned} \frac{dp}{p} - \frac{3}{2u}p + \frac{1}{v} \frac{dv}{ds} &= 0, \quad \ln p - \frac{3}{2} \ln u + \ln v = C \\ \text{or } \ln p &= C + \frac{3}{2} \ln u - \ln v = \ln A + \ln \frac{u^{3/2}}{v} = \ln \frac{Au^{3/2}}{v} \end{aligned}$$

103 Where we choose $C = \ln A$

$$104 \quad p = \frac{Au^{3/2}}{v} = \frac{du}{ds} \quad \text{or} \quad Ads = \frac{vdu}{u^{3/2}}$$

105 Square both side, we get

$$106 \quad A^2 ds^2 = \frac{v^2 du^2}{u^3} \quad (5)$$

107 From equation (3), we see that the first fundamental form of Inverse
108 Gaussian Model is given by:

$$109 \quad ds^2 = \frac{v}{u^3} du^2 + \frac{1}{2v^2} dv^2$$

$$110 \quad \text{Hence, } A^2 ds^2 = \frac{A^2 v}{u^3} du^2 + \frac{A^2}{2v^2} dv^2 \quad (6)$$

111 Put (5) and (6) together we get

$$\begin{aligned} \frac{v^2}{u^3} du^2 &= \frac{A^2 v}{u^3} du^2 + \frac{A^2}{2v^2} dv^2 \\ \text{or } \frac{(v^2 - A^2 v)}{u^3} du^2 &= \frac{A^2}{2v^2} dv^2 \end{aligned} \quad (7)$$

113 By taking the square root of equation (7)

$$\begin{aligned} \pm \sqrt{\frac{(v - A^2)v}{u^3}} du &= \frac{\pm Adv}{\sqrt{2}v} \\ \text{or } \frac{\pm du}{u^{3/2}} &= \frac{\pm Adv}{\sqrt{2}v\sqrt{(v - A^2)v}} \end{aligned}$$

116 Integrating both sides, we get

$$\pm \int u^{-3/2} du \pm \frac{1}{\sqrt{2}} \int \frac{Adv}{v\sqrt{(v-A^2)v}} = B$$

$$\text{or } \pm \frac{2}{\sqrt{u}} \pm \frac{1}{\sqrt{2}} \int \frac{Adv}{v^{3/2}\sqrt{v-A^2}} = B$$

Where A, B are arbitrary constants.

Alternatively, we can find the geodesic equation of the Inverse

Gaussian distribution by solving one partial differential equation.

This idea originated from French mathematician Darboux's theory. A

detailed proof has been given in Chen[9][10]. From section 2, we know

that the coefficient of the first fundamental form of $\nabla Z=1$ is given

$$E = -E\left(\frac{\partial^2 \ln f(x)}{\partial u^2}\right) = \frac{v}{u^3}, \quad F = -E\left(\frac{\partial^2 \ln f(x)}{\partial v \partial u}\right) = 0, \quad G = -E\left(\frac{\partial^2 \ln f(x)}{\partial v^2}\right) = \frac{1}{2v^2}$$

by,

$$\Delta z = 1; \quad \frac{v}{u^3} z_v^2 + \frac{1}{2v^2} z_u^2 = \frac{1}{2u^3 v}$$

To solve the partial differential equation above, we may use the

separable variable method as follows:

By multiple above equation $2u^3v$, we derived

$$2v^2 z_v^2 + \frac{u^3}{v} z_u^2 = 1$$

$$\text{or } u^3 z_u^2 = v(1 - 2v^2 z_v^2) = A^2,$$

Break this into two parts, we get

$$\text{Part 1, } u^3 z_u^2 = A^2 \text{ implies } z_u^2 = \frac{A^2}{u^3}$$

$$\text{or } z_u = \pm \frac{A}{u^{3/2}} = Au^{-3/2} \text{ or } Z = \pm 2Au^{-1/2} = \frac{\pm 2A}{\sqrt{u}} \quad (8)$$

$$\text{and part 2, } v(1 - 2v^2 z_v^2) = A^2$$

$$\text{or } v - A^2 = 2v^3 z_v^2 \text{ so } z_v^2 = \frac{v - A^2}{2v^3}$$

$$z = \pm \frac{1}{\sqrt{2}} \int \frac{\sqrt{v-A^2}}{v^{3/2}} dv \quad (9)$$

The general solution of the geodesic equation combined (8) and (9), hence we get

$$z = \frac{\pm 2A}{\sqrt{u}} \pm \frac{1}{\sqrt{2}} \int \frac{\sqrt{v-A^2}}{v^{3/2}} dv$$

Applying the Darboux Theory, we finally find that the geodesic equation of Inverse Gaussian Distribution is given by

$$\frac{\partial z}{\partial A} = B \text{ or } \pm \frac{2}{\sqrt{u}} \pm \frac{1}{\sqrt{2}} \int \frac{Adv}{v^{3/2} \sqrt{v-A^2}} = B$$

Where A and B are arbitrary constants. This result is the same as the previous one.

4. Deriving The Basic Tensor

The probability density function of two parameters of the Inverse Gaussian distribution is given by

$$f(y, u, v) = \left(\frac{v}{2\pi y^3}\right)^{\frac{1}{2}} \exp\left(-\frac{v}{2u^2 y}(y-u)^2\right) \quad y > 0$$

$$\ln f(y, u, v) = \frac{1}{2} (\ln v - \ln (2\pi y^3)) - \frac{v}{2u^2 y} (y-u)^2$$

From the equation above, we derive the metric tensor components for the distribution as follows,

$$\frac{\partial \ln f}{\partial u} = -\frac{v}{2y} \frac{\partial}{\partial u} \left(\frac{y-u}{u}\right)^2 = \frac{v(y-u)}{u^3}$$

$$\frac{\partial^2 \ln f}{\partial u^2} = \frac{v(-3y+2u)}{u^4},$$

$$\frac{\partial^2 \ln f}{\partial v \partial u} = \frac{y-u}{u^3},$$

$$\frac{\partial \ln f}{\partial v} = \frac{1}{2v} - \frac{(y-u)^2}{2u^2 y}, \quad \frac{\partial^2 \ln f}{\partial v^2} = \frac{-1}{2v^2}$$

For the next step, we need to find the expectation of these second order partial derivatives. The answer is almost straightforward, and we just list them below:

$$E = -E\left(\frac{\partial^2 \ln f}{\partial u^2}\right) = \frac{v}{u^3}, \quad F = -E\left(\frac{\partial^2 \ln f}{\partial v \partial u}\right) = 0, \quad G = -E\left(\frac{\partial^2 \ln f}{\partial v^2}\right) = \frac{1}{2v^2}$$

5. appendix

We list the six well known Christoffel Symbols as follows. For a detailed derivation, see Struik[4] or Grey[5].

$$\Gamma_{11}^1 = \frac{GE_u - 2FF_u + FE_v}{2(EG - F^2)}, \quad \Gamma_{12}^2 = \frac{EG_u - FE_v}{2(EG - F^2)}$$

$$\Gamma_{11}^2 = \frac{2EF_u - EE_v - FE_u}{2(EG - F^2)}, \quad \Gamma_{22}^1 = \frac{2GF_v - GG_u - FG_v}{2(EG - F^2)}$$

$$\Gamma_{12}^1 = \frac{GE_v - FG_u}{2(EG - F^2)}, \quad \Gamma_{22}^2 = \frac{EG_v - 2FF_v + FG_u}{2(EG - F^2)}$$

In general, the solution of the geodesic equation depends upon a pair of partial differential equations as seen below:

$$\frac{d^2 u}{ds^2} + \Gamma_{11}^1 \left(\frac{du}{ds}\right)^2 + 2\Gamma_{12}^1 \left(\frac{du}{ds} \frac{dv}{ds}\right) + \Gamma_{22}^1 \left(\frac{dv}{ds}\right)^2 = 0$$

$$\frac{d^2 v}{ds^2} + \Gamma_{11}^2 \left(\frac{du}{ds}\right)^2 + 2\Gamma_{12}^2 \left(\frac{du}{ds} \frac{dv}{ds}\right) + \Gamma_{22}^2 \left(\frac{dv}{ds}\right)^2 = 0$$

REFERENCE

- [1] Chhikara, R.S. and Folks, J.L. (1989). The Inverse Gaussian Distribution : Theory, Methodology and Applications.

- 169 Marcel Dekker, New York.
- 170 [2] Seshadri V. (1993) The Inverse Gaussian Distribution :
171 A Case Study in Exponential families. Clarendon Press,
172 Oxford, England.
- 173 [3] Balakrishnan N. and Chen W.W.S.(1997). CRC Handbook of
174 Order Statistics from Inverse Gaussian Distributions
175 with Applications. CRC Press, Boca Raton.
- 176 [4] Tweedie, M.C.K.(1945)Inverse statistical variates,
177 Nature, 155, 453.
- 178 [5] Kass R.E. and Vos P.W.(1997) Geometrical Foundations of
179 Asymptotic Inference. John Wiley & Sons, Inc.
- 180 [6] Amari, S-I(1990) Differential-Geometrical Methods in
181 Statistics, Springer, New-York.
- 182 [7] Struik, D.J.(1961) Lectures on Classical Differential
183 Geometry. Second Edition. Dover Publications, Inc.
- 184 [8] Grey, A.(1993) Modern Differential Geometry of Curves and
185 Surfaces. CRC Press, Inc. Boca Raton.
- 186 [9] Chen W.W.S. (2013) A Note On Finding Geodesic Equation
187 of two Parameters Gamma Distribution. Applied
188 Mathematics, www.scirp.org/journal/am ,5, 3511-3517.
- 189 [10] Chen W.W.S. (2014) A note on finding geodesic equation
190 of two parameter Weibull distribution. Theoretical
191 Mathematics & Applications, Vol.4, no. 3, 43-52.